



PII: S0301-0082(96)00055-X

# NEUROIMMUNOMODULATION VIA LIMBIC STRUCTURES — THE NEUROANATOMY OF PSYCHOIMMUNOLOGY

HELGA SUSANNE HAAS and KONRAD SCHAUENSTEIN\*

Department of General and Experimental Pathology, University of Graz Medical School, Mozarigasse 14, A-8010, Graz, Austria

(Received 19 July 1996)

**Abstract**—During the last 20 years, mutual communications between the immune, the endocrine and the nervous systems have been defined on the basis of physiological, cellular, and molecular data. Nevertheless, a major problem in the new discipline “Psychoneuroimmunology” is that controversial data and differences in the interpretation of the results make it difficult to obtain a comprehensive overview of the implications of immunoneuroendocrine interactions in the maintenance of physiological homeostasis, as well as in the initiation and the course of pathological conditions within these systems. In this article, we will first discuss the afferent pathways by which immune cells may affect CNS functions and, conversely, how neural tissues can influence the peripheral immune response. We will then review recent data, which emphasize the (patho)physiological roles of hippocampal–amygdala structures and the nucleus accumbens in neuroimmunomodulation. Neuronal activity within the hippocampal formation, the amygdaloid body, and the ventral parts of the basal ganglia has been examined most thoroughly in studies on neuroendocrine, autonomic and cognitive functions, or at the level of emotional and psychomotor behaviors. The interplay of these limbic structures with components of the immune system and vice versa, however, is still less defined. We will attempt to review and discuss this area of research taking into account recent evidences for neuroendocrine immunoregulation via limbic neuronal systems, as well as the influence of cytokines on synaptic transmission, neuronal growth and survival in these brain regions. Finally, the role of limbic structures in stress responses and conditioning of immune reactivity will be commented. Based on these data, we propose new directions of future research. © 1997 Elsevier Science Ltd. All Rights Reserved.

## CONTENTS

1. Introduction	196
2. Neuroimmune communications	196
2.1. Afferent pathways	199
2.2. Efferent pathways	201
3. The limbic system	202
4. Neuroimmune communications via limbic structures	205
4.1. Lesioning limbic areas affects immune responses	205
4.2. Neuroendocrine immunomodulation at limbic neuronal systems	205
4.3. Immune stimulation evokes electrophysiological changes in neuronal activities in limbic areas	207
4.4. Limbic structures in stress and immunity	209
4.5. Effects of the activated immune system on limbic structures	210
4.6. Conditioning of immune functions — mediated via limbic areas?	211
5. Conclusions	211
Acknowledgements	212
References	212

## ABBREVIATIONS

ACTH	Adrenocorticotropin	CNS	Central nervous system
AP	Area postrema	ConA	Concanavalin A
AVP	Arginine vasopressin	CRH/CRF	Corticotropin-releasing hormone/corticotropin-releasing factor
BBB	Blood–brain barrier	CVO	Circumventricular organ
BDNF	Brain-derived neurotrophic factor	DN cells	Double negative (CD4/CD8) cells (immature T-lymphocytes)
BLA	Basolateral amygdala	DP cells	Double positive (CD4/CD8) cells (immature T-lymphocytes)
BST	Bed nucleus of the stria terminalis	EPSP	Excitatory postsynaptic potential
CA	Cornu Ammonis (field of hippocampal formation)	F344 rat	Fischer 344 rat
CCK	Cholecystokinin	GABA	$\gamma$ -Aminobutyric acid
CeMA	Centromedial amygdala		
CGRP	Calcitonin gene-related peptide		

\* Author for correspondence. Fax: + 43 316 380 9640.

ABBREVIATIONS (*continued*)

GALT	Gut-associated lymphoid tissue	NK cells	Natural killer cells
GM-CSF	Granulocyte-macrophage colony-stimulating factor	NMDA	<i>N</i> -methyl-D-aspartic acid
GR	Glucocorticoid receptor	NO	Nitric oxide
HF	Hippocampal formation	NZB/NZW	New Zealand (black/white) mice
HIV	Human immunodeficiency virus	OB rat	Olfactory bulbectomized rat
HPA axis	Hypothalamic-pituitary-adrenal axis	OVL	Organum vasculosum laminae terminalis
5-HIAA	5-Hydroxyindole acetic acid	PACAP	Pituitary adenylate cyclase activating polypeptide
5-HT	5-Hydroxytryptamine (serotonin)	PAG	Periaqueductal grey
i.c.v.	Intracerebroventricular	PG	Prostaglandin
IFN	Interferon	PHA	Phytohemagglutinin
IL	Interleukin	PNS	Peripheral nervous system
IL-1R	Interleukin-1 receptor	PTP	Post-tetanic potentiation
IL-1RA	Interleukin-1 receptor antagonist	SD rat	Sprague Dawley rat
i.p.	Intraperitoneal	SFO	Subfornical organ
IPSP	Inhibitory postsynaptic potential	SI	Substantia innominata
i.v.	Intravenous	SP cells	Single positive (CD4/CD8) cells (mature T-lymphocytes)
LPS	Lipopolysaccharide	ST	Stria terminalis
LTP	Long-term potentiation	STP	Short-term potentiation
ME	Median eminence	TGF $\beta$	Transforming growth factor $\beta$
MR	Mineralocorticoid receptor	TNF	Tumor necrosis factor
mRNA	Messenger ribonucleic acid	VIP	Vasoactive intestinal polypeptide
NE	Norepinephrine		
NGF	Nerve growth factor		

## 1. INTRODUCTION

Over the past several years, our knowledge of reciprocal intersystem communications involving the immune, the endocrine, the peripheral (PNS) and central nervous systems (CNS) has exponentially grown. This field of science gains the attention of both basic scientists and clinicians to define new pathophysiological concepts in autoimmune (Schauenstein *et al.*, 1987; Sternberg *et al.*, 1989a, 1989b; Mason *et al.*, 1990; Takasu *et al.*, 1990, 1993; Hu *et al.*, 1993; Jafarian Tehrani *et al.*, 1994), allergic (Udem *et al.*, 1995) and infectious diseases (Blum *et al.*, 1992; Duvaux-Miret *et al.*, 1992), stress conditions (Khansari *et al.*, 1990; Hoffman-Goetz and Pedersen, 1994), as well as affective disorders (Irwin *et al.*, 1991; Evans *et al.*, 1992; Maes *et al.*, 1993), and dementia (Irwin *et al.*, 1991; Vandenabeele and Fiers, 1991). Much of our understanding of neuroimmunomodulation is derived from earlier studies documenting the effects of stress on immune functions (Dantzer and Kelley, 1989). The data are partly controversial, which may be due to species differences, to the fact that different qualities and/or doses of stressors may affect immune reactions in different ways (Rinner *et al.*, 1992), and to individual abilities to cope with one and the same stressor (Ader *et al.*, 1995). Besides immunoregulation by the CNS, it has become increasingly evident that the production of neuroendocrine peptides by immune cells is physiologically relevant to both the immune and the nervous system (summarized by Blalock, 1992; and by Blalock, 1994). However, our knowledge about the ascending immune-brain routes, and the function of brain regions, sometimes referred to as "higher brain centers", in neuroimmunological interactions is still fragmentary. This article, besides describing afferent routes by which immune cells may modify CNS functions and summarizing the most prominent efferent pathways involved in brain-immune communications, presents a brief introduction into the broad range of limbic effector functions, and

a detailed description of the specific roles of the hippocampal-amygdala group as well as the ventral striatopallidal system in neuroimmunomodulation (Fig. 1).

## 2. NEUROIMMUNE COMMUNICATIONS

The concept of a dialog between immune cells and neural tissues is based on three requirements:

1. Nerve terminals directly contact immune cells. The noradrenergic as well as the peptidergic innervation of lymphoid organs is best documented in rodents. Within the spleen at both light and electron microscopic levels, norepinephrine (NE)-nerve terminals have been observed in close apposition to lymphocytes (Felten and Olschowka, 1987). The authors described these contacts as an even more regular and intimate membrane relationship than between nerve terminals and smooth muscle cells, and defined these associations as "synaptic-like" contacts, where transmitter release and postsynaptic receptor activation most likely occur. Noradrenergic postganglionic sympathetic nerve fibers also have been detected in primary lymphoid organs, such as bone marrow and thymus (reviewed by Felten *et al.*, 1987), and neuropeptide-containing nerve fibers have been shown to distribute in lymphatic tissues among lymphocytes and mast cells (Bienenstock *et al.*, 1987; Stead *et al.*, 1987; Bellinger *et al.*, 1990). More recent evidence clearly shows an equivalent anatomical link between the nervous and immune system in the cetacean, *Delphinapterus leucas* (beluga whale). Tyrosine hydroxylase and neuropeptide Y positive nerve profiles were observed not only in systemic lymphoid organs, such as spleen, thymus, tonsil, or assorted lymph nodes, but also in gut-associated lymphoid tissue (GALT) (Romano *et al.*, 1994). As to the role of neurotrophic factors in the induction of sympathetic innervation of immune organs a recent *in vitro* study demonstrated that interleukins, such as IL-1.

IL-3 and IL-6, as well as granulocyte-macrophage colony-stimulating factor (GM-CSF), can induce sympathetic neurite outgrowth, either via nerve growth factor (NGF) production or in an NGF-independent manner (Kannan *et al.*, 1996). Although cholinergic signals markedly affect immune functions (Schauenstein *et al.*, 1994), the cholinergic innervation of immune organs is less described and, in the case of the thymus, is a point of controversy (Bullock and Moore, 1981; Fatani *et al.*, 1986; Nance *et al.*, 1987; Micic *et al.*, 1994). However, recent data by Nijima (1995) apparently confirmed the vagal innervation of the rat thymus by means of electrophysiological techniques.

2. Both organ systems share common receptors and ligands. In several studies, remarkable concentrations of binding sites for a number of neurotransmitters or -peptides have been identified on the surface of white blood cells (Gordon *et al.*, 1978; Ovadia and Abramsky, 1987; Hellstrand and Hermodsson, 1990a, 1990b; Lignon *et al.*, 1991; Carr, 1992; Landmann, 1992; Kaneda *et al.*, 1993; Costa *et al.*, 1994). Furthermore, immune cells are found to be equipped to produce not only cytokines, but also

neurohormones, neuropeptides (Akira *et al.*, 1990; Blalock, 1992, 1994) and neurotransmitters, such as catecholamines (Bergquist *et al.*, 1994) and, as reported by our group, acetylcholine (Rinner and Schauenstein, 1993). On the other hand, interleukin (IL)-1 $\beta$  as well as mRNA homologous to IL-1 receptor (IL-1R) type I are expressed in cultured sympathetic ganglia (Freidin *et al.*, 1992; Hart *et al.*, 1993) and IL-1 $\beta$  binding sites have been demonstrated on paraganglia in the rat liver hilus and hepatic vagus (Goehler *et al.*, 1994). IL-6 and IL-6R transcripts were found in postnatal rat sympathetic and sensory ganglia (Gadient and Otten, 1996). IL-6R mRNA has been detected in cultured primary rat sympathetic neurons as well as in the pheochromocytoma cell line PC12, and this expression can be modulated by tumor necrosis factor (TNF)- $\alpha$  (März *et al.*, 1996). Research on nematode infections in small mammals revealed an enhanced expression of IL-1 $\alpha$ , IL-1 $\beta$ , IL-6 and TNF- $\alpha$  mRNA and protein in the longitudinal muscle myenteric plexus of the inflamed intestine of rats (Khan and Collins, 1994). Additionally, a continuously growing number of reports shows that cytokines and their receptors are




Fig. 1. Limbic areas in neuroimmunomodulation.



cholinergic neuronal function (longitudinal muscle myenteric plexus preparations from the rat jejunum/whole preparations of guinea pig ileum, including mucosa) in a dual way, either suppressing or stimulating the electrically evoked release of acetylcholine (Main *et al.*, 1993; Mameya *et al.*, 1995). A similar duality was observed concerning the effects of IL-6 on the evoked release of NE from the rat jejunal myenteric plexus (Rühl *et al.*, 1994). Furthermore, IL-1 $\beta$  has been shown to mediate increases in substance P receptor mRNA in explanted neonatal sympathetic ganglia (Ludlam *et al.*, 1995), IL-1 $\beta$  induces substance P in cultured superior cervical ganglia (Jonakait and Schotland, 1990; Jonakait *et al.*, 1990; Hart *et al.*, 1991), and IL-1 $\alpha$ , as well as TNF- $\alpha$ , differentially modulate both substance P and vasoactive intestinal polypeptide (VIP) biosynthesis in adrenomedullary chromaffin cells (Eskay *et al.*, 1992). In addition, cytokines are known to mediate a wide variety of other effects, such as neuroendocrine responses, neuronal growth, autonomic functions, and behavioral activity (reviewed by Schöbitz *et al.*, 1994a).

### 2.1. Afferent Pathways

Although there is compelling evidence that the peripheral immune system modifies central neural cell functions, the afferent pathways by which immune cells affect the CNS are still far from clear. In the following, two principal mechanisms will be discussed (Fig. 3).

As one mechanism, a humoral route may be involved in neuroimmune communications. Some studies suggested an active transport of cytokines across the blood-brain barrier (BBB) (Gutierrez *et al.*, 1993, 1994; Maness *et al.*, 1995), others supposed an interaction at the level of endothelial cells, which subsequently results in the induction of central prostaglandins (Gottschall *et al.*, 1992a; Van Dam *et al.*, 1993, 1996). Recently it has been found that IL-1 $\beta$ , IL-6 and TNF- $\alpha$  induce changes in the electrical resistance across rat cerebral endothelial cells *in vitro*, and that cytoooxygenase activation within endothelial cells is likely to play a key role in this process (De Vries *et al.*, 1996). Alternatively, it has been proposed that soluble signal mediators, such as immune cytokines, may gain entry to the brain through the circumventricular organs (CVOs), which lack a BBB and provide for the exchange of blood-borne high molecular polypeptides between blood and brain (Stitt, 1990; Blatteis, 1992). The CVOs most frequently examined in neuroimmunology are the median eminence (ME) and the organum vasculosum laminae terminalis (OVLT). It has been suggested that the systemic influence of cytokines on hypothalamic-pituitary-adrenal axis (HPA) activity is exerted primarily at the level of nerve terminals in the ME (Matta *et al.*, 1990; Rivest *et al.*, 1992), and the OVLT has been implicated in the pathogenesis of fever (Stitt, 1990; Blatteis, 1992). Recently, Hare *et al.* (1995) showed that CVOs other than the ME and the OVLT are implicated in the immune-brain dialog. Following *i.v.* treatment with lipopolysaccharide (LPS), an increased number of cells positively staining for Fos could be observed also in the area

postrema (AP) and subfornical organ (SFO). Since the neurons in the CVOs are connected with nearby hypothalamic regions and neighboring brain stem regions, Fos protein expression also was investigated in further parts of the brain. LPS-induced increases in the number of Fos-stained cells could be detected in the major noradrenergic cell groups (A1, A2, A6), in the paraventricular and supraoptic nuclei and in limbic structures, such as the bed nucleus of the stria terminalis (BST), the central nucleus of the amygdaloid body, and the ventral septal area. These observations are similar to those reported by others (Gaykema *et al.*, 1995). As reviewed by Johnson and Gross (1993), it is well established that the CVOs contain a number of hormones, neurotransmitters and transmitter-like factors, e.g. adrenocorticotropin (ACTH), angiotensin, enkephalins, neuropeptide Y, VIP, somatostatin, substance P, epinephrine, NE, acetylcholine, serotonin (5-HT), dopamine,  $\gamma$ -aminobutyric acid (GABA), glutamate and aspartate, as well as high concentrations of respective binding sites. Many of these substances with receptors in the CVOs have been shown to be effective in inducing changes in neuronal firing. Together, these neuronal systems can reach many further parts of the CNS, including limbic forebrain structures. Based on these findings, it is reasonable to assume that CVOs have a crucial role in translating cytokine signals into neuronal activity. Considering the concept of a humoral immune-brain route, however, it should be kept in mind that central illness responses, such as fever, also can be observed without detectable elevations of cytokine blood levels (Kluger, 1991).

A second mechanism, by which immune cells modify CNS functions, may involve the PNS. Such a neural immune-brain route would not require changes in systemic cytokine concentrations. Accordingly, the translation of humoral into nerve signals may occur in the periphery, from where the CNS is activated via the most common afferents of the body, i.e. the vagus nerve and/or the nociceptive afferents, which travel in the sympathetic parts of the PNS.

Evidence continues to accumulate in the literature suggesting a prominent role for the vagus in neuroimmunomodulation. Nijima (1992) reported an increased afferent discharge rate of the hepatic branch of the vagus nerve following administration of IL-1 $\beta$  into the portal venous blood. Subdiaphragmatic vagotomy blocked hyperalgesia, produced by *i.p.* LPS (Watkins *et al.*, 1994a), *i.p.* IL-1 $\beta$  (Watkins *et al.*, 1994b), and by *i.p.* TNF- $\alpha$  (Watkins *et al.*, 1995a). More specifically, the *i.p.* LPS-induced prolonged enhancement of pain responsiveness also is abolished by cutting only the hepatic branch of the vagus (Watkins *et al.*, 1994a), and hepatic macrophages (Kupffer cells) serving as sources for cytokines have been shown to be involved critically in this pathway (Watkins *et al.*, 1994b). Concerning the neurocircuitry of LPS- or cytokine-induced hyperalgesia, data of Watkins *et al.*, 1994a, suggest that afferent information travels via the hepatic vagus to the nucleus tractus solitarius. From here, an efferent pathway arises from the nucleus raphe magnus to the spinal cord via the dorsolateral funiculus involving a spinal substance P, cholecystokinin (CCK), *N*-methyl-D-aspartic acid (NMDA)-nitric oxide (NO)

cascade as well as non-NMDA excitatory amino acid receptors (Watkins *et al.*, 1994c; Wiertelak *et al.*, 1994).

In further support of vagal afferents, subdiaphragmatic vagotomy strongly inhibited endotoxin-induced Fos expression in the paraventricular nucleus

of the hypothalamus, specifically in corticotropin-releasing hormone (CRH)-containing cells, and blocked the ACTH response following low-dose i.p. endotoxin treatment (Gaykema *et al.*, 1995). However, hepatic branch vagotomy did not affect endotoxin-induced Fos expression and, rather,




Fig. 3. Afferent pathways in the immune-brain dialog. Abbreviations: CeMA, centromedial amygdala; BST, bed nucleus of the stria terminata; HF, hippocampal formation; PAG, periaqueductal grey; CVOs, circumventricular organs; ME, median eminence; OVLt, organum vasculosum laminae terminalis; AP, area postrema; SFO, subfornical organ.

potentiated the ACTH response to low-dose endotoxin. Thus, different organ-specific impulses from visceral receptors may reach the CNS through different vagal afferents. This notion is supported by findings demonstrating that the activation of vagal afferents mediates a broad range of responses, such as an IL-1 $\beta$ -induced corticosterone elevation and hypothalamic NE depletion (Fleshner *et al.*, 1995), LPS-induced sickness behavior (Bluthé *et al.*, 1994), LPS-induced central IL-1 $\beta$  mRNA expression (Layé *et al.*, 1995), IL-1-induced hyperthermia (Watkins *et al.*, 1995b), LPS-induced fever (Sehic and Blatteis, 1996) and cytokine-induced conditioned taste aversion (Goehler *et al.*, 1995). Apparently in contrast to Fleshner *et al.* (1995), who demonstrated a vagally mediated IL-1 $\beta$ -induced corticosterone increase, our own results in rats revealed that peripheral cholinergic stimulation with physostigmine on days 1 and 2 before immunization with sheep red blood cells (SRBC) strongly suppressed the rise of plasma corticosterone via muscarinic mechanisms (Rinner and Schauenstein, 1991).

As most of the peripheral pain fibers ascend with the sympathetic innervation, cytokine-induced hyperalgesia also may follow a sympathetic course. In this context, plantar injection of IL-1 $\beta$  was found to enhance neural discharge in dorsal root filaments in response to mechanical and thermal stimulation, and to decrease the threshold for the response to pressure stimulation (Fukuoka *et al.*, 1994). Since substance P mediates peripheral hyperalgesia (Nakamura-Craig and Smith, 1989), and IL-1 $\beta$  regulates the expression of the peptide and its receptor in sympathetic ganglia (see above), cytokines most likely contribute to pain facilitation. On the other hand, interleukins also can induce local analgesic effects. IL-1 $\beta$  and CRF have been found to inhibit pain by releasing opioid peptides from immune cells within inflamed tissue (Stein *et al.*, 1990; Schäfer *et al.*, 1994). Studies on joint inflammation have shown that primary afferents became rapidly sensitized during a developing arthritis (Levine *et al.*, 1986a; Schaible and Schmidt, 1988), and that several neuropeptides, such as substance P, calcitonin gene-related peptide (CGRP), neuropeptide Y, and neurokinin A, appear to be involved in the inflammatory process (Levine *et al.*, 1984; Donaldson *et al.*, 1992; Mapp *et al.*, 1993; Bileviciute *et al.*, 1993, 1995). In addition to these peripheral neurogenic mechanisms, recent data demonstrate that also central, even though different, regulatory mechanisms are involved in the course of adjuvant-induced knee joint monoarthritis. Bileviciute *et al.* (1995) observed various changes in the concentrations of the same neuropeptides in hypothalamus, hippocampus, striatum, and cortex of arthritic rats, but only few correlations between neuropeptide concentrations in brain and periphery. Finally, recent *in vitro* data suggest peptidergic (neurokinin A, VIP) sensory nerve terminals also to be involved in the modulation of antibody responses in the spleen (Hikawa and Takenaka, 1996). However, in spite of the PNS's apparent importance in the immune-brain dialog, it is largely unknown which of the afferent routes and which of the many neurotransmitters are responsible for information processing of a particular type of immune challenge.

## 2.2. Efferent Pathways

Early studies exclusively examined neuroimmune interactions at the level of the hypothalamus. Besedovsky *et al.* (1975) were the first to describe the neuroendocrine feedback regulation of immune responses via glucocorticoid hormones. This glucocorticoid peak at the time of maximal antibody production after antigenic challenge is caused by central activation of the HPA axis. It contributes to the specificity of immune reactions (Besedovsky *et al.*, 1979), and defects of this circuit predispose to unwanted immune reactions, as suggested by data in animal models with spontaneous (Schauenstein *et al.*, 1987), as well as experimentally induced autoimmune diseases (Sternberg *et al.*, 1989a, 1989b; Mason *et al.*, 1990; Hu *et al.*, 1993). The primary event in this pathway appears to be an increased hypothalamic turnover of NE following immune activation (Besedovsky *et al.*, 1983; Zalcman *et al.*, 1991; Shanks *et al.*, 1994), which can give rise to an activation of CRH neurons (Berkenbosch *et al.*, 1987; Sapolsky *et al.*, 1987). Based on these data, it has long been established that the immune system is modulated by hypothalamic-pituitary (HP) axes (Besedovsky *et al.*, 1986; Rivier and Vale, 1989; Chrousos, 1995; Savino and Dardenne, 1995), and the role of central cytokines in the activation of HP axes has been the subject of numerous studies (Harbuz *et al.*, 1992; reviewed by Schöbitz *et al.*, 1994a).

Besides these neuroendocrine pathways, there is evidence for neural routes, by which the brain regulates peripheral immune functions. Several data suggest the sympathetic nervous system to convey information from the CNS to the peripheral immune system (Sundar *et al.*, 1989, 1990; Croiset *et al.*, 1990; Irwin *et al.*, 1990, 1991, 1992; Irwin, 1994), and adrenoceptor-mediated effects on lymphocyte functions have been defined by several groups, including ourselves (Livnat *et al.*, 1987; Felsner *et al.*, 1992, 1995; Sanders, 1995). The specific role of the sympathetic nervous system in the modulation of peripheral immune activity has been underscored by a number of recent findings showing that (i) sympathetic denervation results in a wide variety of strain-, organ-, or even compartment-specific effects, such as increases in cellular proliferation and lymphocyte migration (Madden *et al.*, 1994a), enhanced macrophage phagocytic function (Lyte *et al.*, 1991), reductions in thymus weight and thymocyte numbers (Tsao *et al.*, 1996), thymocyte apoptosis (Tsao *et al.*, 1996), and heterogeneous alterations in B- or T-cell responsiveness, as well as cytokine (IL-2/IFN- $\gamma$ ) production (Lyte *et al.*, 1991; Madden *et al.*, 1994b); (ii) the IL-2-induced enhancement of an IgM plaque-forming cell response requires an intact sympathetic innervation of the spleen and is blocked selectively by the  $\beta$ -adrenergic antagonist propranolol (Zalcman *et al.*, 1994a); (iii) the immunosuppressive effects of stress on humoral and cell-mediated immune responses of rat splenocytes are mediated via adrenergic fibers of the splenic nerve (Wan *et al.*, 1993); and (iv) local NE release is involved in the regulation of IL-6 secretion in the spleen (Straub *et al.*, 1995). Furthermore, there exists

a large body of evidence suggesting a regulatory role for the sympathetic nervous system in the pathogenesis or progression of diseases, such as experimental and clinical arthritis (Levine *et al.*, 1986a, 1986b, 1988; Felten *et al.*, 1992; Lorton *et al.*, 1996), experimental allergic encephalomyelitis (Chelmicka-Schorr *et al.*, 1988, 1989) and experimental autoimmune myasthenia gravis (Agius *et al.*, 1987).

Regarding immunomodulatory effects of vagal efferent fibers, there is increasing evidence that parasympathetic activity plays a prominent role in thymic lymphopoiesis (Singh and Fatani, 1988), apoptosis, as demonstrated by our group (Rinner *et al.*, 1994), and lymphocyte traffic from thymus to peripheral lymphoid organs. Antonica *et al.* (1996) observed an increased export of immature CD4<sup>-</sup>CD8<sup>-</sup>, double positive (DP), and CD4<sup>-</sup>CD8<sup>-</sup>, double negative (DN), cells from mouse thymus following monolateral vagal denervation, and that both DP and DN cells migrate more consistently to lymph nodes and spleen of vagotomized mice compared to controls. In parallel, the percentage of single positive (SP) cells was decreased in peripheral organs. These results, together with the findings of a preeminent role of the thymic microenvironment in positive and negative selection of developing T-cells, and thereby in the avoidance of autoimmunity (Boyd *et al.*, 1993), emphasize the complex interactions between the nervous and immune systems.

As a further efferent mechanism, central LPS as such, or a fragment, could leak through the BBB and induce cytokine production in the periphery. This has been suggested by comparing the time-course of serum IL-6 induction after peripheral or central LPS administration (De Simoni *et al.*, 1992, 1995; Gottschall *et al.*, 1992b). Further descending brain-immune routes were identified by showing that, aside from CRH (Schulte *et al.*, 1994), several other neuropeptides, such as substance P and VIP (Stanisz *et al.*, 1986; Laurenzi *et al.*, 1990; Spangelo *et al.*, 1990; Agro and Stanisz, 1995), can transmit central effects towards the periphery, and there is growing evidence that central opioid pathways are critically involved (De Simoni *et al.*, 1993; Fecho *et al.*, 1996). However, despite intense research efforts, many aspects of the CNS-mediated immunoregulation are still poorly understood. Nor do we exactly know in which way, or to what extent, different brain sites influence specific immune activities. For example, Hernandez *et al.* (1993) reported that injection of morphine into either the third ventricle or the anterior hypothalamus significantly suppressed blood lymphocyte proliferation. Weber and Pert (1989) examined the functional role of the midbrain periaqueductal grey (PAG) in opiate-induced immunosuppression. Microinjection of morphine into the PAG resulted in a rapid suppression of splenic natural killer (NK) cell activity. Injection into the anterior hypothalamus, arcuate nucleus, medial amygdala, medial thalamus, and dorsal hippocampus was without an effect on NK activity. Recently, the neuronal networks connecting the PAG with a large number of brain structures, including the nucleus raphe magnus, ventrolateral medulla, hypothalamus, amygdala, and

prefrontal cortex, have been described in detail by Behbehani (1995). In addition, this author reviewed the pharmacology of PAG-mediated pain, analgesia, anxiety and fear, vocalization or autonomic regulation, which mainly involves opiates, GABAergic interneurons, neurotensin, substance P, excitatory amino acids, catecholamines, 5-HT-containing cell bodies and CCK-responsive neurons. None the less, little or virtually nothing is known whether, how, or to what extent all these neurotransmitters are implicated in brain immunomodulation via specific structures such as the PAG.

The issue of neuroimmunomodulation is complicated further by the fact that, aside from hypothalamus and brain stem regions, additional brain structures are involved in the modulation of peripheral immune responses. Selective alterations of different components of the immune system have been observed in animals subjected to experimental lesions of the cerebral cortex (Renoux *et al.*, 1987), the lateral septal area (Wetmore *et al.*, 1994), the nucleus basalis magnocellularis, the equivalent of Meynert's nucleus in the primate (Cherkaoui *et al.*, 1990), and further limbic structures (see below). In view of the close relations between cholinergic activity and immunity (see above), it should be kept in mind that the basal nucleus of Meynert contains, similar to the septum, not only large cholinergic but also GABAergic neurons, and has been shown to be involved in learning, memory and attention (Lamour *et al.*, 1989; Durkin, 1994; Muir *et al.*, 1994). In addition, septum and nucleus basalis of Meynert serve as one of the diverse neuroanatomical substrates providing for cortical activation or arousal (Riekkinen *et al.*, 1991; Osborne, 1994). However, at present the specific role of both these structures in neuroimmunology has not been defined, and should be subject of future investigations.

What we can conclude from this diversity of information is that (i) the brain as a whole participates in the modulation of peripheral immunity; (ii) humoral and neural pathways convey information from the brain to the peripheral immune system; (iii) neuroimmunomodulation includes brain sites involved in emotional processing and cognitive functions as the neuroanatomical substrate of psychoimmunomodulation.

### 3. THE LIMBIC SYSTEM

The "limbic system" is used as a collective term for various brain structures in the basomedial telencephalon, diencephalon and mesencephalon, all of which have been implicated in cognitive functions, emotions, behavior, neuroendocrine and autonomic responses. The most commonly cited structures comprise the hippocampal formation with the hippocampus, dentate gyrus and subiculum, as well as the amygdaloid body, in particular a recently reconsidered brain region referred to as the "extended amygdala" (centromedial amygdaloid-bed nucleus of the stria terminalis continuum) (Martin *et al.*, 1991; Schmued, 1994; Heimer, 1995a). Aside from these areas, many other parts of the brain, including the old portions of the "rhinencephalon",

the cingulate gyrus, parahippocampal gyrus with the entorhinal cortex and praesubiculum, the ventral parts of the striatal complex, septum, anterior thalamic nuclei, medial thalamic nuclei and the habenula, have been recognized to be "limbic" (Heimer, 1995b). It is generally accepted that all these structures are related to each other as well as to neighboring non-limbic areas by a network of widespread fiber projections providing the basis of highly organized neuronal circuits. However, it is all the same now fairly certain that there does not exist an anatomical or functional unit referable to a "limbic system". Therefore, for clarity's sake, the following chapters will focus on the roles of the hippocampal-amygdala group and the nucleus accumbens in psychoneuroimmunomodulation. As one pathway, the hippocampal formation is related reciprocally with the amygdaloid body via the amygdalohippocampal fibers (Aggleton, 1986; Amaral, 1986; Mello *et al.*, 1992). The extended amygdala itself projects to many regions in the hypothalamus as well as to somatomotor and autonomic areas in the brain stem, with further effects on the spinal cord (Martin *et al.*, 1991; Schmued, 1994; Heimer, 1995a). On the other hand, as shown e.g. in the visual system, information processing also may involve the simultaneous activation of different areas or distributed neuronal networks, including cortico-cortical re-entrant circuits (Bullier and Nowak, 1995). Besides the visual cortex, other re-entrant loops have been recognized, which appear to involve the basal ganglia (Alexander *et al.*, 1989; Parent and Hazrati, 1995). Accordingly, as a second route, a mesolimbic component of such a cortico-basal ganglia-thalamo-cortical re-entrant circuit is considered. The nucleus accumbens is part of the "ventral striatopallidal system" (Heimer, 1995c), which primarily receives inputs from the hippocampal formation, the basolateral amygdala, and temporal, as well as prefrontal cortical areas. Interestingly, some impulses return to the anterior cingulate area via the mediodorsal thalamic nucleus (Heimer, 1995d). The nucleus accumbens itself is known to be of relevance to both cognitive functions (Schacter *et al.*, 1989; Salamone, 1994), as well as the central processing of immune signals (see below). Thus, psychoimmunomodulation appears to be more complex than previously supposed. Figure 4 is an attempt to visualize both routes in one scheme.

In the following, some functional characteristics of limbic structures will be summarized briefly. The hippocampal formation has been attributed with important functions in learning processes (Grossberg and Merrill, 1992), memory and retrieval (Squire and Zola-Morgan, 1989; Mishkin, 1993) and, accordingly, has been implicated in Alzheimer's disease (Hyman *et al.*, 1984; Mullan, 1993; Wakabayashi *et al.*, 1994). The amygdala is critically involved in the processing of emotional stimuli (Davis, 1992), whereby hemispheric differences have been observed (Adamec and Morgan, 1994), and recent evidence suggests that amygdala neurons serve important functions in emotional learning and memory storage (reviewed by Ono *et al.*, 1995). Local infusion of CRF into the amygdala has been

found to affect aggressive as well as investigative behavior in rats (Liang and Lee, 1988; Elkabir *et al.*, 1990). Furthermore, the medial amygdala is well established to modulate defensive rage behavior (Shaikh *et al.*, 1993), as well as predatory attack behavior in the cat (Han *et al.*, 1996a, 1996b). On the one hand, medial amygdaloid stimulation has been shown to facilitate defensive rage behavior elicited from the medial hypothalamus involving a substance P pathway from the medial amygdala to the medial hypothalamus (Shaikh *et al.*, 1993). On the other hand, the medial amygdala suppresses predatory attack behavior elicited from the lateral hypothalamus via a disynaptic pathway which, first, runs from the medial amygdala to the ventromedial hypothalamus utilizing substance P as excitatory neurotransmitter and, second, includes inhibitory GABAergic projections from the ventromedial hypothalamus to the lateral hypothalamus (Han *et al.*, 1996a, 1996b). These data indicate close relationships between limbic areas and different parts of the hypothalamus, supporting the notion of multifunctional neuronal networks involved in the control of behavioral activity. In view of the importance of such distributed organization, it would be of special interest to determine whether similar pathways exist in mediating immune-induced behavioral responses. In addition, hippocampus and amygdala both have been shown to contain receptors for corticotropin-releasing factor (CRF) (Millan *et al.*, 1986; Lovenberg *et al.*, 1995; Avishai-Eliner *et al.*, 1996), glucocorticoids (GR) (Sapolsky *et al.*, 1983; McEwen *et al.*, 1986; De Kloet *et al.*, 1994) and mineralocorticoids (MR) (Cairini *et al.*, 1985; McEwen *et al.*, 1986; De Kloet *et al.*, 1994). It has become clear that the basal and stress-induced HPA activity is coordinately controlled by the high-affinity MRs (Type I receptors) and the low-affinity GRs (Type II receptors) (De Kloet, 1991), and that the hippocampus, particularly the dorsal part, plays a prominent role in modulating the negative feedback effects of glucocorticoids on HPA axis activity (review by Jacobson and Sapolsky, 1991; Feldman and Weidenfeld, 1993). In a series of papers (Beaulieu *et al.*, 1986, 1987; Feldman *et al.*, 1994), the central and medial nucleus of the amygdala were identified as a regulatory sites of ACTH secretion. Back in the seventies, Carroll *et al.* (1976) already argued that a limbic-hypothalamus disinhibition of the pituitary-adrenal axis occurs in patients with depressive disorders. Aside from adrenal steroids, gonadal and thyroid hormones also have been shown to influence hippocampal physiology and morphology (Gould *et al.*, 1991; McEwen and Woolley, 1994; Woolley and McEwen, 1994). However, by far the majority of studies on limbic structures have addressed their roles in the neuroendocrine stress response, and several methods including psychosocial stress, immobilization and cold restraint-induced gastric ulcerogenesis have been used in animal models to examine the effects of stressful stimulation on hippocampus, amygdala, BST, as well as nucleus accumbens (see below). Given these diverse functions of limbic neuronal systems, it is not surprising that a large number of receptor proteins, neuropeptides




Fig. 4. Processing of external stimuli by hippocampal-amygdala and/or ventral striatopallidal structures. Pathway 1 (solid): cortex-“extended amygdala”-(centromedial amygdaloid-bed nucleus of the stria terminalis continuum). Pathway 2 (striped): cortico-basal ganglia thalamo-cortical “re-entrant loop” via the “ventral striatopallidal system”. Additional connections (plain). Abbreviations: CeMA, centromedial amygdala; BLA, basolateral amygdala; BST, bed nucleus of the stria terminalis; HF, hippocampal formation; SI, substantia innominata; ST, stria terminalis; PAG, periaqueductal grey.

and neurotransmitters have been identified in these regions. Current data suggest the involvement of noradrenergic (Beaulieu *et al.*, 1987; Dalmaz *et al.*, 1993; Gobert *et al.*, 1993; Kiss *et al.*, 1995), cholinergic (Finkelstein *et al.*, 1985; Gilad *et al.*, 1985; Dalmaz *et al.*, 1993; Costa *et al.*, 1994; Aigner, 1995; Rieck *et al.*, 1995; Komourian and Quik, 1996; Tice *et al.*, 1996), GABAergic (Rainnie *et al.*, 1991a,

1991b; Mello *et al.*, 1992; Orchinik *et al.*, 1995), serotonergic (Beaulieu *et al.*, 1986; Martin *et al.*, 1992; Holmes *et al.*, 1995; Matsumoto *et al.*, 1995; Zhong and Ciaranello, 1995; Morales *et al.*, 1996), dopaminergic (Beaulieu *et al.*, 1987; Gasbarri *et al.*, 1994; Nistico *et al.*, 1994) and glutamatergic mechanisms (Clark and Cotman, 1992; Liang *et al.*, 1994; Aigner, 1995; Blümcke *et al.*, 1995; Day *et al.*,

1995; Arai and Lynch, 1996), as well as several neuropeptides, including CRF (Ray *et al.*, 1993; Kalin *et al.*, 1994), arginine vasopressin (AVP) (De Vries *et al.*, 1985; Brinton *et al.*, 1994; Hirasawa *et al.*, 1994; Van Wimersma Greidanus and Maigret, 1996), oxytocin (Van Wimersma Greidanus and Maigret, 1996), substance P (Rieck *et al.*, 1995; Han *et al.*, 1996a), VIP (Léránth and Frotscher, 1983; Roberts *et al.*, 1984), pituitary adenylate cyclase activating polypeptide (PACAP) (Palkovits *et al.*, 1995), neurotensin (Roberts *et al.*, 1984), CCK (Roberts *et al.*, 1984; Hamilton and Freeman, 1995), neuropeptide Y (Zachrisson *et al.*, 1995), enkephalins (Rieck *et al.*, 1995), dynorphin (Watanabe *et al.*, 1995a), and somatostatin (Roberts *et al.*, 1984; Obata-Tsuto, 1987; Zachrisson *et al.*, 1995).

Besides these classical neuronal and neuroendocrine components, the expression of cytokines and their receptors has been demonstrated in all limbic areas, as outlined in Table 1. Noteworthy in this regard is that spontaneous autoimmune diseases in the NZB and (NZB/NZW)F1 mouse model have been shown to be associated with a deficit in the density of IL-1R in the dentate gyrus exclusively (Jafarian Tehrani *et al.*, 1994). The IL-1 $\beta$  expression in the hippocampal formation also is affected by chronic exposure to morphine. Using immunocytochemical analysis, Patel *et al.* (1996) recently reported a marked decrease of IL-1 $\beta$  staining in the polymorphic region of the dentate gyrus and CA1-CA3 subfields of the rat hippocampus after chronic treatment with morphine. Finally, classical neurotransmitters such as 5-HT even at very low concentrations were recently shown to induce cytokine mRNA expression in rat primary hippocampal astrocytes (Pousset *et al.*, 1996).

Table 1. Cytokines and their Receptors in Limbic Areas

Receptor	Limbic area
IL-1R	Molecular and pyramidal cell layer of the hippocampus Molecular and granule cell layer of the dentate gyrus Cingulate cortex Entorhinal cortex Subiculum Basolateral amygdala Olfactory bulb Medial habenula Anterior dorsal thalamus Septum Nucleus accumbens
IL-1	Pyramidal cell layer of the hippocampus Granule cell layer of the dentate gyrus Olfactory bulb Bed nucleus of the stria terminalis Substantia innominata
IL-1RA	Dentate gyrus
IL-2R/IL-2	Pyramidal cell layer of the hippocampus Granule cell layer of the dentate gyrus
IL-3R	Hippocampus
IL-3	Hippocampus Medial habenula
IL-6R/IL-6	Hippocampus
IL-8	Hippocampus
TNF- $\alpha$	Hippocampus

## 4. NEUROIMMUNE COMMUNICATIONS VIA LIMBIC STRUCTURES

### 4.1. Lesioning Limbic Areas Affects Immune Responses

Lesions of the dorsal hippocampus or amygdaloid complex were found to produce a transient increase in splenocytes and thymocytes, as well as increased T-cell mitogen responses (Brooks *et al.*, 1982), and hypophysectomy was shown to reverse these effects (Cross *et al.*, 1982). Similarly, lesions of the hippocampus were found to cause differential effects on humoral immunity depending on the hippocampal fields and/or axonal pathways destroyed (Pan and Long, 1993). Alterations of cellular immune responses following small electrolytic lesions in different parts of the brain support the notion that the cingulate cortex, the central and the basomedial nucleus of the amygdala participate in the control of immunoneuroendocrine interactions (Masek *et al.*, 1992). Unilateral dopaminergic lesions of the nucleus accumbens have been shown to result in decreased splenic NK cell activity in left-lesioned mice as compared to right-lesioned animals, and unilateral striatal dopaminergic lesions have been found to impair the proliferation of splenic lymphocytes in the right-lesioned group only (Deleplanque *et al.*, 1994). These results suggest that immunologic reactivity is asymmetrically modulated by central dopaminergic pathways. Even though these data altogether show that structural disturbances within limbic areas affect immune functions, it should be kept in mind that, in view of the widespread and intimate relations of brain structures, results of experimental lesioning protocols should be interpreted with caution.

### 4.2. Neuroendocrine Immunomodulation at Limbic Neuronal Systems

As will be shown in the following, a number of recent experimental data strongly argue for an integration of limbic areas into immunoneuroendocrine circuits. Injection of IL-1 $\beta$  into the lateral cerebral ventricle reduced the affinity of hippocampal MRs in parallel to an enhanced HPA activity, and caused a shift in the MR/GR balance, which has been suggested to be of importance to the regulation of stress responses, behavioral activity, hippocampal excitability in general, long-term potentiation, and immune-induced neuroendocrine effects (De Kloet *et al.*, 1994; Schöbitz *et al.*, 1994b; McEwen and Sapolsky, 1995). It was concluded that this mechanism allows for an individual modulation of HPA axis activity during basal conditions, or in response to different amounts of stressful stimulation, including immune challenge. Another interesting aspect is that prenatal immune challenge has been shown to decrease MR and GR levels in the hippocampus of adult progeny, and to induce abnormalities in HPA axis function (Reul *et al.*, 1994). Furthermore, functional abnormalities in the HPA axis are often observed in pathological conditions, such as Alzheimer's disease, certain depressive illnesses, panic disorder, as well as multiple sclerosis (O'Brien *et al.*, 1993; Michelson *et al.*, 1994; Miller *et al.*, 1994;

Abelson and Curtis, 1996), and in major depression a hyperactivity of the HPA axis has been correlated with an enhanced IL-1 $\beta$  production (Maes *et al.*, 1993). In addition, recent studies addressing steroid hormone effects on neurons and behavior revealed a rich diversity of genomic and non-genomic steroid actions on neuronal properties (for review see Weeks and Levine, 1995). The work by Sapolsky (1986), Sapolsky and Pulsinelli (1985), Sapolsky *et al.* (1985, 1988, 1990) *e.g.*, pointed out the deleterious effects of prolonged glucocorticoid exposure on hippocampal neurons. On the other hand, a deficient immunoneuroendocrine feedback loop via the HPA axis has been implicated in animal models of spontaneous and experimentally induced autoimmune diseases (see above). In this context, an interesting recent study shows a number of immune abnormalities in transgenic mice expressing Type II GR antisense RNA comprising a deficient immunoneuroendocrine dialogue, alterations in thymocyte trafficking/migration, a shift of the T-cell balance towards the CD4 helper subset, and an absence of sex dimorphism in the development of GR mRNA expression and cell-mediated immunity (Morale *et al.*, 1995). Furthermore, Fischer 344 (F344) rats, which are known to mount a significantly higher corticosterone response to stress or to an immune challenge compared to Lewis and Sprague Dawley (SD) rats, are found to exhibit greater stress-induced Type II receptor activation in the cortex, hippocampus, and hypothalamus than SD and Lewis rats (Dhabhar *et al.*, 1995a). In addition, whereas Fischer rats serve as models for aging and cancer (Solleveld *et al.*, 1984; Ghanta *et al.*, 1990), Lewis rats are prone to autoimmune diseases (Sternberg *et al.*, 1989a, 1989b; Mason *et al.*, 1990). However, whether a defect in the hippocampal neuronal system is involved in the pathogenesis or exacerbation of autoimmune diseases needs to be clarified in future experiments.

In another approach to examine the effects of hippocampal IL-1 on brain functions, Linthorst *et al.* (1994) used a microdialysis probe, which allows stress-free intrahippocampal administration of recombinant human IL-1 $\beta$  (hIL-1 $\beta$ ) in close vicinity of field CA3 and the dentate gyrus of rats. The intrahippocampal infusion of hIL-1 $\beta$  caused (i) a stimulation of HPA activity as reflected by increased plasma levels of ACTH, as well as plasma and dialysate levels of corticosterone; (ii) increases in hippocampal extracellular 5-HT levels in parallel with a decrease in behavioral activity; and (iii) an elevation of body temperature. The neural pathways mediating these effects need to be investigated further, but appear to involve serotonergic neurotransmission, possibly the nuclei raphe (Taylor, 1982; Werner and Bienek, 1985; Jacobs and Azmitia, 1992) and projections from hippocampus to nuclei raphe via septum (Swanson *et al.*, 1987; Cui *et al.*, 1993) or extended amygdala (Heimer, 1995e).

In further evidence of neuroimmunologic regulation exerted by limbic structures, Hauger *et al.* (1993) have shown the effects of high intracerebral levels of CRH on the regulation of CRH receptors in amygdala and anterior pituitary as well as on the modulation of immune functions. Chronic CRH

pretreatment resulted in decreased CRH receptor concentrations exclusively in the amygdala, and splenic NK responses became insensitive to an acute intracisternal CRH injection. Although the central mechanisms of neuroendocrine–neuroimmune regulation are not yet known in detail, brain CRH (Irwin, 1994; Irwin *et al.*, 1987) and autonomic nervous system mechanisms (Irwin, 1994; Irwin *et al.*, 1990, 1992) have been shown to be implicated. Moreover, aside from different patterns of sympathetic dysregulation (Roy *et al.*, 1985; Rudorfer *et al.*, 1985; Veith *et al.*, 1994), in depression chronic CRH-hypersecretion has been reported (Nemeroff *et al.*, 1984, 1992), and reduced CRH binding-sites have been found in the frontal cortex (Nemeroff *et al.*, 1988). Both CRH activation and increased sympathetic nervous system activity may contribute to selective immune alterations observed in different types of depressive disorders. Supporting this view, Irwin *et al.* (1991) suggested that the activation of the sympathetic nervous system and elevated plasma levels of neuropeptide Y are associated with a reduction of NK activity in depressed patients. However, the interactions between neuropeptide Y, immunity, and depression appear to be rather complex, since central administration of neuropeptide Y also can result in immunoenhancing effects, as shown in the olfactory bulbectomized (OB) rat model of depression (Song *et al.*, 1996). A subchronic i.c.v. low-dose infusion of neuropeptide Y (i) reversed the deficit in NE concentrations in the amygdala of OB rats; (ii) decreased the concentrations of 5-hydroxyindole acetic acid (5-HIAA) in the hypothalamus; (iii) reduced hyperactivity of OB rats in the open field; (iv) elevated the concentrations of corticosterone; and (v) interestingly increased lymphocyte responses to phytohemagglutinin (PHA) and concanavalin A (ConA), which normally are suppressed in these animals, while not affecting the changes in the differential white blood cell count and the elevated PHA-induced chemiluminescence of mononuclear cells (Song *et al.*, 1996).

As AVP plays a significant role in the immunoneuroendocrine dialog (Nakatsuru *et al.*, 1991; Zelazowski *et al.*, 1993), it is of note that the amygdala, as well as the BST, contain AVP synthesizing cell bodies (De Vries *et al.*, 1985), and nitric oxide synthase has been detected in the amygdala (Pow, 1992). IL-2 has been shown to stimulate AVP release from hypothalamus and amygdala *in vitro* in a calcium-dependent manner, possibly via NO-mediated signaling (Raber and Bloom, 1994). Analogous *in vitro* results have been obtained regarding IL-2-induced CRH release from the hypothalamus as well as from the amygdala (Raber *et al.*, 1995), and it has been established that NE, as well as acetylcholine, enhances the *in vitro* release of CRH (Raber *et al.*, 1995) and AVP (Raber and Bloom, 1994) from the amygdala, involving NO-mediated signaling in the case of acetylcholine.

If limbic areas are in fact involved in the central regulation of the immunoneuroendocrine dialog, the exploration of the neurotransmitters involved in this issue will be of particular interest. Similar to results obtained in the hypothalamus, antigen challenge results in alterations of NE (Carlson *et al.*, 1987;

Zalcman *et al.*, 1991) as well as changes in 5-HT (Carlson *et al.*, 1987) in the hippocampus. Furthermore, dopamine levels in the nucleus accumbens were found decreased following administration of antigen at or about the time of peak immune responses (Zalcman *et al.*, 1991; Shanks *et al.*, 1994). Using *in vivo* microdialysis, Lacosta *et al.* (1994) detected an increase in extracellular dopamine in the nucleus accumbens following immunization. This *in vivo* dopamine elevation is a likely result of an increased release of dopamine from neurons. An interesting point is that immune-induced dopamine alterations in the nucleus accumbens appear to be more pronounced in isolated animals compared to animals housed in groups (Zalcman *et al.*, 1991; Lacosta *et al.*, 1994; Shanks *et al.*, 1994). Other studies have investigated cytokine-specific effects on monoamine activities. According to Zalcman *et al.*, 1994b), peripherally (i.p.) administered IL-1 increased NE turnover in the hypothalamus as well as in the hippocampus, and 5-HT turnover in the hippocampus and prefrontal cortex. IL-6 (i.p.) increased 5-HT activity in the hippocampus as well as the prefrontal cortex, and IL-2 (i.p.) induced a markedly increased hypothalamic NE utilization. All three cytokines modified dopamine activities in the prefrontal cortex (Zalcman *et al.*, 1994b), a brain region most sensitive to stressful stimulation (Roth *et al.*, 1988). Additionally, Zalcman's aforementioned study revealed intriguing results as (i) IL-1-induced increases of plasma corticosterone do not necessarily correlate with IL-1-induced alterations of hypothalamic NE activity; (ii) although IL-2 can markedly enhance NE utilization in the hypothalamus, this does not necessarily result in significant increases of plasma corticosterone concentrations; and (iii) even though IL-6 is able to alter central 5-HT and dopamine activities, it does not obligatorily affect hypothalamic NE activity or plasma corticosterone concentrations. The latter is in line with observations reported by Dunn (1992). The functional significance of these intricate relations will be considered below. Systemically, administered IL-2 also has been found to reduce the *in vivo* dopamine release from the nucleus accumbens (Anisman *et al.*, 1996). As for the amygdala, activation of dopamine D1 receptors has been shown to be involved in the selective enhancement of mitogen (ConA) responsiveness of splenocytes (Nistico *et al.*, 1994). Considering the organization of the basal forebrain cholinergic projections, which allow widespread influence on an array of brain regions, including hippocampus and amygdala (Rieck *et al.*, 1995), and recent investigations confirming the significance of cholinergic activity in learning and memory (Aigner, 1995; Sokolov and Kleschevnikov, 1995), the following data provide convincing evidence that cholinergic neurons in limbic areas have a crucial role in psychoimmunology. As recently summarized by Schauenstein *et al.* (1994), the cholinergic system modulates both immunologic reactivity and the immunoneuroendocrine dialog. Rinner and Schauenstein (1991) have reported that antigenic challenge results in a transient decrease of number and increase in affinity of muscarinic cholinergic receptors in the hippocampus. An important central action of IL-2

has been suggested to be involved in the modulation of hippocampal acetylcholine release (Hanisch *et al.*, 1993), which is in line with previous work by Araujo *et al.* (1989), who have observed that exogenous IL-2 diminished the potassium-evoked release of acetylcholine from rat hippocampal slices. Furthermore, systemically injected IL-1 $\beta$  was found to induce a decrease in extracellular acetylcholine in the hippocampus of freely moving rats (Rada *et al.*, 1991).

To summarize this section, it should be stressed that the hypothalamus is not the only site involved in the modulation of the pituitary-adrenal axis. Besides the evidence that the hippocampus indirectly modulates HPA activity (see above), the central and medial nucleus of the amygdala have been suggested to modulate ACTH secretion (Beaulieu *et al.*, 1986, 1987; Feldman *et al.*, 1994), and to exert differential effects on the CRH content in the ME (Beaulieu *et al.*, 1989; Feldman *et al.*, 1994). Although hypothalamic CRH neurons may be involved in such superior regulatory mechanisms exerted by limbic structures, it seems conceivable that neurosecretory fibers from the extended amygdala and/or hippocampus may project directly to the ME, forming an amygdalo/hippocampo-hypophysial pathway. It remains to be evaluated whether, how, or to what extent external signals (e.g. stress conditions including an immune challenge) are capable to induce the activation of such a route. Alternatively, a number of observations suggested the BST as a site of convergence conveying information from hippocampus and amygdala to hypothalamus or other parts of the brain (Cullinan *et al.*, 1993; Herman *et al.*, 1994; Pacak *et al.*, 1995), and the belief is held that this pathway is involved in the modulation of HPA activity (Dunn, 1987; Feldman *et al.*, 1990; Gray *et al.*, 1993) (Fig. 5). In line with the notion of additional regulatory mechanisms involved in the neuroimmune dialog, several data indicate that HPA axis activation is not an obligatory consequence following antigen challenge (Stenzel-Poore *et al.*, 1993; Shanks *et al.*, 1994). According to Zalcman *et al.*, 1994b) and Dunn (1992), peripheral cytokines not necessarily activate the HPA axis or affect plasma corticosterone concentrations, and results previously obtained in our own laboratory revealed that the corticosterone response induced by immunization with SRBC is abrogated following peripheral cholinergic treatment with physostigmine (Rinner and Schauenstein, 1991). Even though several trivial reasons, such as different experimental conditions or different dose-dependencies of antigen-specific immune responses, may account for such discrepancies, the available data clearly illustrate the high degree of complexity of central neuroendocrine immunomodulation.

#### 4.3. Immune Stimulation Evokes Electrophysiological Changes in Neuronal Activities in Limbic Areas

A large body of information on neuroimmune communications and cognition has been generated by electrophysiological recordings of neurons. Besedovsky *et al.* (1977) were the first to demonstrate increased neuronal firing rates in the ventromedial




Fig. 5. Modulation of the pituitary-adrenal axis via hippocampus and amygdala. Abbreviations: BST, bed nucleus of the stria terminalis; CRH, corticotropin-releasing hormone; ACTH, adrenocorticotropin.

nucleus of the hypothalamus at the time of maximal antibody production following immunization. Similar results were obtained in the hippocampus (Korneva *et al.*, 1985). Recently, effects of immune-derived cytokines on electrophysiological phenomena in limbic areas have been described. IL-1 $\beta$  induced membrane hyperpolarization in neurons of the basolateral amygdala (BLA), most probably indirectly mediated through enhancement of endogenous GABA and accompanied by a decrease in input resistance (Yu and Shinnick-Gallagher, 1994). As stated in the same report, superfusion of IL-1 $\beta$  reduced the amplitudes of EPSPs (excitatory postsynaptic potentials) and blocked fast and slow IPSPs (inhibitory postsynaptic potentials) evoked by stimulation of the stria terminalis and the lateral amygdala possibly via presynaptic mechanisms. In a related study, IL-1 $\beta$  has been found to enhance GABAergic IPSPs in hippocampal neurons of the pyramidal cell layer (Zeise *et al.*, 1992), which would suggest contrary cytokine effects in different brain regions, but could also be due to different experimental approaches used in these two studies. Furthermore,

IL-1 $\beta$  was shown to have excitatory effects on BST neuronal activity possibly via prostaglandin mobilization (Wilkinson *et al.*, 1993), and both IL-1 $\beta$  (Plata-Salamán and Ffrench-Mullen, 1992) as well as IL-2 (Plata-Salamán and Ffrench-Mullen, 1993) depress voltage-dependent calcium currents in CA1 hippocampal neurons.

It has long been established that learning and memory is related to plastic changes of neuronal pathways, and long-term potentiation (LTP) of synaptic transmission is a widely used model of neuronal plasticity (Bliss and Collingridge, 1993). Cytokine effects on electrically induced potentiation of the synaptic transmission have been primarily described in the hippocampus. Interferons have been shown to inhibit the induction of LTP and to attenuate post-tetanic potentiation (PTP), short-term potentiation (STP) as well as the LTP maintenance phase (D'Arcangelo *et al.*, 1991). IL-2 inhibited the induction of both LTP and STP, attenuated PTP and reduced the maintenance of LTP (Tancredi *et al.*, 1990). Tumor necrosis factor affected LTP negatively (Tancredi *et al.*, 1992), and IL-1 $\beta$  inhibited the

potentiation of synaptic transmission in field CA3 of mouse (Katsuki *et al.*, 1990), as well as in field CA1 of rat hippocampal slices (Bellinger *et al.*, 1993). The underlying mechanisms of cytokine-induced effects on LTP remain to be further clarified, but may include NMDA (Collingridge and Davies, 1989; Bashir *et al.*, 1991) and muscarinic receptors (Burgard and Sarvey, 1990; Sokolov and Kleschevnikov, 1995), the protein kinase C system (Farrar and Andersen, 1985; Hu *et al.*, 1987; Malinow *et al.*, 1988; Reymann *et al.*, 1988), sphingosine-sensitive processes (Malinow *et al.*, 1988; Ballou, 1992; Kolesnick and Golde, 1994), the G protein system (Evans *et al.*, 1987; Goh and Pennefather, 1989; Harnett and Rigley, 1992), as well as endogenous NO (Schuman and Madison, 1991; Watanabe *et al.*, 1995b). Besides mono- and lymphokines, several growth factors have been shown to influence the generation of LTP. Both epidermal growth factor and basic fibroblast growth factor were found to promote hippocampal LTP either in fimbria-fornix-lesioned rats or anaesthetized rats (Ishiyama *et al.*, 1991; Abe *et al.*, 1992). Finally, it should be considered that induction criteria for LTP are not only studied in the hippocampal formation but also in other limbic parts of the brain. Recently, it was suggested that neuronal activities in the ipsilateral BLA are required in part for the induction of LTP in the dentate gyrus *in vivo* (Ikegaya *et al.*, 1995). Moreover, recent work supports the hypothesis of a cooperation of hippocampal formation and amygdala providing electrophysiological evidence of direct neuronal connections between the BLA and the dentate gyrus (Ikegaya *et al.*, 1996).

#### 4.4. Limbic Structures in Stress and Immunity

With the emergence of interdisciplinary research areas, such as psychoneuroimmunology, new insights into the concept of stress as a stereotyped and non-specific response (alarm, resistance, exhaustion) to aversive or noxious stimuli were obtained, indicating the involvement of an array of regulatory mechanisms including corticosteroids, monoamines, acetylcholine, endogenous opioids, melatonin, several pituitary hormones and cytokines (Khansari *et al.*, 1990; Hoffman-Goetz and Pedersen, 1994). As initially mentioned, it is generally accepted that different stressors alter immunity in different ways, and this research is reviewed in detail elsewhere (Dantzer and Kelley, 1989; DeGiorgi and Altomare, 1992; Rinner *et al.*, 1992; Fricchione and Stefano, 1994; Irwin, 1994; Kort, 1994; Dhabhar *et al.*, 1995b). Noteworthy in this regard is an intriguing finding, which suggests that the adrenal gland may be the source of plasma IL-6 following exposure to physical and psychological stressors, and that  $\beta$ -adrenergic receptor stimulation is not involved in mediating this effect (Zhou *et al.*, 1993). Basal IL-6 secretion in cultures of splenic cells and peripheral blood mononuclear cells was not enhanced after repeated footshocks, and mitogen-induced IL-6 production was decreased in cultures of immune cells following stress. Thus, the elevation of plasma IL-6 may reflect a hormonal response of the organism to arousal. Whether psychosocial stress influences the

expression of cytokine mRNA and/or cytokine synthesis in limbic areas is not yet clear. With the exception of Pousset *et al.* (1996) (see also above), who recently observed the induction of IL-6, TNF- $\alpha$  and transforming growth factor  $\beta$  (TGF- $\beta$ ) mRNA expression in cultured rat hippocampal astrocytes by low concentrations ( $10^{-10}$ – $10^{-12}$  M) of serotonin, non-inflammatory stimuli appear to induce IL-1 $\beta$  gene expression in the hypothalamus exclusively (Minami *et al.*, 1991). Endogenous glucocorticoids are known to modulate the central effects of cytokines (Morrow *et al.*, 1993a; Goujon *et al.*, 1995). However, studies addressing the mechanisms and the specific site of action are rather limited to date. Despite such gaps in our knowledge, a considerable number of data describe relations between stress effects including levels of catecholamines in blood and different nuclei of the amygdaloid body on the basis of lesion or stimulation experiments (Henke, 1980a, 1980b, 1985; Johansson *et al.*, 1981; Galeno and Brody, 1983; Morrow *et al.*, 1993b). Electrolytic lesions of the central nucleus of the amygdala attenuated the increase in dopamine turnover in the frontal cortex normally seen following mild stress (Davis *et al.*, 1994). The question of the underlying neural pathways mediating this response is not resolved. However, dopaminergic cells in the ventral tegmental area, which in turn project to the prefrontal cortex may be involved (Maeda and Mogenson, 1981; Wallace *et al.*, 1992). CRH-NE interactions at the central nucleus of the amygdala appear to mediate gastric cytoprotective effects during stress implicating  $\beta$ -adrenoceptor action (Ray *et al.*, 1993). An increase in NE levels also was observed in the BST of rats during 2 hr of immobilization (Pacak *et al.*, 1995). In another study, an increased number of CRH-binding sites was found in the amygdala of tree shrews following psychosocial stress (Fuchs and Flügge, 1994). Kalin *et al.* (1994) have observed that restraint stress increased levels of CRH mRNA in the rat amygdala. However, a more recent study failed to replicate this result, and it was suggested that type or duration of stressors and species differences may account for selective responses of CRH neurons in the amygdala (Makino *et al.*, 1995). Support to this notion came from other studies, indicating that different neuronal networks, particularly dopaminergic activity in the nucleus accumbens, are involved in the adaptation to aversive or pleasurable repeated experiences (Le Moal and Simon, 1991; Imperato *et al.*, 1992). The concept that increased dopamine turnover, metabolism and release in mesocorticolimbic areas is a strain-specific and time-dependent response to stress-inducing stimuli, such as physical restraint, footshock, or social defeat stress, has been experimentally confirmed by a number of investigations (Cabib *et al.*, 1988; Roth *et al.*, 1988; Abercrombie *et al.*, 1989; Puglisi-Allegra *et al.*, 1991; Tidey and Miczek, 1996). Furthermore, it is well established that the hippocampus, particularly the hippocampal cholinergic system, has a crucial role in the processing of stressful stimuli (Finkelstein *et al.*, 1985; Gilad *et al.*, 1985; Tajima *et al.*, 1996). Chronic psychosocial stress also has been found to induce morphological alterations in CA1 and CA3 hippocampal pyramidal

neurons of tree shrews (Fuchs *et al.*, 1995). However, the functional relationship between stress-induced HPA axis activation and abovementioned neuronal systems is still a point of debate. Imperato *et al.* (1991) e.g. have observed in rats that restraint stress and freeing the animals both stimulated release and metabolism of dopamine in the nucleus accumbens and prefrontal cortex, as well as the release of acetylcholine in the hippocampus. After habituation to the respective situation these alterations declined. Corticosterone levels were elevated during the whole period of restraint stress and decreased by freeing the animals. Moreover, this study provides evidence that these dopamine and acetylcholine changes in limbic structures occur independently of HPA axis activation. Thus, different forms of stressful stimulation appear to be processed in limbic brain regions implicating emotional experiences, and selective neurochemical changes in limbic structures in turn may cause selective central and peripheral neuronal, neuroendocrine and, most probably, immune alterations.

A final point to be discussed is that brief or prolonged periods of stress have been shown to influence memory formation in different ways (reviewed by McEwen and Sapolsky, 1995). Considering glutamate receptors as putative candidates involved in learning and memory (Advokat and Pellegrin, 1992; Aigner, 1995), changes in excitatory amino acid receptor binding have been demonstrated in the hippocampus following different qualities of stressful stimulation (single social stress experience, acute tail-shock stress) (Tocco *et al.*, 1991; Krugers *et al.*, 1993). However, contrary results were obtained in response to repeated restraint stress (Watanabe *et al.*, 1995a). Further detailed research is needed to study the implication of glutamate receptors in stress responses and cognition, and future experiments should examine the specific role of these receptors in alterations of immune responses in association with psychosocial stress. Even though, at present, nothing is known about such an interaction, first — albeit indirect — lines of evidence appear to point in this direction. In experiments on embryonic mammalian hippocampal neurons, Mayer and Vyklicky (1989) observed that the T-cell mitogen ConA selectively reduces desensitization of quisqualate receptors. The most compelling evidence for an interaction between limbic structures, the glutamatergic system and the immune system is found in a recent report by Hardin-Pouzet *et al.* (1996). The authors described an enhanced expression of glutamate dehydrogenase in cultured hippocampal astrocytes following treatment with TNF- $\alpha$  or IL-1 $\alpha$ . This enzyme is involved in glutamate metabolism and helps to maintain a low (non-toxic) extracellular glutamate concentration.

#### 4.5. Effects of the Activated Immune System on Limbic Structures

Consistent with the notion of a link between limbic areas and immunity, peripheral endotoxin (LPS) treatment has been found to result in decreased [<sup>125</sup>I]-IL-1 $\alpha$  binding in the hippocampus (Takao *et al.*, 1994) and dentate gyrus (Haour *et al.*, 1990). Furthermore, LPS, IL-1 $\alpha$  or  $\beta$  can increase prosta-

glandin (PG)E2 biosynthesis in the dorsal hippocampus of mice involving IL-1 receptor activation (Weidenfeld *et al.*, 1995), and LPS decreases the affinity of hippocampal MRs (Schöbitz *et al.*, 1994b). The LPS-induced increases in 5-HT metabolism in the hypothalamus and hippocampus can be asymmetrically expressed (Delrue *et al.*, 1994), supporting observations on brain and behavioral lateralization, which have been associated with immune reactivity (review by Neveu, 1992). Central LPS has been found to induce different patterns of cytokine mRNA expression in the brain including the hippocampus (De Simoni *et al.*, 1995). In addition to what has been observed with bacterial endotoxin, muramyl dipeptide, a microbial product with adjuvant activity, which has been shown to be implicated in steep regulation (Krueger and Majde, 1995), can alter neuronal activity in the hypothalamus as well as the hippocampus (Dougherty and Dafny, 1990).

Immunologic reactivity also can interfere indirectly with the production of neurotransmitters, and could have neurobiological as well as psychological consequences within limbic structures. Werner-Felmayer *et al.* (1989) have found that an IFN- $\gamma$ -induced pteridine biosynthesis is associated with an enhanced tryptophan metabolism to form kynurenine. This reduces the availability of 5-HT and may cause the production of metabolites with neurotoxic properties. In this context, a chronically activated immune system, as is observed in HIV patients, has been correlated with reduced serum levels of tryptophan (Fuchs *et al.*, 1990) and with decreased whole blood levels of 5-HT (Launay *et al.*, 1989). Interestingly, both results have been shown to be associated with neurologic/psychiatric symptoms in HIV patients.

Additional evidence that HIV infection affects limbic areas came from studies in rats. The i.c.v. infusion of recombinant gp120, as well as injection directly into the dorsal hippocampus, induced IL-1 activity in the brain (Sundar *et al.*, 1991), whereby the injection into the hippocampus was more effective than i.c.v. infusion. This was associated with an enhanced HPA axis activity, reduced NK cell activity and lymphocyte mitogen responses.

Finally, immune cell-derived cytokines were found to exert effects on growth and survival of neurons. Hippocampal neurons, as well as glial cells, respond to IL-1 stimulation in culture by enhanced production of nerve growth factor (NGF) (Friedman *et al.*, 1990; Spranger *et al.*, 1990) and i.c.v. injection of IL-1 was found to increase NGF mRNA in the hippocampus (Spranger *et al.*, 1990). Recombinant IFN- $\gamma$  has been reported to promote the differentiation of cultured embryonic hippocampal neurons at physiologically relevant concentrations (Barish *et al.*, 1991). Additional lines of evidence indicate that IL-6 acts as a neurotrophic factor (Hama *et al.*, 1989, 1991; Kushima and Hatanaka, 1992; Kushima *et al.*, 1992). Moreover, IL-6 has been recognized to protect cultured rat hippocampal neurons against glutamate neurotoxicity (Yamada and Hatanaka, 1994), which has been suggested to be caused, in part, by NMDA receptor/channel activation (Rothman and Olney, 1987; Yamada and Hatanaka, 1994). This is in agreement with Toulmond *et al.* (1992)'s *in vivo* findings in rats regarding a neuroprotective role of

IL-6 against neurotoxic effects of NMDA on striatal cholinergic neurons. Sarder *et al.* (1996) observed differential effects of IL-2 and IL-6 on neuronal morphology and neurite regeneration in cultured hippocampal cells. IL-2 promoted both elongation and branching of neurites, whereas IL-6 only promoted elongation of neuronal processes. Furthermore, IL-2 significantly increased the number and the length of the branches, which emerge from the proximal part of a damaged neurite, and it also increased the total length and number of non-damaged neurites. In contrast, IL-6 did not show any significant effect on these parameters in laser beam-damaged neurons. In all, the results of the present and preceding studies clearly indicate that immune-cytokines selectively affect neuronal growth, survival and neuritogenesis. On the other hand, direct actions of cytokines, or interactions between cytokines and neurotrophic factors may have detrimental effects on limbic neuronal systems. Besides NGF, one of the best known neurotrophic factors is brain-derived neurotrophic factor (BDNF). Both of them have been considered as agents that presumably act on cholinergic neurons of the basal forebrain (Whittemore and Sieger, 1987; Phillips *et al.*, 1990) and, because of this, probably are able to protect neurons from neurodegenerative diseases. In this context, Lapchak *et al.* (1993) reported that systemic IL-1 $\beta$  decreases BDNF mRNA expression in the rat hippocampal formation, and this mechanism is thought to be involved in the neuropathology of Alzheimer's disease. Chronic treatment with recombinant IL-2 has been used to determine the cytokine effect on the CNS of aged mice as compared to adult animals (Nemni *et al.*, 1992). IL-2-mediated neuronal cell losses and degenerative changes limited to hippocampal regions became apparent in aged mice only, and were correlated with mnemonic dysfunctions.

#### 4.6. Conditioning of Immune Functions — Mediated Via Limbic Areas?

Based on the work of Pavlov (1928), conditioned neutral stimuli, such as saccharin, lithium chloride or environmental stimuli, that were previously paired with an unconditioned stimulus, e.g. an immunosuppressive drug, subsequently are able by themselves to modulate an immune response (conditioned response). A detailed review of this issue is presented in Ader and colleagues' *Psychoneuroimmunology* (1991) — the underlying mechanisms mediating these phenomena remain largely speculative. Endogenous increases of glucocorticoids caused by stressful conditioning paradigms may be involved in conditioned immunosuppression (Kelley *et al.*, 1985; Lysle *et al.*, 1988). However, other studies did not support this hypothesis (Ader *et al.*, 1979; King *et al.*, 1987; Peeke *et al.*, 1987). Endogenous opioids, such as  $\beta$ -endorphin, and the HPA axis, are believed to play an essential role in the conditioned enhancement of NK cell activity (Hsueh *et al.*, 1994, 1995). In line with this view, arecoline, a muscarinic cholinergic agent, which previously has been shown to stimulate HPA activity via central pathways (Calogero *et al.*, 1989), has recently been identified as a drug that serves as an unconditioned stimulus in a Pavlovian

conditioning paradigm (Demissie *et al.*, 1995). The pairing of arecoline with a novel odor (camphor) could be used to trigger a conditioned response, whereby the subsequent exposure to the odor alone was sufficient to raise NK activity. In a related study, centrally injected IFN- $\beta$  also has been found to serve as an unconditioned stimulus for the conditioning of an NK cell response (Solvason *et al.*, 1993); however, the central pathways mediating this effect still remain to be examined. Other data suggest the involvement of endogenous CRH (Perez and Lysle, 1995), and the  $\beta$ -adrenergic system (Lysle *et al.*, 1991; Luecken and Lysle, 1992) in the conditioned modulation of different types of immune responses. Whether and how limbic areas actually participate in the central modulation of conditioned immune responses is not yet formally known. Evidence in favor of this notion came from a study by Pezzone *et al.* (1992). An auditory cue as conditioned stimulus following electric footshock (the unconditioned stimulus) was able to induce Fos expression in different parts of the brain including the medial amygdaloid nucleus. In addition, both of these stimuli were capable to suppress immune functions. Unfortunately, it cannot be excluded that Fos activation in the amygdala just reflects the role of this structure in an (un)conditioned fear response. Thus, these studies should be extended by using non-aversive conditioning paradigms. However, based on the data summarized herein, and in view of the well-known role of limbic areas in learning processes, as well as in the processing of associative emotional factors, we propose that neuronal activities in the hippocampal formation and extended amygdala are critically involved in these phenomena.

## 5. CONCLUSIONS

In this review, we have charted for the first time an integrated view of the neuroanatomy of psychoimmunology based on the description of afferent/efferent pathways in the neuroimmune dialog, as well as on the major functions of limbic areas and their putative roles in brain-immune communications. It stands to reason that psychoneuroimmunology cannot provide a universal explanation of a given disease, but it may enhance our understanding of psychosomatic aspects of immune disorders, such as immune deficiency states or autoimmune diseases. Additionally, it may open up new pathophysiological concepts of different types of depression and dementia. However, considering the conspicuous complexity of neuronal networks and neuroendocrine mechanisms on the one hand, and the complexity of immune functions on the other, it is not surprising that the results obtained are diverse and often contradictory. We can only conclude that (i) humoral and neural, most probably vagal, afferents link the peripheral immune system with the brain; (ii) humoral and neural efferents, such as the HPA axis, a number of neuropeptides and autonomic pathways link the CNS with the peripheral immune system; (iii) limbic structures represent the neuroanatomical substrates for psychoimmunomodulation as they serve as superior regulators of neuroendocrine and

autonomic centers allowing an individual and selective modulation of learning processes, memory, emotional behavior, stress and immune responses.

Given these results, immune dysfunctions may (i) arise from a dysregulation of the parasympathetic division of the nervous system; (ii) from an MR/GR imbalance in the hippocampus leading either to an enhanced or an impaired HPA activity; (iii) from changes in number, affinity or signaling of cytokine receptors in the CNS; and (iv) from alterations in release and turnover of all the neurotransmitter material mentioned, ranging from NE to excitatory amino acids. More detailed knowledge of the pharmacology, particularly of the "newcomers" of transmitters, as well as their physiological roles in brain and immune functions, will be required to clarify further the extensive network of brain-immune interactions. Additional studies at the molecular level will be needed to determine precisely cytokine effects on neurotransmission. Cholinergic and GABAergic responses should be studied in more detail. Finally, a potential field of interest would be the investigation of glutamatergic mechanisms in neuroimmunology. Such investigations should lead to a better understanding of neuroimmunomodulation at limbic structures to provide the scientific basis to Juvenal's "*Orandum est ut sit mens sana in corpore sano*".

**Acknowledgements**—The research from our laboratory reported in this article was supported by the Austrian Science Foundation, projects 7509, 7038, and 9925. We wish to thank Drs Peter Liebmann and Peter Marquardt for their review of this manuscript. Thanks also to Mr Heinrich Haas for his advice on the illustrations.

## REFERENCES

- Abe, K., Ishiyama, J. and Saito, H. (1992) Effects of epidermal growth factor and basic fibroblast growth factor on generation of long-term potentiation in the dentate gyrus of fimbria-fornix-lesioned rats. *Brain Res.* **593**, 335–338.
- Abelson, J. L. and Curtis, G. C. (1996) Hypothalamic pituitary-adrenal axis activity in panic disorder. *Arch. Gen. Psychiat.* **53**, 323–331.
- Abercrombie, E. D., Keefe, K. A., DiFrischia, D. F. and Zigmond, M. J. (1989) Differential effects of stress on *in vivo* dopamine release in striatum, nucleus accumbens, and medial prefrontal cortex. *J. Neurochem.* **52**, 1655–1658.
- Adamec, R. E. and Morgan, H. D. (1994) The effect of kindling of different nuclei in the left and right amygdala on anxiety in the rat. *Physiol. Behav.* **55**, 1–12.
- Ader, R., Cohen, N. and Grota, L. J. (1979) Adrenal involvement in conditioned immunosuppression. *Int. J. Immunopharmacol.* **1**, 141–145.
- Ader, R., Felten, D. L. and Cohen, N. (1991) *Psychoneuroimmunology*, second edition. Academic Press: New York.
- Ader, R., Cohen, N. and Felten, D. (1995) Psychoneuroimmunology: interactions between the nervous system and the immune system. *Lancet* **345**, 99–103.
- Advokat, C. and Pellegrin, A. I. (1992) Excitatory amino acids and memory — evidence from research on Alzheimer's disease and behavioural pharmacology. *Neurosci. Behav. Rev.* **16**, 13–24.
- Aggleton, J. P. (1986) A description of the amygdalo-hippocampal interconnections in the macaque monkey. *Exp. Brain Res.* **64**, 515–526.
- Agius, M. A., Checinski, M. E., Richman, D. P. and Chelmicka-Schorr, E. (1987) Sympathectomy enhances the severity of experimental autoimmune myasthenia gravis (EAMG). *J. Neuroimmunol.* **16**, 11–12.
- Agro, A. and Stanis, A. M. (1995) Neuroimmunomodulation: classical and non-classical cellular activation. *Adv. Neuroimmunol.* **5**, 311–319.
- Aigner, T. G. (1995) Pharmacology of memory: cholinergic glutamatergic interactions. *Curr. Opin. Neurobiol.* **5**, 155–160.
- Akira, S., Hirano, T., Taga, T. and Kishimoto, T. (1990) Biology of multifunctional cytokines: IL 6 and related molecules (IL 1 and TNF). *FASEB J.* **4**, 2860–2867.
- Alexander, G. E., DeLong, M. R. and Strick, P. (1989) Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu. Rev. Neurosci.* **9**, 357–381.
- Amaral, D. G. (1986) Amygdalohippocampal and amygdalocortical projections in the primate brain. *Adv. Exp. Med. Biol.* **203**, 3–17.
- Anisman, H., Kokkinidis, L. and Merali, Z. (1996) Interleukin-2 decreases accumbal dopamine efflux and responding for rewarding lateral hypothalamic stimulation. *Brain Res.* **731**, 1–11.
- Antonica, A., Ayroldi, E., Magni, F. and Paolucci, N. (1996) Lymphocyte traffic changes induced by monolateral vagal denervation in mouse thymus and peripheral organs. *J. Neuroimmunol.* **64**, 115–122.
- Arai, A. and Lynch, G. (1996) Response to repetitive stimulation of AMPA receptors in patches excised from fields CA1 and CA3 of the hippocampus. *Brain Res.* **716**, 202–206.
- Araujo, D. M., Lapchak, P. A., Collier, B. and Quirion, R. (1989) Localization of interleukin-2 immunoreactivity and interleukin-2 receptors in the rat brain: interaction with the cholinergic system. *Brain Res.* **498**, 257–266.
- Avishai-Eliner, S., Yi, S.-J. and Baram, T. Z. (1996) Developmental profile of messenger RNA for the corticotropin-releasing hormone receptor in the rat limbic system. *Dev. Brain Res.* **91**, 159–163.
- Ballou, L. R. (1992) Sphingolipids and cell function. *Immunol. Today* **13**, 339–341.
- Barish, M. E., Mansdorf, N. B. and Raissdana, S. S. (1991) Gamma-interferon promotes differentiation of cultured cortical and hippocampal neurons. *Dev. Biol.* **144**, 412–423.
- Bashir, Z. I., Alford, S., Davies, S. N., Randall, A. D. and Collingridge, G. L. (1991) Long-term potentiation of NMDA receptor-mediated synaptic transmission in the hippocampus. *Nature* **349**, 156–158.
- Beaulieu, S., Di Paolo, T. and Barden, N. (1986) Control of ACTH secretion by the central nucleus of the amygdala: implication of the serotonergic system and its relevance to glucocorticoid delayed negative feedback mechanism. *Neuroendocrinology* **44**, 247–254.
- Beaulieu, S., Di Paolo, T., Cote, J. and Barden, N. (1987) Participation of the central amygdaloid nucleus in the response of adrenocorticotropin secretion to immobilization stress: opposing roles of the noradrenergic and dopaminergic systems. *Neuroendocrinology* **45**, 37–46.
- Beaulieu, S., Pelletier, G., Vaudry, H. and Barden, N. (1989) Influence of the central nucleus of the amygdala on the content of corticotropin-releasing factor in the median eminence. *Neuroendocrinology* **49**, 255–261.
- Behbehani, M. M. (1995) Functional characteristics of the midbrain periaqueductal gray. *Prog. Neurobiol.* **46**, 575–605.
- Bellinger, D. L., Lorton, D., Romano, T. D., Olschowka, J. A., Felten, S. Y. and Felten, D. L. (1990) Neuropeptide innervation of lymphoid organs. *Ann. NY Acad. Sci.* **594**, 17–33.
- Bellinger, F. P., Madamba, S. and Siggins, G. R. (1993) Interleukin-1 $\beta$  inhibits synaptic strength and long-term potentiation in the rat CA1 hippocampus. *Brain Res.* **628**, 227–234.
- Bergquist, J., Tarkowski, A., Ekman, R. and Ewing, A. (1994) Discovery of endogenous catecholamines in lymphocytes and evidence for catecholamine regulation of lymphocyte function via an autocrine loop. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 12912–12916.
- Berkenbosch, F., Van Oers, J., Del Rey, A., Tilders, F. and Besedovsky, H. O. (1987) Corticotropin-releasing factor-producing neurons in the rat activated by interleukin-1. *Science* **238**, 524–526.
- Besedovsky, H. O., Sorkin, E., Keller, M. and Müller, J. (1975) Changes in blood hormone levels during the immune response. *Proc. Soc. Exp. Biol. Med.* **150**, 466–470.
- Besedovsky, H., Sorkin, E., Felix, D. and Haas, H. (1977) Hypothalamic changes during the immune response. *Eur. J. Immunol.* **7**, 323–325.
- Besedovsky, H. O., Del Rey, A. and Sorkin, E. (1979) Antigenic competition between horse and sheep red blood cells is a

- hormone-dependent phenomenon. *Clin. Exp. Immunol.* **37**, 106-113.
- Besedovsky, H. O., Del Rey, A., Sorkin, E., Da Prada, M., Burri, R. and Honegger, C. (1983) The immune response evokes changes in brain noradrenergic neurons. *Science* **221**, 564-566.
- Besedovsky, H. O., Del Rey, A., Sorkin, E. and Dinarello, C. A. (1986) Immunoregulatory feedback between interleukin-1 and glucocorticoid hormones. *Science* **233**, 652-654.
- Bienenstock, J., Tomioka, M., Matsuda, H., Stead, R. H., Quinonez, G., Simon, G. T., Coughlin, M. D. and Denburg, J. A. (1987) The role of mast cells in inflammatory processes: evidence for nerve-mast cell interactions. *Int. Arch. Allergy Immunol.* **82**, 238-243.
- Bileviciute, I., Lundeberg, T., Ekblom, A. and Theodorsson, E. (1993) Bilateral changes of substance P, neurokinin A, calcitonin gene-related peptide- and neuropeptide Y-like immunoreactivity in rat knee joint synovial fluid during acute monoarthritis. *Neurosci. Lett.* **153**, 37-40.
- Bileviciute, I., Stenfors, C., Theodorsson, E., Beckman, M. and Lundeberg, T. (1995) Significant changes in neuropeptide concentrations in the brain of normotensive (WKY) and spontaneously hypertensive (SHR) rats following knee joint monoarthritis. *Brain Res.* **704**, 71-78.
- Blalock, J. E. (1992) Production of peptide hormones and neurotransmitters by the immune system (Eds K. Ishizaka, P. J. Lachmann and B. H. Waksman). *Chem. Immunol. (Neuroimmunoenocrinol.)* **52**, 1-24.
- Blalock, J. E. (1994) The syntax of immune-neuroendocrine communication. *Immunol. Today* **15**, 504-511.
- Blatteis, C. M. (1992) Role of the OVLTL in the febrile response to circulating pyrogens. *Prog. Brain Res.* **91**, 409-412.
- Bliss, T. V. P. and Collingridge, G. L. (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* **361**, 31-39.
- Blum, A. M., Metwali, A., Mathew, R. C., Cook, G., Elliott, D. and Weinstock, J. V. (1992) Granuloma T lymphocytes in murine schistosomiasis mansoni have somatostatin receptors and respond to somatostatin with decreased IFN- $\gamma$  secretion. *J. Immunol.* **149**, 3621-3626.
- Blümcke, I., Wolf, H. K., Hof, P. R., Morrison, J. H. and Wiestler, O. D. (1995) Regional distribution of the AMPA glutamate receptor subunits GluR2(4) in human hippocampus. *Brain Res.* **682**, 239-244.
- Bluthé, R.-M., Walter, V., Parnet, P., Layé, S., Lestace, J., Verrier, D., Poole, S., Stenning, B. E., Kelley, K. W. and Dantzer, R. (1994) Lipopolysaccharide induces sickness behaviour in rats by a vagal mediated mechanism. *C.R. Acad. Sci. Paris Ser. III* **317**, 499-503.
- Boyd, R. L., Tucek, C. L., Godfrey, D. I., Izon, D. J., Wilson, T. J., Davidson, N. J., Bean, A. G. D., Ladyman, H. M., Ritter, M. A. and Hugo, P. (1993) The thymic microenvironment. *Immunol. Today* **14**, 445-459.
- Brinton, R. D., Monreal, A. W. and Fernandez, J. G. (1994) Vasopressin-induced neurotrophism in cultured hippocampal neurons via V1 receptor activation. *J. Neurobiol.* **25**, 380-394.
- Brooks, W. H., Cross, R. J., Roszman, T. L. and Markesbery, W. R. (1982) Neuroimmunomodulation: neural anatomical basis for impairment and facilitation. *Ann. Neurol.* **12**, 56-61.
- Bullier, J. and Nowak, L. G. (1995) Parallel versus serial processing: new vistas on the distributed organization of the visual system. *Curr. Opin. Neurobiol.* **5**, 497-503.
- Bullock, K. and Moore, R. Y. (1981) Innervation of the thymus gland by brain stem and spinal cord in mouse and rat. *Am. J. Anat.* **162**, 157-166.
- Burgard, E. C. and Sarvey, J. M. (1990) Muscarinic receptor activation facilitates the induction of long-term potentiation (LTP) in the rat dentate gyrus. *Neurosci. Lett.* **116**, 34-39.
- Cabib, S., Kempf, E., Schleaf, C., Oliverio, A. and Puglisi-Allegra, S. (1988) Effects of immobilization stress on dopamine and its metabolites in different brain areas of the mouse: role of genotype and stress duration. *Brain Res.* **441**, 153-160.
- Calogero, A. E., Kamilaris, T. C., Gomez, M. T., Johnson, E. O., Tartaglia, M. E., Gold, P. W. and Chrousos, G. P. (1989) The muscarinic cholinergic agonist arecoline stimulates the rat hypothalamic-pituitary-adrenal axis through a centrally mediated corticotropin-releasing hormone-dependent mechanism. *Endocrinology* **125**, 2445-2453.
- Carlson, S. L., Felten, D. L., Livnat, S. and Felten, S. Y. (1987) Alterations of monoamines in specific central autonomic nuclei following immunization in mice. *Brain Behav. Immun.* **1**, 52-63.
- Carr, D. J. J. (1992) Neuroendocrine peptide receptors on cells of the immune system (Eds K. Ishizaka, P. J. Lachmann and B. H. Waksman). *Chem. Immunol. (Neuroimmunoenocrinol.)* **52**, 84-105.
- Carroll, B. J., Curtis, G. C. and Mendels, J. (1976) Neuroendocrine regulation in depression I. Limbic system-adrenocortical dysfunction. *Arch. Gen. Psychiat.* **33**, 1039-1044.
- Chelmicka-Schorr, E., Checinski, M. and Arnason, B. G. W. (1988) Chemical sympathectomy augments the severity of experimental allergic encephalomyelitis. *J. Neuroimmunol.* **17**, 347-350.
- Chelmicka-Schorr, E., Kwasniewski, M. N., Thomas, B. E. and Arnason, B. G. W. (1989) The  $\beta$ -adrenergic agonist isoproterenol suppresses experimental allergic encephalomyelitis in Lewis rats. *J. Neuroimmunol.* **25**, 203-207.
- Cherkaoui, J., Mayo, W., Neveu, P. J., Kelley, K. W., Vitiello, S., Le Moal, M. and Simon, H. (1990) The nucleus basalis is involved in brain modulation of the immune system in rats. *Brain Res.* **516**, 345-348.
- Chrousos, G. P. (1995) The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation. *New Engl. J. Med.* **332**, 1351-1362.
- Clark, A. S. and Cotman, C. W. (1992) Adrenal hormone effects on hippocampal excitatory amino acid binding. *Brain Res.* **585**, 161-168.
- Coirini, H., Margariños, A. M., De Nicola, A. F., Rainbow, T. C. and McEwen, B. S. (1985) Further studies of brain aldosterone binding sites employing new mineralocorticoid and glucocorticoid receptor markers *in vitro*. *Brain Res.* **361**, 212-216.
- Collingridge, G. L. and Davies, S. N. (1989) NMDA receptors and long-term potentiation in the hippocampus. In: *The NMDA Receptor*, pp. 123-135. Eds J. C. Watkins and G. L. Oxford University Press: Oxford, U.K.
- Costa, P., Traver, D. J., Auger, C. B. and Costa, L. G. (1994) Expression of cholinergic muscarinic receptor subtypes mRNA in rat blood mononuclear cells. *Immunopharmacology* **28**, 113-123.
- Croiset, G., Heijnen, C. J., Van der Wal, W. E., De Boer, S. F. and De Wied, D. (1990) A role for the autonomic nervous system in modulating the immune response during mild emotional stimuli. *Life Sci.* **46**, 419-425.
- Cross, R. J., Brooks, W. H., Roszman, T. L. and Markesbery, W. R. (1982) Hypothalamic-immune interactions. Effect of hypophysectomy on neuroimmunomodulation. *J. Neurol. Sci.* **53**, 557-566.
- Cui, Y., Lee, T. F. and Wang, L. C. (1993) Thermoregulatory responses following injection of 5-hydroxytryptamine into the septohippocampal complex in rats. *Pharmacol. Biochem. Behav.* **45**, 935-939.
- Cullinan, W. E., Herman, J. P. and Watson, S. J. (1993) Ventral subicular interaction with the hypothalamic paraventricular nucleus: evidence for a relay in the bed nucleus of the stria terminalis. *J. Comp. Neurol.* **332**, 1-20.
- Dalmaz, C., Introicollison, J. B. and Mcgaugh, J. L. (1993) Noradrenergic and cholinergic interactions in the amygdala and the modulation of memory storage. *Behav. Brain Res.* **58**, 167-174.
- Dantzer, R. and Kelley, K. W. (1989) Stress and immunity: an integrated view of relationships between the brain and the immune system. *Life Sci.* **44**, 1995-2008.
- D'Arcangelo, G., Grassi, F., Ragozzino, D., Santoni, A., Tancredi, V. and Eusebi, F. (1991) Interferon inhibits synaptic potentiation in rat hippocampus. *Brain Res.* **564**, 245-248.
- Davis, M. (1992) The role of the amygdala in fear and anxiety. *Annu. Rev. Neurosci.* **15**, 353-375.
- Davis, M., Hitchcock, J. M., Bowers, M. B., Berridge, C. W., Melia, K. R. and Roth, R. H. (1994) Stress-induced activation of prefrontal cortex dopamine turnover: blockade by lesions of the amygdala. *Brain Res.* **664**, 207-210.
- Day, N. C., Williams, T. L., Ince, P. G., Kamboj, R. K., Lodge, D. and Shaw, P. J. (1995) Distribution of AMPA-selective glutamate receptor subunits in the human hippocampus and cerebellum. *Mol. Brain Res.* **31**, 17-32.
- DeGiorgi, L. and Altomare, D. (1992) Modification of the immune system following mental and physical stress. *Bull. Inst. Pasteur* **90**, 99-107.
- De Kloet, E. R. (1991) Brain corticosteroid receptor balance and homeostatic control. *Front. Neuroendocrinol.* **12**, 95-164.

- De Kloet, E. R., Oitzl, M. S. and Schöbitz, B. (1994) Cytokines and the brain corticosteroid receptor balance: relevance to pathophysiology of neuroendocrine immune communication. *Psychoneuroendocrinology* **19**, 121-134.
- Deleplanque, B., Vitiello, S., Le Moal, M. and Neveu, P. J. (1994) Modulation of immune reactivity by unilateral striatal and mesolimbic dopaminergic lesions. *Neurosci. Lett.* **166**, 216-220.
- Delrue, C., Deleplanque, B., Rouge-Pont, F., Vitiello, S. and Neveu, P. J. (1994) Brain monoaminergic, neuroendocrine, and immune responses to an immune challenge in relation to brain and behavioral lateralization. *Brain Behav. Immun.* **8**, 137-152.
- Demissie, S., Rogers, C. F., Hiramoto, N. S., Ghanta, V. K. and Hiramoto, R. N. (1995) Arecoline a muscarinic cholinergic agent conditions central pathways that modulate natural killer cell activity. *J. Neuroimmunol.* **59**, 57-63.
- De Simoni, M. G., De Luigi, A., Gemma, C., Faggioni, R., Gatti, S., Manfridi, A., Sironi, M. and Ghezzi, P. (1992) Central nervous system-mediated induction of systemic IL-6 and TNF. In: *Pathophysiology and Pharmacology of Cytokines*, pp. 143-148. Eds A. Mantovani and P. Ghezzi. Biomedical Press: New York.
- De Simoni, M. G., De Luigi, A., Gemma, C., Sironi, M., Manfridi, A. and Ghezzi, P. (1993) Modulation of systemic interleukin-6 induction by central interleukin-1. *Am. J. Physiol.* **265**, R739-R742.
- De Simoni, M. G., Del Bo, R., De Luigi, A., Simard, S. and Forloni, G. (1995) Central endotoxin induces different patterns of interleukin (IL)-1 $\beta$  and IL-6 messenger ribonucleic acid expression and IL-6 secretion in the brain and periphery. *Endocrinology* **136**, 897-902.
- De Vries, G. J., Buijs, R. M., Van Leeuwen, F. W., Caffé, A. R. and Swaab, D. F. (1985) The vasopressin innervation of the brain in normal and castrated rats. *J. Comp. Neurol.* **233**, 236-254.
- De Vries, H. E., Blom-Roosemalen, M. C. M., van Oosten, M., de Boer, A. G., van Berkel, T. J. C., Breimer, D. D. and Kuiper, J. (1996) The influence of cytokines on the integrity of the blood brain barrier *in vitro*. *J. Neuroimmunol.* **64**, 37-43.
- Dhabhar, F. S., Miller, A. H., McEwen, B. S. and Spencer, R. L. (1995a) Differential activation of adrenal steroid receptors in neural and immune tissues of Sprague Dawley, Fischer 344, and Lewis rats. *J. Neuroimmunol.* **56**, 77-90.
- Dhabhar, F. S., Miller, A. H., McEwen, B. S. and Spencer, R. L. (1995b) Effects of stress on immune cell distribution. Dynamics and hormonal mechanisms. *J. Immunol.* **154**, 5511-5527.
- Donaldson, L. F., Harmar, A. J., McQueen, D. S. and Seckl, J. R. (1992) Increased expression of preprotachykinin, calcitonin gene-related peptide, but not vasoactive intestinal peptide messenger RNA in dorsal root ganglia during the development of adjuvant monoarthritis in the rat. *Mol. Brain Res.* **16**, 143-149.
- Dougherty, P. M. and Dafny, N. (1990) Muramyl-dipeptide, a macrophage-derived cytokine, alters neuronal activity in hypothalamus and hippocampus but not in the dorsal raphe/periaqueductal gray of rats. *J. Neuroimmunol.* **28**, 201-208.
- Dunn, A. J. (1992) The role of interleukin-1 and tumor necrosis factor- $\alpha$  in the neurochemical and neuroendocrine responses to endotoxin. *Brain Res. Bull.* **29**, 807-812.
- Dunn, J. D. (1987) Plasma corticosterone responses to electrical stimulation of the bed nucleus of the stria terminalis. *Brain Res.* **407**, 327-331.
- Durkin, T. P. (1994) Spatial working memory over long retention intervals: dependence on sustained cholinergic activation in the septohippocampal or nucleus basalis magnocellularis-cortical pathways?. *Neuroscience* **62**, 681-693.
- Duvaux-Miret, O., Stefano, G. B., Smith, E. M., Dissous, C. and Capron, A. (1992) Immunosuppression in the definitive and intermediate hosts of the human parasite *Schistosoma mansoni* by release of immunoreactive neuropeptides. *Proc. Natl. Acad. Sci. U.S.A.* **89**, 778-781.
- Elkabar, D. R., Wyatt, M. E., Vellucci, S. V. and Herbert, J. (1990) The effects of separate or combined infusions of corticotropin-releasing factor and vasopressin either intraventricularly or into the amygdala on aggressive and investigative behavior in the rat. *Regul. Peptides* **28**, 199-214.
- Eskay, R. L., Elden, L. E. and Hsu, C.-M. (1992) Interleukin-1 $\alpha$  and tumor necrosis factor- $\alpha$  differentially regulate enkephalin, vasoactive intestinal polypeptide, neurotensin, and substance P biosynthesis in chromaffin cells. *Endocrinology* **130**, 2252-2258.
- Evans, D. L., Folds, J. D., Petitto, J. M., Golden, R. N., Pedersen, C. A., Corrigan, M., Gilmore, J. H., Silva, S. G., Quade, D. and Ozer, H. (1992) Circulating natural killer cell phenotypes in men and women with major depression. *Arch. Gen. Psychiat.* **49**, 388-395.
- Evans, S. W., Beckner, S. K. and Farrar, W. L. (1987) Stimulation of specific GTP binding and hydrolysis activities in lymphocyte membrane by interleukin-2. *Nature* **325**, 166-168.
- Farrar, W. L. and Andersen, W. B. (1985) Interleukin-2 stimulates association of protein kinase C with plasma membrane. *Nature* **315**, 233-235.
- Farrar, W. L., Hill, J. M., Harel-Bellan, A. and Vinocour, M. (1987) The immune logical brain. *Immunol. Rev.* **100**, 361-378.
- Fatani, J. A., Quayyum, M. A., Mehta, L. and Singh, U. (1986) Parasympathetic innervation of the thymus: a histochemical and immunocytochemical study. *J. Anat.* **147**, 115-119.
- Fecho, K., Maslonek, K. A., Dykstra, L. A. and Lysle, D. T. (1996) Assessment of the involvement of central nervous system and peripheral opioid receptors in the immunomodulatory effects of acute morphine treatment in rats. *J. Pharmacol. Exp. Ther.* **276**, 626-636.
- Feldman, S. and Weidenfeld, J. (1993) The dorsal hippocampus modifies the negative effect of glucocorticoids on the adrenocortical and median eminence CRF-41 responses to photic stimulation. *Brain Res.* **614**, 227-232.
- Feldman, S., Conforti, N. and Saphier, D. (1990) The preoptic area and bed nucleus of the stria terminates are involved in the effects of the amygdala on adrenocortical secretion. *Neuroscience* **37**, 775-779.
- Feldman, S., Conforti, N., Itzik, A. and Weidenfeld, J. (1994) Differential effect of amygdaloid lesions on CRF-41, ACTH and corticosterone responses following neural stimuli. *Brain Res.* **658**, 21-26.
- Felsner, P., Hofer, D., Rinner, I., Mangge, H., Gruber, M., Korsatko, W. and Schauenstein, K. (1992) Continuous *in vivo* treatment with catecholamines suppresses *in vitro* reactivity of rat peripheral blood T-lymphocytes via  $\alpha$ -mediated mechanisms. *J. Neuroimmunol.* **37**, 47-57.
- Felsner, P., Hofer, D., Rinner, I., Porta, S., Korsatko, W. and Schauenstein, K. (1995) Adrenergic suppression of peripheral blood T cell reactivity in the rat is due to activation of peripheral  $\alpha_2$ -receptors. *J. Neuroimmunol.* **57**, 27-34.
- Felten, D. L., Felten, S. Y., Bellinger, D. L., Carlson, S. L., Ackerman, K. D., Madden, K. S., Olshchowa, J. A. and Livnat, S. (1987) Noradrenergic sympathetic neural interactions with the immune system: structure and function. *Immunol. Rev.* **100**, 225-260.
- Felten, D. L., Felten, S. Y., Bellinger, D. L. and Lorton, D. (1992) Noradrenergic and peptidergic innervation of secondary lymphoid organs: role in experimental rheumatoid arthritis. *Eur. J. Clin. Invest.* **22**, Suppl. 1, 37-41.
- Felten, S. Y. and Olshchowa, J. (1987) Noradrenergic sympathetic innervation of the spleen: II. Tyrosine hydroxylase (TH)-positive nerve terminals form synaptic-like contacts on lymphocytes in the splenic white pulp. In: *Neuroimmunomodulation*, pp. 70-74. Eds J. R. Perez-Polo, K. Bulloch, R. H. Angeletti, G. A. Hashim and J. de Vellis. Alan R. Liss, Inc.: New York.
- Finkelstein, Y., Koffler, B., Rabey, J. M. and Gilad, G. M. (1985) Dynamic of cholinergic synaptic mechanisms in the rat hippocampus after stress. *Brain Res.* **343**, 314-319.
- Fleshner, M., Goehler, L. E., Hermann, J., Reltou, J. K., Maier, S. F. and Watkins, L. R. (1995) Interleukin-1 $\beta$  induced corticosterone elevation and hypothalamic NE depletion is vagally mediated. *Brain Res. Bull.* **37**, 605-610.
- Frei, K., Bodmer, S., Schwerdel, C. and Fontana, A. (1985) Astrocytes of the brain synthesize interleukin-3-like factors. *J. Immunol.* **135**, 4044-4047.
- Freidin, M., Bennett, M. V. L. and Kessler, J. A. (1992) Cultured sympathetic neurons synthesize and release the cytokine interleukin 1 $\beta$ . *Proc. Natl. Acad. Sci. U.S.A.* **89**, 10440-10443.
- Fricchione, G. L. and Stefano, G. P. (1994) The stress response and autoimmunoregulation. *Adv. Neuroimmunol.* **4**, 13-27.
- Friedman, W. J., Larkfors, L., Ayer-LeLievre, C., Ebendal, T., Olson, L. and Persson, H. (1990) Regulation of B-nerve growth factor expression by inflammatory mediators in hippocampal cultures. *J. Neurosci.* **Res. **27**, 374-388.**
- Fuchs, D., Moeller, A. A., Reibnegger, G., Stoeckle, E., Werner, E. R. and Wächter, H. (1990) Decreased serum tryptophan in

- patients with HIV-1 infection correlates with increased serum neopterin and with neurologic/psychiatric symptoms. *J. Acq. Immun. Def. Synd.* **3**, 873-876.
- Fuchs, E. and Flügge, G. (1994) Modulation of binding sites for corticotropin-releasing hormone by chronic psychosocial stress. *Psychoneuroendocrinology* **20**, 33-51.
- Fuchs, E., Uno, H. and Flügge, G. (1995) Chronic psychosocial stress induces morphological alterations in hippocampal pyramidal neurons of the tree shrew. *Brain Res.* **673**, 275-282.
- Fukuoka, H., Kawatani, M., Hisamitsu, T. and Takeshige, C. (1994) Cutaneous hyperalgesia induced by peripheral injection of interleukin-1 $\beta$  in the rat. *Brain Res.* **657**, 133-140.
- Gadient, R. A. and Otten, U. (1994) Expression of interleukin-6 (IL-6) and interleukin-6 receptor (IL-6R) mRNAs in rat brain during postnatal development. *Brain Res.* **637**, 10-14.
- Gadient, R. A. and Otten, U. (1996) Postnatal expression of interleukin-6 (IL-6) and IL-6 receptor (IL-6R) mRNAs in rat sympathetic and sensory ganglia. *Brain Res.* **724**, 41-46.
- Galeno, T. M. and Brody, M. J. (1983) Hemodynamic responses to amygdaloid stimulation in spontaneously hypertensive rats. *J. Physiol.* **245**, R281-R286.
- Gasbarri, A., Verney, C., Innocenzi, R., Campana, E. and Pacitti, C. (1994) Mesolimbic dopaminergic neurons innervating the hippocampal formation in the rat: a combined retrograde tracing and immunohistochemical study. *Brain Res.* **668**, 71-79.
- Gaykema, R. P. A., Dijkstra, I. and Tilders, F. J. H. (1995) Subdiaphragmatic vagotomy suppresses endotoxin-induced activation of hypothalamic corticotropin-releasing hormone neurons and ACTH secretion. *Endocrinology* **136**, 4717-4720.
- Ghanta, N. R., Haseman, J. K., Grumbine, S., Crawford, D. D. and Eustis, S. L. (1990) Growth, body weight, survival, and tumor trends in F344/N rats during an eleven-year period. *Toxicol. Pathol.* **18**, 61-70.
- Gilad, G. M., Mahom, B. D., Finkelstein, Y., Koffler, B. and Gilad, V. (1985) Stress-induced activation of the hippocampal cholinergic system and the pituitary-adrenocortical axis. *Brain Res.* **347**, 404-408.
- Gobert, A., Widdowson, P., Renouard, A., Rivet, J. M., Bervoets, K. and Millan, R. J. (1993) Multiple  $\alpha_2$ -adrenoceptor subtypes: an examination of the subtype modulating noradrenaline synthesis in rat hippocampus. *J. Neurochem.* **61S**, S25A.
- Goehler, L. E., Relton, J., Maier, S. F. and Watkins, L. R. (1994) Biotinylated interleukin-1 receptor antagonist (IL-1ra) labels paraventricular nuclei in the rat liver hilus and hepatic vagus. *Proc. Soc. Neurosci. Abstr.* **20**, 956.
- Goehler, L. E., Busch, C. R., Tartaglia, N., Relton, J., Sisk, D., Maier, S. F. and Watkins, L. R. (1995) Blockade of cytokine induced conditioned taste aversion by subdiaphragmatic vagotomy: further evidence for vagal mediation of immune-brain communication. *Neurosci. Lett.* **185**, 163-166.
- Goetzl, E. J. and Sreedharan, S. P. (1992) Mediators of communication and adaptation in the neuroendocrine and immune system. *FASEB J.* **6**, 2646-2652.
- Goh, J. W. and Pennefather, P. S. (1989) A pertussis toxin-sensitive G protein in hippocampal long term potentiation. *Science* **244**, 980-983.
- Gordon, M. A., Cohen, J. J. and Wilson, B. (1978) Muscarinic cholinergic receptors in murine lymphocytes: demonstration by direct binding. *Proc. Natl. Acad. Sci. U.S.A.* **75**, 2902-2904.
- Gottschall, P. E., Komaki, G. and Arimura, A. (1992a) Interleukin-1 beta activation of central nervous system. In: *Interleukin-1 in the Brain*, pp. 27-50. Eds Rothwell and Dantzer. Pergamon Press: New York.
- Gottschall, P. E., Komaki, G. and Arimura, A. (1992b) Increased circulating interleukin-1, and interleukin-6 after intracerebroventricular injection of lipopolysaccharide. *Neuroendocrinology* **56**, 935-938.
- Goujon, E., Parnet, P., Cremona, S. and Dantzer, R. (1995) Endogenous glucocorticoids down regulate central effects of interleukin-1 $\beta$  on body temperature and behaviour in mice. *Brain Res.* **702**, 173-180.
- Gould, E., Woolley, C. S. and McEwen, B. S. (1991) The hippocampal formation: morphological changes induced by thyroid, gonadal and adrenal hormones. *Psychoneuroendocrinology* **16**, 67-84.
- Gray, T. S., Piechowski, R. A., Yracheta, J. M., Rittenhouse, P. A., Bethea, C. L. and Van de Kar, L. D. (1993) Ibotenic acid lesions in the bed nucleus of the stria terminalis attenuate conditioned stress-induced increases in prolactin ACTH and corticosterone. *Neuroendocrinology* **57**, 517-524.
- Grossberg, S. and Merrill, J. W. L. (1992) A neural network model of adaptively timed reinforcement learning and hippocampal dynamics. *Cogn. Brain Res.* **1**, 3-38.
- Gutierrez, E. G., Banks, W. A. and Kastin, A. J. (1993) Murine tumor necrosis factor alpha is transported from blood to brain in the mouse. *J. Neuroimmunol.* **47**, 169-176.
- Gutierrez, E. G., Banks, W. A. and Kastin, A. J. (1994) Blood-borne interleukin-1 receptor antagonist crosses the blood-brain barrier. *J. Neuroimmunol.* **55**, 153-160.
- Hama, T., Miyamoto, M., Tsukui, H., Nishio, C. and Hatanaka, H. (1989) Interleukin-6 as a neurotrophic factor for promoting the survival of cultured basal forebrain cholinergic neurons from postnatal rats. *Neurosci. Lett.* **104**, 340-344.
- Hama, T., Kushima, Y., Miyamoto, M., Kubota, M., Takei, N. and Hatanaka, H. (1991) Interleukin-6 improves the survival of mesencephalic catecholaminergic and septal cholinergic neurons from postnatal, two-week-old rats in cultures. *Neuroscience* **40**, 445-452.
- Hamilton, M. E. and Freeman, A. S. (1995) Effects of administration of cholecystokinin into the VTA on DA overflow in nucleus accumbens and amygdala of freely moving rats. *Brain Res.* **688**, 134-142.
- Han, Y., Shaikh, M. B. and Siegel, A. (1996a) Medial amygdaloid suppression of predatory attack behavior in the cat: I. Role of a substance P pathway from the medial amygdala to the medial hypothalamus. *Brain Res.* **716**, 59-71.
- Han, Y., Shaikh, M. B. and Siegel, A. (1996b) Medial amygdaloid suppression of predatory attack behavior in the cat: II. Role of a GABAergic pathway from the medial to the lateral hypothalamus. *Brain Res.* **716**, 72-83.
- Hanisch, U. K., Seto, D. and Quirion, R. (1993) Modulation of hippocampal acetylcholine release: a potent central action of interleukin-2. *J. Neurosci.* **13**, 3368-3374.
- Haour, F. G., Ban, E. M., Milon, G. M., Baran, D. and Fillion, G. M. (1990) Brain interleukin-1 receptors: characterization and modulation after lipopolysaccharide injection. *Prog. Neuroimmunoenocrinol.* **3**, 196-204.
- Harbuz, M. Z., Stephanou, A., Sarlis, N. and Lightman, S. L. (1992) The effects of recombinant interleukin (IL)-1 alpha, IL-1 beta or IL-6 on hypothalamo-pituitary-adrenal axis activation. *J. Endocrinol.* **133**, 349-355.
- Hardin-Pouzet, H., Giraudon, P., Bernard, A., Derrington, E., Belin, M. F. and Didier-Bazet, M. (1996) Cytokines are increased in the rat hippocampus after serotonergic neuron degeneration and upregulate the expression of GDH, an enzyme involved in glutamate detoxification. *J. Neuroimmunol.* **69**, 117-122.
- Hare, A. S., Clarke, G. and Tolchard, S. (1995) Bacterial lipopolysaccharide-induced changes in FOS protein expression in the rat brain: correlation with neuroendocrine and thermoregulatory changes. *J. Neuroendocrinol.* **7**, 791-799.
- Harnett, M. and Rigley, K. (1992) The role of G-proteins versus protein tyrosine kinases in the regulation of lymphocyte activation. *Immunol. Today* **13**, 482-486.
- Hart, R. P., Shadiack, A. M. and Jonakait, G. M. (1991) Substance P gene expression is regulated by interleukin-1 in cultured sympathetic ganglia. *J. Neurosci. Res.* **29**, 282-291.
- Hart, R. P., Liu, C., Shadiack, A. M., McCormack, R. J. and Jonakait, G. M. (1993) An mRNA homologous to interleukin-1 receptor type I is expressed in cultured rat sympathetic ganglia. *J. Neuroimmunol.* **44**, 49-56.
- Hauger, R. L., Irwin, M. R., Lorang, M., Aguilera, G. and Brown, M. R. (1993) High intracerebral levels of CRH result in CRH receptor downregulation in the amygdala and neuroimmune desensitization. *Brain Res.* **616**, 283-292.
- Heimer, L. (1995a) Amygdaloid body and extended amygdala. Ch. 20. In: *The Human Brain and Spinal Cord. Functional Neuroanatomy and Dissection Guide, Second Edition*, pp. 415-421. Ed. L. Heimer. Springer: New York.
- Heimer, L. (1995b) Hippocampal formation. Ch. 21. In: *The Human Brain and Spinal Cord. Functional Neuroanatomy, and Dissection Guide, Second Edition*, pp. 423-431. Ed. L. Heimer. Springer: New York.
- Heimer, L. (1995c) Basal ganglia and related basal forebrain structures. Ch. 16. In: *The Human Brain and Spinal Cord*.

- Functional Neuroanatomy and Dissection Guide, Second Edition*, pp. 339/352. Ed. L. Heimer. Springer: New York.
- Heimer, L. (1995d) Basal ganglia and related basal forebrain structures, Ch. 16. In: *The Human Brain and Spinal Cord. Functional Neuroanatomy and Dissection Guide, Second Edition*, pp. 346-347. Ed. L. Heimer. Springer: New York.
- Heimer, L. (1995e) Brain stem, monoaminergic pathways, and reticular formation. Ch. 10. In: *The Human Brain and Spinal Cord. Functional Neuroanatomy and Dissection Guide, Second Edition*, pp. 227-229. Ed. L. Heimer. Springer: New York.
- Hellstrand, K. and Hermodsson, S. (1990a) Enhancement of human natural killer cell cytotoxicity by serotonin: role of non-T/CD16+ NK cells, monocytes and 5-HT-1A receptors. *Cell. Immunol.* **12**, 199-214.
- Hellstrand, K. and Hermodsson, S. (1990b) Monocyte mediated suppression of human natural killer cell cytotoxicity: regulation by serotonergic 5-HT-1A receptors. *Scand. J. Immunol.* **32**, 183-192.
- Henke, P. G. (1980a) The amygdala and restraint ulcers in rats. *J. Comp. Physiol. Psychol.* **94**, 313-323.
- Henke, P. G. (1980b) Facilitation and inhibition of gastric pathology after lesions in the amygdala of rats. *Physiol. Behav.* **25**, 575-579.
- Henke, P. G. (1985) The amygdala and forced immobilization of rats. *Behav. Brain Res.* **16**, 19-24.
- Herman, J. P., Cullinan, W. E. and Watson, S. J. (1994) Involvement of the bed nucleus of the stria terminalis in tonic regulation of paraventricular hypothalamic CRH and AVP mRNA expression. *J. Neuroendocrinol.* **6**, 433-442.
- Hernandez, M. C., Flores, L. R. and Bayer, B. M. (1993) Immunosuppression by morphine is mediated by central pathways. *J. Pharmacol. Exp. Ther.* **267**, 1336-1341.
- Hikawa, N. and Takenaka, T. (1996) Sensory neurons regulate immunoglobulin secretion of spleen cells: cellular analysis of bidirectional communications between neurons and immune cells. *J. Neuroimmunol.* **70**, 191-198.
- Hirasawa, A., Nakayama, Y., Ishiharada, N., Honda, K., Saito, R., Tsujimoto, G., Takano, Y. and Kamiya, H. (1994) Evidence for the existence of vasopressin V2 receptor mRNA in rat hippocampus. *Biochem. Biophys. Res. Commun.* **205**, 1702-1706.
- Hoffman-Goetz, L. and Pedersen, B. K. (1994) Exercise and the immune system: a model of the stress response?. *Immunol. Today* **15**, 382-387.
- Holmes, M. C., French, K. L. and Seeki, J. R. (1995) Modulation of serotonin and corticosteroid receptor gene expression in the rat hippocampus with circadian rhythm and stress. *Mol. Brain Res.* **28**, 186-192.
- Hsueh, C.-M., Tying, S. K., Hiramoto, R. N. and Ghanta, V. K. (1994) Efferent signal(s) responsible for the conditioned augmentation of natural killer cell activity. *Neuroimmunomodulation* **1**, 74-81.
- Hsueh, C.-M., Chen, S.-F., Ghanta, V. K. and Hiramoto, R. N. (1995) Expression of the conditioned NK cell activity is  $\beta$ -endorphin dependent. *Brain Res.* **678**, 76-82.
- Hu, G.-Y., Hvalby, & Oslash;, Walaas, S. I., Albert, K. A., Skjeflo, P., Andersen, P. and Greengard, P. (1987) Protein kinase C injection into hippocampal pyramidal cells elicits features of long term potentiation. *Nature* **328**, 426-429.
- Hu, Y., Dietrich, H., Herold, M., Heinrich, P. C. and Wick, G. (1993) Disturbed immuno-endocrine communication via the hypothalamo-pituitary-adrenal axis in autoimmune disease. *Int. Arch. Allergy Immunol.* **102**, 232-241.
- Hyman, B. T., Van Hoesen, G. W., Damasio, A. R. and Barnes, C. L. (1984) Alzheimer's disease: cell-specific pathology isolates the hippocampal formation. *Science* **225**, 1168-1170.
- Ikegaya, Y., Saito, H. and Abe, K. (1995) Requirement of basolateral amygdala neuron activity for the induction of long-term potentiation in the dentate gyrus *in vivo*. *Brain Res.* **671**, 351-354.
- Ikegaya, Y., Saito, H. and Abe, K. (1996) Dentate gyrus field potentials evoked by stimulation of the basolateral amygdaloid nucleus in anesthetized rats. *Brain Res.* **718**, 53-60.
- Imperato, A., Puglisi-Allegra, S., Casolini, P. and Angelucci, L. (1991) Changes in brain dopamine and acetylcholine release during and following stress are independent of the pituitary-adrenocortical axis. *Brain Res.* **538**, 111-117.
- Imperato, A., Angelucci, L., Casolini, P., Zocchi, A. and Puglisi-Allegra, S. (1992) Repeated stressful experiences differently affect limbic dopamine release during and following stress. *Brain Res.* **577**, 194-199.
- Irwin, M. (1994) Stress-induced immune suppression: role of brain corticotropin releasing hormone and autonomic nervous system mechanisms. *Adv. Neuroimmunol.* **4**, 29-47.
- Irwin, M. R., Vale, W. and Britton, K. T. (1987) Central corticotropin-releasing factor suppresses natural killer cytotoxicity. *Brain Behav. Immun.* **1**, 81-87.
- Irwin, M., Hauger, R. L., Jones, L., Provencio, M. and Britton, K. T. (1990) Sympathetic nervous system mediates central corticotropin-releasing factor induced suppression of natural killer cytotoxicity. *J. Pharmacol. Exp. Ther.* **255**, 101-107.
- Irwin, M., Brown, M., Patterson, T., Hauger, R., Maseovich, A. and Grant, I. (1991) Neuropeptide Y and natural killer cell activity: findings in depression and Alzheimer caregiver stress. *FASEB J.* **5**, 3100-3107.
- Irwin, M., Hauger, R. and Brown, M. (1992) Central corticotropin releasing hormone activates the sympathetic nervous system and reduces immune function: increased responsiveness of the aged rat. *Endocrinology* **131**, 1047-1053.
- Ishiyama, J., Saito, H. and Abe, K. (1991) Epidermal growth factor and basic fibroblast growth factor promote the generation of long-term potentiation in the dentate gyrus of anesthetized rats. *Neurosci. Res.* **12**, 403-411.
- Jacobs, B. L. and Azmitia, E. C. (1992) Structure and function of the brain serotonin system. *Physiol. Rev.* **72**, 165-229.
- Jacobson, L. and Sapolsky, R. (1991) The role of the hippocampus in feedback regulation of the hypothalamic-pituitary-adrenocortical axis. *Endocr. Rev.* **12**, 118-134.
- Jafarian Tehrani, M., Hu, Y., Marquette, C., Dietrich, H., Haour, F. and Wick, G. (1994) Interleukin-1 receptor deficiency in brains from NZB and (NZB/NZW)F1 autoimmune mice. *J. Neuroimmunol.* **53**, 91-99.
- Johansson, G., Olsson, K., Haggendal, J., Jonsson, L. and Thoren-Tolling, K. (1981) Effect of amygdectomy on stress-induced myocardial necrosis and blood levels of catecholamines in pigs. *Acta Physiol. Scand.* **113**, 553-555.
- Johnson, A. K. and Gross, P. M. (1993) Sensory circumventricular organs and brain homeostatic pathways. *FASEB J.* **7**, 678-686.
- Jonakait, G. M. and Schotland, S. (1990) Conditioned medium from activated splenocytes increases substance P in sympathetic ganglia. *J. Neurosci. Res.* **26**, 24-30.
- Jonakait, G. M., Schotland, S. and Hart, R. P. (1990) Interleukin-1 specifically increases substance P in injured sympathetic ganglia. *Ann. NY Acad. Sci.* **594**, 222-230.
- Kalin, N. H., Takahashi, L. K. and Chen, F.-L. (1994) Restraint stress increases corticotropin-releasing hormone mRNA content in the amygdala and paraventricular nucleus. *Brain Res.* **656**, 182-186.
- Kaneda, T., Kitamura, Y. and Nomura, Y. (1993) Presence of m3 subtype muscarinic acetylcholine receptors and receptor-mediated increases in the cytoplasmic concentration of Ca<sup>2+</sup> in Jurkat, a human leukemic helper T lymphocyte line. *Mol. Pharmacol.* **43**, 356-364.
- Kannan, Y., Bienenstock, J., Ohta, M., Stanisz, A. M. and Stead, R. H. (1996) Nerve growth factor and cytokines mediate lymphoid tissue-induced neurite outgrowth from mouse superior cervical ganglia *in vitro*. *J. Immunol.* **156**, 313-320.
- Katsuki, H., Nakai, Y., Akaji, K. and Satoh, M. (1990) Interleukin-1 beta inhibits long-term potentiation in the CA3 region of mouse hippocampal slices. *Eur. J. Pharmacol.* **181**, 323-326.
- Kelley, K. W., Dantzer, R., Mormede, P., Salmon, H. and Aynaud, J. (1985) Conditioned taste aversion suppresses induction of delayed-type hypersensitivity immune reactions. *Physiol. Behav.* **34**, 193-198.
- Khan, I. and Collins, S. M. (1994) Expression of cytokines in the longitudinal muscle myenteric plexus of the inflamed intestine of rat. *Gastroenterology* **107**, 691-700.
- Khansari, D. N., Murgu, A. J. and Faith, R. E. (1990) Effects of stress on the immune system. *Immunol. Today* **11**, 170-175.
- King, M. G., Husband, A. J. and Kusnecov, A. W. (1987) Behaviourally conditioned immunosuppression using anti-lymphocyte serum: duration of effect and role of corticosteroids. *Med. Sci. Res.* **15**, 407-408.
- Kiss, J. P., Zsilla, G., Mike, A., Zelles, T., Toth, E., Lajtha, A. and Vizi, E. S. (1995) Subtype-specificity of the presynaptic

- $\alpha$ -adrenoceptors modulating hippocampal norepinephrine release in rat. *Brain Res.* **674**, 238-244.
- Kluger, M. J. (1991) Fever: role of pyrogens and cryogens. *Physiol. Rev.* **71**, 93-127.
- Kolesnick, R. and Golde, D. W. (1994) The sphingomyelin pathway in tumor necrosis factor and interleukin-1 signaling. *Cell* **77**, 325-328.
- Komourian, J. and Quik, M. (1996) Characterization of nicotinic receptors in immortalized hippocampal neurons. *Brain Res.* **718**, 37-45.
- Korneva, E. A., Klimenko, V. M. and Shkhinek, E. K. (1985) *Neurohumoral Maintenance of Immune Homeostasis*. University of Chicago Press: Chicago. Translated and edited by S. A. Corson and E. O'L. Corson.
- Kort, W. J. (1994) The effect of chronic stress on the immune response. *Adv. Neuroimmunol.* **4**, 1-11.
- Krueger, J. M. and Majde, J. A. (1995) Cytokines and sleep. *Int. Arch. Allergy Immunol.* **106**, 97-100.
- Krugers, H. J., Koolhaas, J. M., Bohus, B. and Korf, J. (1993) A single social stress-experience alters glutamate receptor-binding in rat hippocampal CA3 area. *Neurosci. Lett.* **154**, 73-77.
- Kushima, Y. and Hatanaka, H. (1992) Interleukin-6 and leukemia inhibitory factor promote the survival of acetylcholinesterase-positive neurons in culture from embryonic rat spinal cord. *Neurosci. Lett.* **143**, 110-114.
- Kushima, Y., Hama, T. and Hatanaka, H. (1992) Interleukin-6 as a neurotrophic factor for promoting the survival of cultured catecholaminergic neurons in a chemically defined medium. *Neurosci. Res.* **13**, 267-280.
- Lacosta, S., Merahi, Z., Zalcman, S. and Anisman, H. (1994) Time-dependent *in vitro* mesolimbic dopamine variations following antigenic challenge. *Brain Res.* **664**, 225-230.
- Lamour, Y., Bassant, M. H., Senut, M. C. and Dutar, P. (1989) Brain aging, Alzheimer's disease and the cholinergic system. In: *Neurotransmission and Cerebrovascular Function*, Vol. 1, pp. 3-28. Eds J. Seylaz and E. T. MacKenzie. Elsevier: Amsterdam.
- Landmann, R. (1992) Beta-adrenergic receptors in human leukocyte subpopulations. *Eur. J. Clin. Invest.* **22**, Suppl. 1, 30-36.
- Lapchak, P., Araujo, D., Quirion, R. and Beaudet, A. (1991) Immunohistochemical localization of interleukin 2-like immunoreactivity and interleukin 2 receptors (Tac antigen-like immunoreactivity) in rat brain. *Neuroscience* **44**, 173-184.
- Lapchak, P. A., Araujo, D. M. and Hefti, F. (1993) Systemic interleukin-1 $\beta$  decreases brain-derived neurotrophic factor messenger RNA expression in the rat hippocampal formation. *Neuroscience* **53**, 297-301.
- Launay, J.-M., Copel, L., Callebert, J., Corvaia, N., Bricaire, F., Laplanche, J.-L., Saal, F. and Peries, J. (1989) Serotonin and human immunodeficiency viruses. *Nouv. Rev. Fr. Hematol.* **31**, 159-161.
- Laurenzi, M. A., Persson, M. A. A., Dalsgaard, C. J. and Haegerstrand, A. (1990) The neuropeptide substance P stimulates production of interleukin-1 in human blood monocytes: activated cells are preferentially influenced by the neuropeptide. *Scand. J. Immunol.* **31**, 529-533.
- Layé, S., Bluthé, R. M., Kent, S., Combe, C., Medina, C., Parnet, P., Kelley, K. W. and Dantzer, R. (1995) Subdiaphragmatic vagotomy blocks the induction of interleukin-1 $\beta$  mRNA in the brain of mice in response to peripherally administered lipopolysaccharide. *Am. J. Physiol. (Regul. Integr. Comp. Physiol.)* **268**, R1327-R1331.
- Le Moal, M. and Simon, H. (1991) Mesocorticolimbic dopaminergic network: functional and regulatory roles. *Physiol. Rev.* **71**, **1**, 155-234.
- Léránth, C. and Frotscher, M. (1983) Commissural afferents to the rat hippocampus terminate on vasoactive intestinal polypeptide-like immunoreactive non-pyramidal neurons. An EM immunocytochemical degeneration study. *Brain Res.* **276**, 357-361.
- Levine, J. D., Clark, R., Devor, M., Helms, C., Moskowitz, M. A. and Basbaum, A. I. (1984) Intraneuronal substance P contributes to the severity of experimental arthritis. *Science* **226**, 547-552.
- Levine, J. D., Dardick, S. J., Roizen, M. F., Helms, C. and Basbaum, A. I. (1986a) Contribution of sensory afferents and sympathetic efferents to joint injury in experimental arthritis. *J. Neurosci.* **6**, 3423-3429.
- Levine, J. D., Fye, K., Heller, P., Basbaum, A. I. and Whiting-O'Keefe, Q. (1986b) Clinical response to regional intravenous guanethidine in patients with rheumatoid arthritis. *J. Rheumatol.* **13**, 1040-1043.
- Levine, J. D., Coderre, T. J., Helms, C. and Basbaum, A. I. (1988)  $\beta$ -Adrenergic mechanisms in experimental arthritis. *Proc. Natl. Acad. Sci. U.S.A.* **85**, 4553-4556.
- Liang, K. C. and Lee, E. H. Y. (1988) Intra-amygdala injections of corticotropin-releasing factor facilitate inhibitory avoidance learning and reduce exploratory behavior in rats. *Psychopharmacology* **96**, 232-236.
- Liang, K. C., Hon, W. and Davis, M. (1994) Pre- and post-training infusion of *N*-methyl-D-aspartate receptor antagonists into the amygdala impair memory in an inhibitory avoidance task. *Behav. Neurosci.* **108**, 241-253.
- Licinio, J., Wong, M. and Gold, P. W. (1991) Localization of interleukin-1 receptor antagonist mRNA in rat brain. *Endocrinology* **129**, 562-564.
- Licinio, J., Wong, M. L. and Gold, P. W. (1992) Neutrophil-activating peptide-1/interleukin-8 mRNA is localized in rat hypothalamus and hippocampus. *Neuroreport* **3**, 753-756.
- Lignon, M. F., Bernad, N. and Martinez, J. (1991) Pharmacological characterization of type B cholecystokinin binding sites on human Jurkat T lymphocyte cell line. *Mol. Pharmacol.* **39**, 615-620.
- Linthorst, A. C. E., Flachskamm, C., Holsboer, F. and Reul, J. M. H. M. (1994) Local administration of recombinant human interleukin-1 $\beta$  in the rat hippocampus increases serotonergic neurotransmission, hypothalamic-pituitary-adrenocortical axis activity, and body temperature. *Endocrinology* **135**, 520-532.
- Livnat, S., Madden, K. S., Felten, D. L. and Felten, S. Y. (1987) Regulation of the immune system by sympathetic neural mechanisms. *Prog. Neuropsychopharmacol. Biol. Psychiat.* **11**, 145-152.
- Lorton, D., Bellinger, D., Ducloux, M., Felten, S. Y. and Felten, D. L. (1996) Application of 6-hydroxydopamine into the fatpads surrounding the draining lymph nodes exacerbates adjuvant-induced arthritis. *J. Neuroimmunol.* **64**, 103-113.
- Lovenberg, T. W., Liaw, C. W., Grigoriadis, D. E., Clevenger, W., Chalmers, D. T., De Souza, E. B. and Oltersdorf, T. (1995) Cloning and characterization of a functionally distinct corticotropin-releasing factor receptor subtype from rat brain. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 836-840.
- Ludlam, W. H., Chandross, K. J. and Kessler, J. A. (1995) LIF- and IL-1 $\beta$ -mediated increases in substance P receptor mRNA in axotomized, explanted or dissociated sympathetic ganglia. *Brain Res.* **685**, 12-20.
- Luecken, L. J. and Lysle, D. T. (1992) Evidence for the involvement of  $\beta$ -adrenergic receptors in conditioned immunomodulation. *J. Neuroimmunol.* **38**, 209-220.
- Lysle, D. T., Cunnick, J. E., Fowler, H. and Rabin, B. (1988) Pavlovian conditioning of shock-induced suppression of lymphocyte reactivity: acquisition, extinction, and preexposure effects. *Life Sci.* **42**, 2185-2194.
- Lysle, D. T., Cunnick, J. E. and Maslonek, K. A. (1991) Pharmacological manipulation of immune alterations induced by an aversive conditioned stimulus: evidence for a  $\beta$ -adrenergic receptor-mediated Pavlovian conditioning process. *Behav. Neurosci.* **105**, 443-449.
- Lyte, M., Ernst, S., Driemeyer, J. and Baissa, B. (1991) Stain-specific enhancement of splenic T cell mitogenesis and macrophage phagocytosis following peripheral axotomy. *J. Neuroimmunol.* **31**, 1-8.
- Madden, K. S., Felten, S. Y., Felten, D. L., Hardy, C. A. and Livnat, S. (1994a) Sympathetic nervous system modulation of the immune system. II. Induction of lymphocyte proliferation and migration *in vivo* by chemical sympathectomy. *J. Neuroimmunol.* **49**, 67-75.
- Madden, K. S., Moynihan, J. A., Brenner, G. J., Felten, S. Y., Felten, D. L. and Livnat, S. (1994b) Sympathetic nervous system modulation of the immune system. III. Alterations in T and B cell proliferation and differentiation *in vitro* following chemical sympathectomy. *J. Neuroimmunol.* **49**, 77-87.
- Maeda, H. and Mogenson, C. J. (1981) Electrophysiological responses of neurons of the ventral tegmental area to electrical stimulation of amygdala and lateral septum. *Neuroscience* **6**, 367-376.
- Maes, M., Bosmans, E., Meltzer, H. Y., Scharpe, S. and Suy, E. (1993) Interleukin-1 beta: a putative mediator of HPA axis hyperactivity in major depression?. *Am. J. Psychiat.* **150**, 1189-1193.

- Main, C., Blennerhassett, P. and Collins, S. M. (1993) Human recombinant interleukin  $\beta$  suppresses acetylcholine release from rat myenteric plexus. *Gastroenterology* **104**, 1648–1654.
- Makino, S., Schulkin, J., Smith, M. A., Pacák, K., Palkovits, M. and Gold, P. W. (1995) Regulation of corticotropin-releasing hormone receptor messenger ribonucleic acid in the rat brain and pituitary by glucocorticoids and stress. *Endocrinology* **136**, 4517–4525.
- Malinow, R., Madison, D. V. and Tsien, R. W. (1988) Persistent protein kinase activity underlying long-term potentiation. *Nature* **335**, 820–824.
- Mameya, S.-I., Sawa, T. and Taniyama, K. (1995) Arachidonic acid cascade and stimulation of acetylcholine release by human recombinant interleukin- $\beta$  in guinea pig ileum. *J. Pharmacol. Exp. Ther.* **275**, 319–324.
- Maness, L. M., Banks, W. A., Zadina, J. E. and Kastin, A. J. (1995) Selective transport of blood-borne interleukin-1 $\alpha$  into the posterior division of the septum of the mouse brain. *Brain Res.* **700**, 83–88.
- Mapp, P. J., Terenghi, G., Walsh, D. A., Chen, S. T., Cruwys, S. C., Garrett, N., Kidd, B. L., Polak, J. M. and Blake, D. R. (1993) Monoarthritis in the rat knee induces bilateral and time-dependent peptide immunoreactivity in the spinal cord. *Neuroscience* **57**, 1091–1096.
- Martin, K. F., Hannon, S., Phillips, J. and Heal, D. J. (1992) Opposing roles for 5-HT $_B$  and 5-HT $_T$  receptors in the control of 5-HT release in rat hippocampus *in vivo*. *Br. J. Pharmacol.* **106**, 139–142.
- Martin, L. J., Powers, R. E., Dellovade, T. L. and Price, D. L. (1991) The bed nucleus-amygdala continuum in human and monkey. *J. Comp. Neurol.* **309**, 445–485.
- März, P., Gadiant, R. A. and Otten, U. (1996) Expression of interleukin-6 receptor (IL-6R) and gp130 mRNA in PC12 cells and sympathetic neurons: modulation by tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ). *Brain Res.* **706**, 71–79.
- Masek, K., Petrovicky, P. and Seifert, J. (1992) An introduction to the possible role of central nervous system structures in neuroendocrine-immune systems interaction. *Int. J. Immunopharmacol.* **14**, 317–322.
- Mason, D., MacPhee, I. and Antoni, F. (1990) The role of the neuroendocrine system in determining genetic susceptibility to experimental allergic encephalomyelitis in the rat. *Immunology* **70**, 1–5.
- Matsumoto, M., Yoshioka, M., Togashi, H., Tochiwara, M., Ikeda, T. and Saito, H. (1995) Modulation of norepinephrine release by serotonergic receptors in the rat hippocampus as measured by *in vivo* microdialysis. *J. Pharmacol. Exp. Ther.* **272**, 1044–1051.
- Matta, S. G., Singh, J., Newton, R. and Sharp, B. M. (1990) The adrenocorticotropin response to interleukin- $\beta$  instilled into the rat median eminence depends on the local release of catecholamines. *Endocrinology* **127**, 2175–2181.
- Mayer, M. L. and Vyklicky, L. Jr. (1989) Concanavalin A selectively reduces desensitization of mammalian neuronal quisqualate receptors. *Proc. Natl. Acad. Sci. U.S.A.* **86**, 1411–1415.
- McEwen, B. S. and Sapolsky, R. M. (1995) Stress and cognitive function. *Curr. Opin. Neurobiol.* **5**, 205–216.
- McEwen, B. S. and Woolley, C. S. (1994) Estradiol and progesterone regulate neuronal structure and synaptic connectivity in adult as well as developing brain. *Exp. Gerontol.* **29**, 431–436.
- McEwen, B. S., DeKloet, E. R. and Rostene, W. (1986) Adrenal steroid receptors and actions in the nervous system. *Physiol. Rev.* **66**, 1121–1137.
- Mello, L. E. A. M., Tan, A. M. and Finch, D. M. (1992) GABAergic synaptic transmission in projections from the basal forebrain and hippocampal formation to the amygdala: an *in vivo* iontophoretic study. *Brain Res.* **587**, 41–48.
- Michelson, D., Stone, L., Galliven, E., Magiakou, M. A., Chrousos, G. P., Sternberg, E. M. and Gold, P. W. (1994) Multiple sclerosis is associated with alterations in hypothalamic-pituitary-adrenal axis function. *J. Clin. Endocrinol. Metab.* **79**, 848–853.
- Micic, M., Lepasovic, G. and Ugresic, N. (1994) Relationship between monoaminergic and cholinergic innervation of the rat thymus during aging. *J. Neuroimmunol.* **49**, 205–212.
- Millan, M. A., Jacobowitz, D. M., Hauger, R. L., Catt, K. J. and Aguilera, G. (1986) Distribution of corticotropin-releasing factor receptors in primate brain. *Proc. Natl. Acad. Sci. U.S.A.* **83**, 1921–1925.
- Miller, A. H., Sastry, G., Speranza, A. J. Jr, Lawlor, B. A., Mohs, R. C., Ryan, T. M., Gabriel, S. M., Serby, M., Schmeidler, J. and Davis, K. L. (1994) Lack of association between cortisol hypersecretion and non-suppression on the DST in patients with Alzheimer's disease. *Am. J. Psychiat.* **151**, 267–270.
- Minami, M., Kuraishi, Y., Yamaguchi, T., Nakai, S., Hirai, Y. and Satoh, M. (1991) Immobilization stress induces interleukin-1 beta mRNA in the rat hypothalamus. *Neurosci. Lett.* **123**, 254–256.
- Mishkin, M. (1993) Cerebral memory circuits. In: *Exploring Brain Functions: Models in Neuroscience*, pp. 113–125. Eds T. A. Poggio and D. A. Glaser. Wiley: New York.
- Morale, M. C., Batticane, N., Gallo, F., Barden, N. and Marchetti, B. (1995) Disruption of hypothalamic-pituitary-adrenocortical system in transgenic mice expressing type II glucocorticoid receptor antisense ribonucleic acid permanently impairs T cell function: effects on T cell trafficking and T cell responsiveness during postnatal development. *Endocrinology* **136**, 3949–3960.
- Morales, M., Battenberg, E., de Lecea, L., Sanna, P. P. and Bloom, F. E. (1996) Cellular and subcellular immunolocalization of the type 3 serotonin receptor in the rat central nervous system. *Mol. Brain Res.* **36**, 251–260.
- Morrow, L. E., McClellan, J. L., Conn, C. A. and Kluger, M. J. (1993a) Glucocorticoids alter fever and IL-6 responses to psychological stress and to lipopolysaccharide. *Am. J. Physiol. (Regul. Integr. Comp. Physiol.)* **264**, R1010–R1016.
- Morrow, N. S., Grijalva, C. V., Geiselman, P. J. and Novin, D. (1993b) Effects of amygdaloid lesions on gastric erosion formation during exposure to activity-stress. *Physiol. Behav.* **53**, 1043–1048.
- Muir, J. L., Everitt, B. J. and Robbins, T. W. (1994) AMPA-induced excitotoxic lesions of the basal forebrain: a significant role for the cortical cholinergic system in attentional function. *J. Neurosci.* **14**, 2313–2326.
- Mullan, M. (1993) The molecular pathology of Alzheimer's disease. *Int. Rev. Psychiat.* **5**, 351–362.
- Nakamura-Craig, M. and Smith, T. W. (1989) Substance P and peripheral inflammatory hyperalgesia. *Pain* **38**, 91–98.
- Nakatsuru, K., Ohgo, S., Oki, Y. and Matsukura, S. (1991) Interleukin-1 (IL-1) stimulates arginine vasopressin (AVP) release from superfused rat hypothalamo-neurohypophyseal complexes independently of cholinergic mechanism. *Brain Res.* **554**, 38–45.
- Nance, D. M., Hopkins, D. A. and Bieger, D. (1987) Re-investigation of the innervation of the thymus gland in mice and rats. *Brain Behav. Immun.* **1**, 134–147.
- Nemeroff, C. B., Widerlov, E. and Bissette, G. (1984) Elevated concentrations of CSF corticotropin-releasing factor-like immunoreactivity in depressed patients. *Science* **226**, 1342–1344.
- Nemeroff, C. B., Owens, M. J., Bissette, G. and Stanley, M. (1988) Reduced corticotropin releasing factor binding sites in the frontal cortex of suicide victims. *Arch. Gen. Psychiat.* **45**, 577–582.
- Nemeroff, C. B., Krishnan, K. R. R., Reed, D., Leder, R., Beam, C. and Dumnick, N. R. (1992) Adrenal gland enlargement in major depression. *Arch. Gen. Psychiat.* **49**, 384–387.
- Nemni, R., Iannaccone, S., Quattrini, A., Smirne, S., Sessa, M., Lodi, M., Erminio, C. and Canal, N. (1992) Effect of chronic treatment with recombinant interleukin-2 on the central nervous system of adult and old mice. *Brain Res.* **591**, 248–252.
- Neveu, P. J. (1992) Asymmetrical brain modulation of the immune response. *Brain Res. Rev.* **17**, 101–107.
- Nijijima, A. (1992) The afferent discharges from sensors for interleukin-1 $\beta$  in the hepato-portal system in the anaesthetized rat. *J. Physiol.* **446**, 236P.
- Nijijima, A. (1995) An electrophysiological study on the vagal innervation of the thymus in the rat. *Brain Res. Bull.* **38**, 319–323.
- Nijijima, A., Hori, T., Aou, S. and Oomura, Y. (1991) The effects of interleukin  $\beta$  on the activity of adrenal, splenic and renal sympathetic nerves in the rat. *J. Auton. Nerv. Syst.* **36**, 183–192.
- Nijijima, A., Hori, T., Katafuchi, T. and Ichijo, T. (1995) The effect of interleukin- $\beta$  on the efferent activity of the vagus nerve to the thymus. *J. Auton. Nerv. Syst.* **54**, 137–144.
- Nistico, G., Caroleo, M. C., Arbitrio, M. and Pulvirenti, L. (1994) Dopamine D1 receptors in the amygdala enhance the immune response in the rat. *Ann. NY Acad. Sci.* **741**, 316–323.
- Obata-Tsuto, H.-L. (1987) Light and electron microscopic study of somatostatin-like immunoreactive neurons in rat hippocampus. *Brain Res. Bull.* **18**, 613–620.
- O'Brien, J. T., Ames, D. and Schweitzer, I. (1993) HPA axis function in depression and dementia: a review. *Int. J. Geriatr. Psychiat.* **8**, 887–898.

- Ono, T., Nishijo, H. and Uwano, T. (1995) Amygdala role in conditioned associative learning. *Prog. Neurobiol.* **46**, 401-422.
- Orchinik, M., Welland, N. G. and McEwen, B. S. (1995) Chronic exposure to stress levels of corticosterone alters GABA<sub>A</sub> receptor subunit mRNA levels in rat hippocampus. *Mol. Brain Res.* **34**, 29-37.
- Osborne, P. G. (1994) A GABAergic mechanism in the medial septum influences cortical arousal and locomotor activity but not a previously learned spatial discrimination task. *Neurosci. Lett.* **173**, 63-66.
- OVADIA, H. and ABRAMSKY, O. (1987) Dopamine receptors on isolated membranes of rat lymphocytes. In: *Neuroimmunomodulation*, pp. 70-74. Eds J. R. Perez-Polo, K. Bulloch, R. H. Angeletti, G. A. Hashim and J. de Vellis. Alan R. Liss, Inc.: New York.
- Pacak, K., McCarty, R., Palkovits, M., Kopin, I. J. and Goldstein, D. S. (1995) Effects of immobilization on *in vivo* release of norepinephrine in the bed nucleus of the stria terminalis in conscious rats. *Brain Res.* **688**, 242-246.
- Palkovits, M., Somogyvári-Vigh, A. and Arimura, A. (1995) Concentrations of pituitary adenylate cyclase activating polypeptide (PACAP) in human brain nuclei. *Brain Res.* **699**, 116-120.
- Pan, Q. and Long, J. (1993) Lesions of the hippocampus enhance or depress humoral immunity in rats. *Neuroreport* **4**, 864-866.
- Parent, A. and Hazrati, L.-N. (1995) Functional anatomy of the basal ganglia. I. The cortico-basal ganglia-thalamo-cortical loop. *Brain Res. Rev.* **20**, 91-127.
- Patel, N. A., Romero, A. A., Zadina, J. E. and Chang, S. L. (1996) Chronic exposure to morphine attenuates expression of interleukin-1 $\beta$  in the rat hippocampus. *Brain Res.* **712**, 340-344.
- Pavlov, I. P. (1928) *Lectures on Conditioned Reflexes*. Liveright: New York.
- Peeke, H. V. S., Ellman, G., Dark, K., Salfi, M. and Reus, V. J. (1987) Cortisol and behaviorally conditioned histamine release. *Ann. NY Acad. Sci.* **496**, 583-587.
- Perez, L. and Lysic, D. T. (1995) Corticotropin-releasing hormone is involved in conditioned stimulus-induced reduction of natural killer cell activity but not in conditioned alterations in cytokine production or proliferative responses. *J. Neuroimmunol.* **63**, 1-8.
- Petitot, J. M. and Huang, Z. (1994) Molecular cloning of a partial cDNA of the interleukin-2 receptor- $\beta$  in normal mouse brain: *in situ* localization in the hippocampus and expression by neuroblastoma cells. *Brain Res.* **650**, 140-145.
- Pezzone, M. A., Lee, W.-S., Hoffman, G. E. and Rabin, B. S. (1992) Induction of c-Fos immunoreactivity in the rat forebrain by conditioned and unconditioned aversive stimuli. *Brain Res.* **597**, 41-50.
- Phillips, H. S., Hains, J. M., Laramee, G. R., Rosenthal, A. and Winslow, J. W. (1990) Widespread expression of BDNF but not NT-3 by target areas of basal forebrain cholinergic neurons. *Science* **250**, 290-294.
- Plata-Salamán, C. R. and Ffrench-Mullen, J. M. H. (1992) Interleukin-1 $\beta$  depresses calcium currents in CA1 hippocampal neurons at pathophysiological concentrations. *Brain Res. Bull.* **29**, 221-223.
- Plata-Salamán, C. R. and Ffrench-Mullen, J. M. (1993) Interleukin-2 modulates calcium currents in dissociated hippocampal CA1 neurons. *Neuroreport* **4**, 579-581.
- Pousset, F., Fournier, J., Legoux, P., Keane, P., Shire, D. and Soubrie, P. (1996) Effect of serotonin on cytokine mRNA expression in rat hippocampal astrocytes. *Mol. Brain Res.* **38**, 54-62.
- Pow, D. V. (1992) NADPH-diaphorase (nitric oxide synthase) staining in the rat supraoptic nucleus is activity dependent: possible functional implications. *J. Neuroendocrinol.* **4**, 377-380.
- Puglisi-Allegra, S., Imperato, A., Angelucci, L. and Cabib, S. (1991) Acute stress induces time-dependent responses in dopamine mesolimbic system. *Brain Res.* **554**, 217-222.
- Raber, J. and Bloom, F. E. (1994) IL-2 induces vasopressin release from the hypothalamus and the amygdala: role of nitric oxide-mediated signaling. *J. Neurosci.* **14**, 6187-6195.
- Raber, J., Koob, G. F. and Bloom, F. E. (1995) Interleukin-2 (IL-2) induces corticotropin-releasing-factor (CRF) release from the amygdala and involves a nitric oxide-mediated signaling: comparison with the hypothalamic response. *J. Pharmacol. Exp. Ther.* **272**, 815-824.
- Rada, P., Mark, G. P., Vitek, M. P., Mangano, R. M., Blume, A. J., Beer, B. and Hoebel, B. G. (1991) Interleukin-1 $\beta$  decreases acetylcholine measured by microdialysis in the hippocampus of freely moving rats. *Brain Res.* **550**, 287-290.
- Rainnie, D. G., Asprodini, E. K. and Shinnick-Gallagher, P. (1991a) Excitatory transmission in the basolateral amygdala. *J. Neurophysiol.* **66**, 986-998.
- Rainnie, D. G., Asprodini, E. K. and Shinnick-Gallagher, P. (1991b) Inhibitory transmission in the basolateral amygdala. *J. Neurophysiol.* **66**, 999-1009.
- Ray, A., Henke, P. G., Gulati, K. and Sen, P. (1993) The amygdaloid complex, corticotropin releasing factor and stress-induced gastric ulcerogenesis in rats. *Brain Res.* **624**, 286-290.
- Renoux, G., Biziere, K., Renoux, M., Bardos, P. and Degenne, D. (1987) Consequences of bilateral brain neocortical ablation on Imithiol-induced immunostimulation in mice. *Ann. NY Acad. Sci.* **496**, 346-353.
- Reul, J. M., Stec, I., Wieggers, G. J., Labeur, M. S., Linthorst, A. C., Arzt, E. and Holtsboer, F. (1994) Prenatal immune challenge alters the hypothalamic-pituitary-adrenocortical axis in adult rats. *J. Clin. Invest.* **93**, 2600-2607.
- Reymann, K. G., Frey, U., Jork, R. and Matthies, H. (1988) Polymyxin B, an inhibitor of protein kinase C, prevents the maintenance of synaptic long-term potentiation in hippocampal CA<sub>1</sub> neurons. *Brain Res.* **440**, 305-314.
- Rieck, R. W., Nabors, C. C., Updyke, B. V. and Kratz, K. E. (1995) Organization of the basal forebrain in the cat: localization of L-enkephalin, substance P, and choline acetyltransferase immunoreactivity. *Brain Res.* **672**, 237-250.
- Riekkinen, P., Buzsáki, G., Riekkinen, P. Jr, Soininen, H. and Partanen, J. (1991) The cholinergic system and EEG slow waves. *Electroencephalogr. Clin. Neurophysiol.* **78**, 89-96.
- Rinner, I. and Schauenstein, K. (1991) The parasympathetic nervous system takes part in the immuno-neuroendocrine dialogue. *J. Neuroimmunol.* **34**, 165-172.
- Rinner, I. and Schauenstein, K. (1993) Detection of choline acetyltransferase activity in lymphocytes. *J. Neurosci. Res.* **35**, 188-191.
- Rinner, I., Schauenstein, K., Mangge, H., Porta, S. and Kvetnansky, R. (1992) Opposite effects of mild and severe stress on *in vitro* activation of rat peripheral blood lymphocytes. *Brain Behav. Immun.* **6**, 130-140.
- Rinner, I., Kukulansky, T., Felsner, P., Skreiner, E., Globerson, A., Kasai, M., Hirokawa, K., Korsatko, W. and Schauenstein, K. (1994) Cholinergic stimulation modulates apoptosis and differentiation of murine thymocytes via a nicotinic effect on thymic epithelium. *Biochem. Biophys. Res. Com.* **203**, 1057-1062.
- Rivest, S., Torres, G. and Rivier, C. (1992) Differential effects of central and peripheral injection of interleukin-1 $\beta$  on brain *c-fos* expression and neuroendocrine functions. *Brain Res.* **587**, 13-23.
- Rivier, C. and Vale, W. (1989) In the rat, interleukin-1 $\alpha$  acts at the level of the brain and the gonads to interfere with gonadotropin and sex steroid secretion. *Endocrinology* **124**, 2105-2109.
- Roberts, G. W., Woodhams, P. L., Polak, J. M. and Crow, T. J. (1984) Distribution of neuropeptides in the limbic system of the rat: the hippocampus. *Neuroscience* **11**, 35-77.
- Romano, T. A., Felten, S. Y., Olschowka, J. A. and Felten, D. L. (1994) Noradrenergic and peptidergic innervation of lymphoid organs in the beluga, *Delphinapterus leucas*: an anatomical link between the nervous and immune systems. *J. Morphol.* **221**, 243-259.
- Roth, R. H., Tam, S.-Y., Ida, Y., Yang, J.-X. and Deutch, A. Y. (1988) Stress and mesocorticolimbic dopamine system. *Ann. NY Acad. Sci.* **537**, 138-147.
- Rothman, S. M. and Olney, J. W. (1987) Excitotoxicity and the NMDA receptor. *Trends Neurosci.* **10**, 299-302.
- Roy, A., Pickar, D., Linnoila, M. and Potter, W. Z. (1985) Plasma norepinephrine level in affective disorders. *Arch. Gen. Psychiat.* **42**, 1181-1185.
- Rudorfer, M. V., Ross, R. J., Linnoila, M., Sherer, M. A. and Potter, W. Z. (1985) Exaggerated orthostatic responsiveness of plasma norepinephrine in depression. *Arch. Gen. Psychiat.* **42**, 1186-1192.
- Rühl, A., Hurst, S. and Collins, S. M. (1994) Synergism between interleukins 1 $\beta$  and 6 on noradrenergic nerves in rat myenteric plexus. *Gastroenterology* **107**, 993-1001.
- Salamone, J. D. (1994) The involvement of nucleus accumbens dopamine in appetitive and aversive motivation. *Behav. Brain Res.* **61**, 117-133.
- Sanders, V. M. (1995) The role of adrenoceptor-mediated signals in

- the modulation of lymphocyte function. *Adv. Neuroimmunol.* **5**, 283–298.
- Sapolsky, R. M. (1986) Glucocorticoid toxicity in the hippocampus: temporal aspects of synergy with kainic acid. *Neuroendocrinology* **43**, 440–444.
- Sapolsky, R. M. and Pulsinelli, W. A. (1985) Glucocorticoids potentiate ischemic injury to neurons: therapeutic implications. *Science* **229**, 1397–1400.
- Sapolsky, R. M., McEwen, B. S. and Rainbow, T. C. (1983) Quantitative autoradiography of [<sup>3</sup>H]corticosterone in rat brain. *Brain Res.* **271**, 331–334.
- Sapolsky, R. M., Krey, L. C. and McEwen, B. S. (1985) Prolonged glucocorticoid exposure reduces hippocampal neuron number: implications for aging. *J. Neurosci.* **5**, 1222–1227.
- Sapolsky, R., Rivier, C., Yamamoto, G., Plotsky, P. and Vale, W. (1987) Interleukin-1 stimulates the secretion of hypothalamic corticotropin-releasing factor. *Science* **238**, 522–524.
- Sapolsky, R. M., Packan, D. R. and Vale, W. W. (1988) Glucocorticoid toxicity in the hippocampus: *in vitro* demonstration. *Brain Res.* **453**, 367–371.
- Sapolsky, R. M., Uno, H., Rebert, C. S. and Finch, C. E. (1990) Hippocampal damage associated with prolonged glucocorticoid exposure in primates. *J. Neurosci.* **10**, 2897–2902.
- Sarder, M., Abe, K., Saito, H. and Nishiyama, N. (1996) Comparative effect of IL-2 and IL-6 on morphology of cultured hippocampal neurons from fetal rat brain. *Brain Res.* **715**, 9–16.
- Savino, W. and Dardenne, M. (1995) Immune neuroendocrine interactions. *Immunol. Today* **16**, 318–322.
- Schacter, G. B., Yang, C. R., Innis, N. K. and Mogenson, G. J. (1989) The role of the hippocampal-nucleus accumbens pathway in radial-arm maze performance. *Brain Res.* **494**, 339–349.
- Schäfer, M., Carter, L. and Stein, C. (1994) Interleukin-1 $\beta$  and corticotropin-releasing factor inhibit pain by releasing opioids from immune cells in inflamed tissue. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 4219–4223.
- Schaible, H. G. and Schmidt, R. F. (1988) Time course of mechanosensitivity changes in articular afferents during a developing arthritis. *J. Neurophysiol.* **60**, 2180–2195.
- Schauenstein, K., Faessler, R., Dietrich, H., Schwarz, S., Kroemer, G. and Wick, G. (1987) Disturbed immune endocrine communication in autoimmune diseases. Lack of corticosterone response to immune signals in obese strain chickens with spontaneous autoimmune thyroiditis. *J. Immunol.* **139**, 1830–1833.
- Schauenstein, K., Rinner, I., Felsner, P., Hofer, D., Mangge, H., Kreiner, E., Liebmann, P. and Globerson, A. (1994) The role of the adrenergic/cholinergic balance in the immune-neuroendocrine circuit. In: *Advances in Psychoneuroimmunology*, pp. 349–356. Eds I. Berczi and J. Szélenyi. Plenum Press: New York.
- Schmued, L. C. (1994) Diagonal ventral forebrain continuum has overlapping telencephalic inputs and brainstem outputs which may represent loci for limbic/autonomic integration. *Brain Res.* **667**, 175–191.
- Schöbitz, B., De Kloet, E. R. and Holsboer, F. (1994a) Gene expression and function of interleukin 1, interleukin 6 and tumor necrosis factor in the brain. *Prog. Neurobiol.* **44**, 397–432.
- Schöbitz, B., Sutanto, W., Carey, M. P., Holsboer, F. and De Kloet, E. R. (1994b) Endotoxin and interleukin 1 decrease the affinity of hippocampal mineralocorticoid (type I) receptor in parallel to activation of the hypothalamic-pituitary-adrenal axis. *Neuroendocrinology* **60**, 124–133.
- Schulte, H. M., Bamberger, C. M., Elsen, H., Herrmann, G., Bamberger, A. M. and Barth, J. (1994) Systemic interleukin-1 $\alpha$  and interleukin-2 secretion in response to acute stress and to corticotropin-releasing hormone in humans. *Eur. J. Clin. Invest.* **24**, 773–777.
- Schuman, E. M. and Madison, D. V. (1991) A requirement for the intercellular messenger nitric oxide in long-term potentiation. *Science* **254**, 1503–1506.
- Sehic, E. and Blatteis, C. M. (1996) Blockade of lipopolysaccharide-induced fever by subdiaphragmatic vagotomy in guinea pigs. *Brain Res.* **726**, 160–166.
- Selami, S. and de Beaurepaire, R. (1995) Hypothalamic and thalamic sites of action of interleukin-1 $\beta$  on food intake, body temperature and pain sensitivity in the rat. *Brain Res.* **694**, 69–77.
- Shaikh, M. B., Steinberg, A. and Siegel, A. (1993) Evidence that substance P is utilized in medial amygdaloid facilitation of defensive rage behavior in the cat. *Brain Res.* **625**, 283–294.
- Shanks, N., Francis, D., Zalzman, S., Meaney, M. J. and Anisman, H. (1994) Alterations in central catecholamines associated with immune responding in adult and aged mice. *Brain Res.* **666**, 77–87.
- Singh, U. and Fatani, J. (1988) Thymic lymphopoiesis and cholinergic innervation. *Thymus* **11**, 3–13.
- Sokolov, M. V. and Kleschevnikov, A. M. (1995) Atropine suppresses associative LTP in the CA1 region of rat hippocampal slices. *Brain Res.* **672**, 281–284.
- Solleveld, H. A., Haseman, J. K. and McConnell, E. E. (1984) Natural history of body weight gain, survival, and neoplasia in the F344 rat. *J. Natl. Cancer Inst.* **72**, 929–940.
- Solvason, H. B., Ghanta, V. K. and Hiramoto, R. N. (1993) The identity of the unconditioned stimulus to the central nervous system is interferon- $\beta$ . *J. Neuroimmunol.* **45**, 75–82.
- Song, C., Earley, B. and Leonard, B. E. (1996) The effects of central administration of neuropeptide Y on behavior, neurotransmitter, and immune functions in the olfactory bulbectomized rat model of depression. *Brain, Behav. Immun.* **10**, 1–16.
- Spangelo, B. L., Isakson, P. C. and MacLeod, R. M. (1990) Production of interleukin-6 by anterior pituitary cells is stimulated by increased intracellular adenosine 3',5'-monophosphate and vasoactive intestinal peptide. *Endocrinology* **127**, 403–409.
- Spranger, M., Lindholm, D., Bandtlow, C., Heumann, R., Grahn, H., Naher-Noe, M. and Thoenen, H. (1990) Regulation of nerve growth factor (NGF) synthesis in the rat central nervous system: comparison between the effects of interleukin-1 and various growth factors in astrocyte cultures and *in vivo*. *Eur. J. Neurosci.* **2**, 69–76.
- Squire, L. R. and Zola-Morgan, S. (1989) The medial temporal lobe memory system. *Science* **253**, 1380–1386.
- Stanisz, A. M., Befus, D. and Bienenstock, J. (1986) Differential effects of VIP, substance P, and somatostatin on immunoglobulin synthesis and proliferations by lymphocytes from Peyer's patches, mesenteric lymph nodes, and spleen. *J. Immunol.* **136**, 152–156.
- Stead, R. H., Tomioka, M., Quinonez, G., Simon, G. T., Felten, S. Y. and Bienenstock, J. (1987) Intestinal mucosal mast cells in normal and nematode-infected rat intestines are in intimate contact with peptidergic nerves. *Proc. Natl. Acad. Sci. U.S.A.* **84**, 2975–2979.
- Stein, C., Hassan, A. H. S., Przewlocki, R., Gramsch, C., Peter, K. and Herz, A. (1990) Opioids from immunocytes interact with receptors on sensory nerves to inhibit nociception in inflammation. *Proc. Natl. Acad. Sci. U.S.A.* **87**, 5935–5939.
- Stenzel-Poore, M., Vale, W. W. and Rivier, C. (1993) Relationship between antigen-induced immune stimulation and activation of the hypothalamic-pituitary-adrenal axis in the rat. *Endocrinology* **132**, 1313–1318.
- Sternberg, E. M., Hill, J. M., Chrousos, G. P., Kamilaris, T., Listwak, S. J., Gold, P. W. and Wilder, R. L. (1989a) Inflammatory mediator-induced hypothalamic-pituitary-adrenal axis activation is defective in streptococcal cell wall arthritis-susceptible Lewis rats. *Proc. Natl. Acad. Sci. U.S.A.* **86**, 2374–2378.
- Sternberg, E. M., Young, W. S., Bernadini, R., Calogero, A. E., Chrousos, G. P., Gold, P. W. and Wilder, R. L. (1989b) A central nervous system defect in biosynthesis of corticotropin releasing hormone is associated with susceptibility to streptococcal cell wall-induced arthritis in Lewis rats. *Proc. natn. Acad. Sci. U.S.A.* **86**, 4771–4775.
- Stitt, J. T. (1990) Passage of immunomodulators across the blood-brain barrier. *Yale J. Biol. Med.* **63**, 121–131.
- Straub, R. H., Lang, B., Falk, W., Schölmerich, J. and Singer, E. A. (1995) *In vitro* superfusion method for the investigation of nerve-immune cell interaction in murine spleen. *J. Neuroimmunol.* **61**, 53–60.
- Sundar, S. K., Becker, K. J., Cierpial, M. A., Carpenter, M. D., Rankin, L. A., Fleener, S. L., Ritchie, J. C., Simpson, P. E. and Weiss, J. M. (1989) Intracerebroventricular infusion of interleukin 1 rapidly decreases peripheral cellular immune responses. *Proc. Natl. Acad. Sci. U.S.A.* **86**, 6398–6402.
- Sundar, S. K., Cierpial, M. A., Kilts, C., Ritchie, J. C. and Weiss, J. M. (1990) Brain IL-1-induced immunosuppression occurs through activation of both pituitary-adrenal axis and sympathetic nervous system by corticotropin-releasing factor. *J. Neurosci.* **10**, 11, 3701–3706.
- Sundar, S. K., Cierpial, M. A., Kamaraju, L. S., Long, S., Hsieh, S., Lorenz, C., Aaron, M., Ritchie, J. C. and Weiss, J. M. (1991)

- Human immunodeficiency virus glycoprotein (gp120) infused into rat brain induces interleukin-1 to elevate pituitary-adrenal activity and decrease peripheral cellular immune responses. *Proc. Natl. Acad. Sci. U.S.A.* **88**, 11246-11250.
- Swanson, L. W., Köhler, C. and Björklund, A. (1987) The limbic region. I. The septohippocampal system. In: *Handbook of Chemical Neuroanatomy*, Vol. 5, Part 1, pp. 125-277. Eds A. Björklund, T. Hökfelt and L. W. Swanson. Elsevier: Amsterdam.
- Tajima, T., Endo, H., Suzuki, Y., Ikari, H., Gotoh, M. and Iguchi, A. (1996) Immobilization stress-induced increase of hippocampal acetylcholine and of plasma epinephrine, norepinephrine and glucose in rats. *Brain Res.* **720**, 155-158.
- Takao, T., Nakata, H., Tojo, C., Kurokawa, H., Nishioka, T., Hashimoto, K. and De Souza, E. B. (1994) Regulation of interleukin-1 receptors and hypothalamic-pituitary-adrenal axis by lipopolysaccharide treatment in the mouse. *Brain Res.* **649**, 265-270.
- Takasu, N., Komiya, I., Nagasawa, Y., Asawa, T. and Yamada, T. (1990) Exacerbation of autoimmune thyroid dysfunction after unilateral adrenalectomy in patients with Cushing's syndrome due to an adrenocortical adenoma. *N. Engl. J. Med.* **322**, 1708-1712.
- Takasu, N., Ohara, N., Yamada, T. and Komiya, I. (1993) Development of autoimmune thyroid dysfunction after bilateral adrenalectomy in a patient with Carney's complex and after removal of ACTH producing pituitary adenoma in a patient with Cushing's disease. *J. Endocrinol. Invest.* **16**, 697-702.
- Tancredi, V., Zona, C., Velotti, F., Eusebi, F. and Santoni, A. (1990) Interleukin-2 suppresses established long-term potentiation and inhibits its induction in the rat hippocampus. *Brain Res.* **525**, 149-151.
- Tancredi, V., D'Arcangelo, G., Grassi, F., Tarroni, P., Palmieri, G., Santoni, A. and Eusebi, F. (1992) Tumor necrosis factor alters synaptic transmission in rat hippocampal slices. *Neurosci. Lett.* **146**, 176-178.
- Taylor, D. C. M. (1982) The effects of nucleus raphe magnus lesions on an ascending thermal pathway in the rat. *J. Physiol. (London)* **326**, 309-318.
- Tice, M. A. B., Hashemi, T., Taylor, L. A. and McQuade, R. D. (1996) Distribution of muscarinic receptor subtypes in rat brain from postnatal to old age. *Dev. Brain Res.* **92**, 70-76.
- Tidey, J. W. and Miczek, K. A. (1996) Social defeat stress selectively alters meso-corticolimbic dopamine release: an *in vivo* microdialysis study. *Brain Res.* **721**, 140-149.
- Tocco, G., Shors, T. J., Baudry, M. and Thompson, R. F. (1991) Selective increase of AMPA binding to the AMPA/quisqualate receptor in the hippocampus in response to acute stress. *Brain Res.* **559**, 168-171.
- Toulmond, S., Vige, X., Fage, D. and Benavides, J. (1992) Local infusion of interleukin-6 attenuates the neurotoxic effects of NMDA on rat striatal cholinergic neurons. *Neurosci. Lett.* **144**, 49-52.
- Tsao, C.-W., Cheng, J.-T., Shen, C.-L. and Lin, Y.-S. (1996) 6-Hydroxydopamine induces thymocyte apoptosis in mice. *J. Neuroimmunol.* **65**, 91-95.
- Udem, B. J., Riccio, M. M., Weinreich, D., Ellis, J. L. and Myers, A. C. (1995) Neurophysiology of mast cell-nerve interactions in the airways. *Int. Arch. Allergy Immunol.* **107**, 199-201.
- Van Dam, A.-M., Brouns, M., Man-A-Hing, W. and Berkenbosch, F. (1993) Immunocytochemical detection of prostaglandin E2 in microvasculature and in neurons of rat brain after administration of bacterial endotoxin. *Brain Res.* **613**, 331-336.
- Van Dam, A.-M., De Vries, H. E., Kuiper, J., Zijlstra, F. J., De Boer, A. G., Tilders, F. J. H. and Berkenbosch, F. (1996) Interleukin-1 receptors on rat brain endothelial cells: a role in neuroimmune interaction?. *FASEB J.* **10**, 351-356.
- Vandenabeele, P. and Fiers, W. (1991) Is amyloidogenesis during Alzheimer's disease due to an IL-1/IL-6-mediated "acute phase response" in the brain?. *Immunol. Today* **12**, 217-219.
- Van Wimersma Greidanus, T. J. B. and Maigret, C. (1996) The role of limbic vasopressin and oxytocin in social recognition. *Brain Res.* **713**, 153-159.
- Veith, R. C., Lewis, N., Linares, O. A., Barnes, R. F., Raskind, M. A., Villacres, E. C., Murburg, M. M., Ashleigh, E. A., Castillo, S., Peskind, E. R., Pascualy, M. and Halter, J. B. (1994) Sympathetic nervous system activity in major depression. Basal and desipramine-induced alterations in plasma norepinephrine kinetics. *Arch. Gen. Psychiat.* **51**, 411-422.
- Wakabayashi, K., Honer, W. G. and Masliah, E. (1994) Synapse alterations in the hippocampal-entorhinal formation in Alzheimer's disease with and without Lewy body disease. *Brain Res.* **667**, 24-32.
- Wallace, D. M., Magnuson, D. J. and Gray, T. S. (1992) Organization of amygdaloid projections to brainstem dopaminergic, noradrenergic, and adrenergic cell groups in the rat. *Brain Res. Bull.* **28**, 447-454.
- Wan, W., Vriend, C. Y., Wetmore, L., Gartner, J. G., Greenberg, A. H. and Nance, D. M. (1993) The effects of stress on splenic immune function are mediated by the splenic nerve. *Brain Res. Bull.* **30**, 101-105.
- Watanabe, Y., Wailand, N. G. and McEwen, B. S. (1995a) Effects of adrenal steroid manipulations and repeated restraint stress on dynorphin mRNA levels and excitatory amino acid receptor binding in hippocampus. *Brain Res.* **680**, 217-225.
- Watanabe, Y., Saito, H. and Abe, K. (1995b) Nitric oxide is involved in long-term potentiation in the medial but not lateral amygdala neuron synapses *in vitro*. *Brain Res.* **688**, 233-236.
- Watkins, L. R., Wiertelak, E. P., Goehler, L. E., Mooney-Heiberger, K., Martinez, J., Furness, L., Smith, K. P. and Maier, S. F. (1994a) Neurocircuitry of illness-induced hyperalgesia. *Brain Res.* **639**, 283-299.
- Watkins, L. R., Wiertelak, E. P., Goehler, L. E., Smith, K. P., Martin, D. and Maier, S. F. (1994b) Characterization of cytokine-induced hyperalgesia. *Brain Res.* **654**, 15-26.
- Watkins, L. R., Wiertelak, E. P., Furness, L. E. and Maier, S. F. (1994c) Illness-induced hyperalgesia is mediated by spinal neuropeptides and excitatory amino acids. *Brain Res.* **664**, 17-24.
- Watkins, L. R., Goehler, L. E., Reltton, J., Brewer, M. T. and Maier, S. F. (1995a) Mechanisms of tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) hyperalgesia. *Brain Res.* **692**, 244-250.
- Watkins, L. R., Goehler, L. E., Reltton, J. K., Taraglia, N., Silbert, L., Martin, D. and Maier, S. F. (1995b) Blockade of interleukin-1 induced hyperthermia by subdiaphragmatic vagotomy: evidence for vagal mediation of immune-brain communication. *Neurosci. Lett.* **183**, 27-31.
- Weber, R. J. and Pert, A. (1989) The periaqueductal gray matter mediates opiate-induced immunosuppression. *Science* **245**, 188-190.
- Weeks, J. C. and Levine, R. B. (1995) Steroid hormone effects on neurons subserving behavior. *Curr. Opin. Neurobiol.* **5**, 809-815.
- Weidenfeld, J., Crumeyrolle-Arias, M. and Haour, F. (1995) Effect of bacterial endotoxin and interleukin-1 on prostaglandin biosynthesis by the hippocampus of mouse brain: role of interleukin-1 receptors and glucocorticoids. *Neuroendocrinology* **62**, 39-46.
- Werner, J. and Bienc, A. (1985) The significance of nucleus raphe dorsalis and centrals for thermoafferent signal transmission to the preoptic area of the rat. *Exp. Brain Res.* **59**, 543-547.
- Werner-Felmayer, G., Werner, E. R., Fuchs, D., Hausen, A., Reibnegger, G. and Wachter, H. (1989) Characteristics of interferon induced tryptophan metabolism in human cells *in vitro*. *Biochim. Biophys. Acta* **1012**, 140-147.
- Wetmore, L., Green-Johnson, J., Gartner, J. G., Sanders, V. and Nance, D. M. (1994) The effects of kainic acid-induced lesions in the lateral septal area on cell-mediated immune function. *Brain Behav. Immun.* **8**, 341-354.
- Whittemore, S. R. and Sieger, A. (1987) The expression, localization and functional significance of  $\beta$ -nerve growth factor in the central nervous system. *Brain Res. Rev.* **12**, 439-464.
- Wiertelak, E. P., Furness, L. E., Watkins, L. R. and Maier, S. F. (1994) Illness-induced hyperalgesia is mediated by a spinal NMDA-nitric oxide cascade. *Brain Res.* **664**, 9-16.
- Wilkinson, M. F., Mathieson, W. B. and Pittman, Q. J. (1993) Interleukin-1 $\beta$  has excitatory effects on neurons of the bed nucleus of the stria terminalis. *Brain Res.* **625**, 342-346.
- Woolley, C. S. and McEwen, B. S. (1994) Estradiol regulates hippocampal dendritic spine density via an N-methyl-D-aspartate receptor-dependent mechanism. *J. Neurosci.* **14**, 7680-7687.
- Yamada, M. and Hatanaka, H. (1994) Interleukin-6 protects cultured rat hippocampal neurons against glutamate-induced cell death. *Brain Res.* **643**, 173-180.
- Yu, B. and Shinnick-Gallagher, P. (1994) Interleukin-1 $\beta$  inhibits synaptic transmission and induces membrane hyperpolarization in amygdala neurons. *J. Pharmacol. Exp. Ther.* **271**, 590-600.
- Zachrisson, O., Mathé, A. A., Stenfors, C. and Lindfors, N. (1995)

- Limbic effects of repeated electroconvulsive stimulation on neuropeptide Y and somatostatin mRNA expression in the rat brain. *Mol. Brain Res.* **31**, 71-85.
- Zalcman, S., Shanks, N. and Anisman, H. (1991) Time-dependent variations of central norepinephrine and dopamine following antigen administration. *Brain Res.* **557**, 69-76.
- Zalcman, S., Green-Johnson, J. M., Murray, L., Wan, W., Nance, D. M. and Greenberg, A. H. (1994a) Interleukin-2-induced enhancement of an antigen-specific IgM plaque-forming cell response is mediated by the sympathetic nervous system. *J. Pharmacol. Exp. Ther.* **271**, 977-982.
- Zalcman, S., Green-Johnson, J. M., Murray, L., Nance, D. M., Dyck, D., Anisman, H. and Greenberg, A. H. (1994b) Cytokine-specific central monoamine alterations induced by interleukin-1, -2 and -6. *Brain Res.* **643**, 40-49.
- Zeise, M. L., Madamba, S. and Siggins, G. R. (1992) Interleukin-1 $\beta$  increases synaptic inhibition in rat hippocampal pyramidal neurons *in vitro*. *Regul. Peptides* **39**, 1-7.
- Zelazowski, P., Patchev, V. K., Zelazowska, E. B., Chrousos, G. P., Gold, P. W. and Sternberg, E. M. (1993) Release of hypothalamic corticotropin-releasing hormone and arginine-vasopressin by interleukin-1 $\beta$  and  $\alpha$ -MSH: studies in rats with different susceptibility to inflammatory disease. *Brain Res.* **631**, 22-26.
- Zhong, P. and Ciaranello, R. D. (1995) Transcriptional regulation of hippocampal 5-HT<sub>1A</sub> receptors by corticosteroid hormones. *Mol. Brain Res.* **29**, 23-34.
- Zhou, D., Kusnecov, A. W., Shurin, M. R., DePaoli, M. and Rabin, B. S. (1993) Exposure to physical and psychological stressors elevates plasma interleukin 6: relationship to the activation of hypothalamic-pituitary-adrenal axis. *Endocrinology* **133**, 2523-2530.