

Rehabilitation Approaches to Hemineglect

Randolph S. Marshall, MD, MS

Background: Hemineglect is a difficult neurologic condition to rehabilitate. It arises predominantly from right brain injury, and manifests heterogeneously in clinical deficits such as poor visual exploration to the left, inaccurate assessment of the midpoint of a line, left limb hypokinesia, and anosognosia. Most of the cognitive dysfunction produced by hemineglect is because of an asymmetric distribution of attention, either with respect to extrapersonal space or to an object being viewed. Many treatments draw on hemineglect theory to attempt to mediate the basic asymmetry of attention.

Review Summary: Treatment approaches can be divided into 2 main categories. Extrinsic or “top-down” approaches require active participation of the patient under the guidance of a therapist. The most common approach of this type is visual scanning therapy in which the patient is continually instructed to move the gaze leftward into the neglected space. Intrinsic or “bottom-up” approaches manipulate stimulus characteristics, sensory input, or the brain directly in an attempt to alter the interhemispherical attentional imbalance. Examples of this approach include vestibular stimulation of the left side, sensory activation of the left limb, and transcranial magnetic stimulation of the overactive left hemisphere. Combined approaches such as prism adaptation have also shown good results.

Conclusions: Hemineglect is a complicated disorder that poses challenges to treatment. A paucity of clinical trial evidence limits our ability to extrapolate experimental mediation of hemineglect to globally improved functioning. Nonetheless, many treatment approaches appear promising. Underlying neuroscience may help guide future treatment approaches.

Key Words: hemineglect, rehabilitation, hemi-inattention, visual scanning therapy, prism adaptation

(*The Neurologist* 2009;15: 185–192)

Hemineglect is a heterogeneous syndrome that can produce major disability in patients who acquire the disorder from brain injury such as stroke. Its presence is vexing for rehabilitation specialists, as the elements of the syndrome often prevent full participation in therapy. Amelioration of hemineglect is crucial not only to reduce the basic visual-spatial dysfunction, but also to maximize engagement in other aspects of therapy such as motor recovery, gait training, and general functional recovery.^{1–3} This review surveys the major treatment approaches to hemineglect, which have been tested over the last 50 years. Both rationally designed treatments based on underlying theories of neglect and purely empirical approaches will be discussed. Proof of efficacy has been difficult because few well-designed randomized clinical trials have been completed, but there is compelling evidence that several approaches to hemineglect treatment are useful.

CLINICAL AND THEORETICAL ASPECTS OF HEMINEGLECT

The reported incidence of hemineglect in stroke patients is approximately 23%, ranging from 8% to 81%, depending on definitions, patient selection, and assessment tools.^{4,5} Neglect is much more common in right brain injury than left⁶; right hemispherical predominance is most likely because of specialized functions of the right hemisphere as opposed to an inability to measure hemineglect due to aphasia produced by left hemisphere injury.

The underlying deficit in hemineglect is inattention to 1 hemispace. Clinically, one can identify a variety of deficits which may occur in isolation or in combination, but all arise from the basic asymmetry in attentional field. Hemineglect may manifest as “egocentric” (viewer-centered reference frame) or “allocentric” (object-centered reference frame). Examples of egocentric manifestations include unilateral extinction to double simultaneous sensory stimulation, thought to occur as the result of the inability of the left stimulus to compete for salience when the attention to the right hemispace contains an equivalent stimulus.⁷ Asymmetric orienting response similarly occurs when a patient orients his or her gaze to the non-neglected field when stimuli are presented to both sides.⁸ This obligatory shift in orientation may be because of both reduced attention to the neglected side as well as to an inability to disengage from stimuli in the non-neglected field.⁹ Asymmetric drawing or copying and right-shifted marking of the midpoint on a line bisection test are examples of allocentric neglect. In such a deficit, the object is perceived asymmetrically regardless of its location in egocentric space. Another way of classifying the different deficits is by “sensory/perceptual” neglect versus “motor/exploratory” neglect.¹⁰ Most manifestations of egocentric neglect are motor/exploratory, and those with an allocentric base are generally sensory. The terms “intentional” and “attentional,” respectively have been applied to this dichotomy as well.^{11,12} Unilateral limb hypokinesia is a demonstration of motor neglect that affects peripersonal space or body perception.^{13,14} Asomatognosia is another example of this type of distorted hemi-inattention of self.¹⁵ For most patients the hemineglect syndrome clusters its individual deficits according to the basic divisions described here. Any single element or combination, however, may be seen in a given patient.

The underlying deficit in hemineglect is inattention to 1 hemispace.

In addition to the elements clearly caused by some sort of hemi-inattention, there are often additional, more global deficits of cognitive function that accompany hemineglect to make the treatment additionally challenging. Anosognosia is seen in hemineglect patients in up to 20% to 58% of cases.^{16,17} A lack of self-monitoring because of the unawareness of the deficit makes cooperative, mindful work with a therapist difficult. Other higher order cognitive dysfunction also may occur from right brain injury. Personality change can range from apathy to disinhibition to agitation.¹⁸ Emo-

From the Stroke Division, The Neurological Institute of New York, Columbia University Medical Center, New York, New York.

Reprints: Randolph S. Marshall, MD, MS, Stroke Division, The Neurological Institute of New York, Columbia University Medical Center, 710 w 168th St., New York, NY 10032. E-mail: rsm2@columbia.edu.

Copyright © 2009 by Lippincott Williams & Wilkins
ISSN: 1074-7931/09/1504-0185

DOI: 10.1097/NRL.0b013e3181942894

tional dysregulation is common, often appearing in the setting of the impaired self-monitoring.¹⁹ Inability to multitask has also been recently reported.²⁰ Finally, although patients' perception may be abnormal, and it can be pointed out to them (eg, this arm that you claim is not yours can be traced back to your body), they are often unable to reconcile logic of the situation with abnormal perceptions. One patient, for example, reported that his hospital room was the "only one to have telephones with 2 columns of numbers." On a line bisection task, a hemineglect patient may be trained to move the gaze leftward to identify a cue at the left extent of a line, confirming that the whole line is in view, and yet, that patient may still bisect the line to the right of midpoint as the gaze tracks rightward.^{21,22} Furthermore, abnormal use information about stimuli depending on its context has been demonstrated.²³ Line bisection behavior on a given length line was shown to be altered by the introduction of different length lines—the perceived midpoint shifting leftward on that line when longer lines were mixed in the set and the perceived midpoint shifting rightward when shorter lines were mixed in. This "context effect" seems to be unique to right brain injury.

A lack of self-monitoring because of the unawareness of the deficit makes cooperative, mindful work with a therapist difficult.

Another demonstration of the dissociation between perception and logic was reported by Ramachandran et al.²⁴ Twelve patients with left hemineglect caused by right hemisphere stroke were approached at the bedside on their right side with a mirror. All patients were able to identify the object as a mirror. The investigator stood behind the mirror, holding it. A second investigator, standing on the patient's left side, held out a pen or a candy close enough that the patient would be able to reach it with the nonparetic right hand, but so that it could be seen in the mirror. When the patient affirmed that the object was seen, he or she was asked to reach out and grab it. A substantial subset of patients, despite acknowledging the presence of the mirror, repeatedly reached for the object's reflection on the right, even going so far as to say, "it's behind the mirror." Thus right brain injury results in more than just a hemi-inattention. The layering on of other cognitive deficits adds complexity to the syndrome, producing additional challenges to devising effective treatments.

The taxonomy of treatment approaches is based on the phenomenology and underlying theories of hemineglect. Brain (1941),²⁵ Denny-Brown et al (1954),²⁶ and Critchley (1950)²⁷ promoted the role of the parietal lobes in integrating and synthesizing sensory information. Bisiach et al (1983)²⁸ theorized the deficit as an abnormal internal representation, suggesting that hemineglect patients had a defect in transforming retinotopic to cephalocentric to corpocentric reference frames. Kinsbourne (1977)²⁹ and later Heilman and Van den Abell (1980),³⁰ Chatterjee (1995),³¹ and Anderson (1996),³² held that hemineglect was explained by an asymmetric orientation in extrapersonal space. The thought was that gaze and attention were asymmetrically represented, with the right hemisphere directing attention both contralaterally and ipsilaterally, and the left hemisphere directing attention contralaterally only. Thus, with a left brain injury attention could still be directed to the right with the right hemisphere, but with right brain injury the left side of space could not be accessed ipsilaterally with the intact left hemisphere. Others have stressed the importance of poor general atten-

tional arousal that occurs with right brain injury as an important component of the neglect syndrome.³³ Clinically, global inattention often appears as part of the syndrome and establishes itself as the most significant limitation on intervention. The dopaminergic system mediates general attention,³⁴ and its receptors seem to be asymmetrically lateralized to the right hemisphere.³⁵

Clinically, global inattention often appears as part of the syndrome and establishes itself as the most significant limitation on intervention.

As most clinical observations and psychophysical studies support the idea of asymmetric attention being the most important underlying deficit in hemineglect, the physiologic basis for hemispherical asymmetry has been elusive. In a functional imaging study by Corbetta et al, 11 patients with right ventral frontal brain injury and left hemineglect underwent imaging in the acute, impaired phase and upon recovery.³⁶ In-scanner performance of a directional cue target detection task resulted in an abnormal balance of activity in fronto-parietal regions remote from the lesions. Relative hyperactivity in the left superior parietal lobule compared with homologous regions in the right hemisphere correlated with missed targets and slowness of response in left visual space. As neglect spontaneously improved in these patients, a more normal balance of activity in both the dorsal and ventral regions was observed.

Beyond the notion of hemispherical asymmetry, intrahemispheric anatomic specificity has also been shown to apply to subtypes of hemineglect, with perceptual deficits correlating with posterior parietal injury and motor deficits correlating more with frontal lesions.^{10,37,38} In 1 study, Binder et al¹⁰ divided 35 patients with right brain injury into 2 behavioral categories, those who had markedly asymmetric performance on a target cancellation test but performed in the normal range on a line bisection task, and those who had an abnormal line bisection test and were relatively normal on the cancellation task. When the lesions of those with cancellation abnormality only were overlapped on an anatomic template and the lesions of those who did not fit in that category were subtracted from the overlap, the resulting lesion location was frontal opercular, anterior insular, and basal ganglia. When the group with line bisection abnormalities was assessed in the same way, the temporo-parietal-occipital junction was identified as relevant.

CATEGORIZATION OF TREATMENT APPROACHES

Proposed approaches to treat hemineglect have been available for more than 50 years. The treatment types can be divided into basic categories, each of which will be discussed in this review. One approach can be classified as extrinsic or "top-down," meaning that external cues and guidance are employed predominantly to engage the conscious and purposeful involvement of the patient. These treatments rely heavily on the participation of a therapist who provides continuous feedback, encouragement, and training. Cooperation and conscious effort on the part of the patient is required for success. Treatments that follow such an approach include visual scanning training (VST),^{39–46} sustained attention training,^{47,48} and mental imagery training.^{49–51} A second approach to treatment attempts to affect the attentional system at a preconscious level by manipulating endogenous components of the neural axis, mostly sensory input. This approach follows the idea that a disordered perceptual and attentional system can be corrected without the

TABLE 1. Classification of Treatment Approaches

Treatment Approach	Examples of Therapy	References
Extrinsic or “top-down”	Visual scanning training (VST)	21,38–45
	Sustained attention training (SAT)	46,47
	Mental imagery training (MIT)	48–50
Intrinsic or “bottom-up”	Prism glasses	51
	Phasic alerting	84
	Trunk rotation	43,52,53
	Eye patching	54–56
	Limb activation	21,47,57–59,107–109
	Optokinetic/caloric stimulation	64,65,89–93
	Transcutaneous electrical stimulation	87,88
	Neck muscle vibration	60–63,94
Combined	Direct brain stimulation (rTMS)	80–82
	Prism adaptation	66–75,96,97
	Stimulation plus VST	63,110

conscious participation of the patient, and thus have been termed “intrinsic,” or bottom-up, treatments. Such treatments may manipulate visual-spatial input such as with prism glasses,⁵² trunk rotation,^{44,53,54} or eye patching.^{55–57} Others alter the somatosensory system such as with limb activation,^{22,58–60} neck muscle vibration,^{61–64} and optokinetic stimulation.^{65,66} A combination of top-down and bottom-up approaches has also been used, exemplified by the prism adaptation approach,^{67–76} which will be discussed in more detail below. A final treatment approach attempts to alter brain function directly, either with pharmacologic agents^{77–80} or with brain stimulation using repetitive transcranial magnetic stimulation (rTMS).^{81–83} A Table 1 of treatment approach classification is provided below. Individual treatments warrant particular attention.

Extrinsic or Top-Down Treatments

VST is the most popular and widely used treatment for hemineglect. Part of its appeal is the obvious relationship between an observable deficit—inability to find and identify objects in the neglected hemispace and the intervention—active directions to search the left hemispace. Family members and caretakers can easily grasp the concept and are able to assist in the effort. This approach can be done with no special equipment or setting, working on the principle that a preserved verbal language system in the intact left hemisphere can be engaged to mediate the visual-spatial deficits created by the injured right hemisphere. At its most basic, the technique involves a therapist sitting to the patient’s neglected side to issue instructions urging scanning of the eyes leftward throughout training. As with most extrinsic therapy, the program is methodical and progressive. In a reading exercises, for example, an anchoring cue such as a red line is placed on the left side of the page, and the patient is asked to look at the cue before beginning the text reading.²² The text advances from simple to complex, with repeated scanning instructions and visual cues used throughout. Mental imagery is often employed, such as the instructions: “Imagine you are a lighthouse like the one you see in this picture. Let your eyes sweep the environment like a lighthouse beam from right to left (Fig. 1).” Patients are encouraged to self-direct the imagery, verbally repeating the lighthouse strategy during the course of therapy. The lighthouse strategy was tested formally by Niemeier et al.⁵¹ Sixteen patients with left hemineglect from right brain stroke were given the lighthouse strategy over the course of therapy and compared with




FIGURE 1. Lighthouse strategy of mental imagery for use with visual scanning therapy.

identical patients who were not given this training. The investigators showed that the patients trained with lighthouse strategy were significantly better on the Mesulam target cancellation test as well as on the attention item on the Functional Autonomy Rating Scale ($P < 0.007$ on repeated measures ANOVA). Another study combined VST with trunk rotation.⁴⁴ In this study by Wiart et al, 22 hemineglect patients were subject to either traditional physical and occupational training or VST with “trunk rotation.” For the trunk rotation, patients wore a harness with a pointer attached to an arm that extended from the back up over the head and down to a point just above the visual horizontal meridian. Patients in the experimental group were instructed in a scanning strategy to bring both the eyes (retinocentric frame) and the pointer (body-centric frame) to the left in search of objects to the left of midline. Compared with controls, patients receiving the visual scanning and trunk rotation improved on 3 neglects tests and the Functional Independence Measure (FIM), and these results seemed to persist after completing the 1 month of training (Fig 1., lighthouse).

VST is the most popular and widely used treatment for hemineglect.

Intrinsic or Bottom-Up Approaches

Unlike the top-down approaches, bottom-up strategies take advantage of the brain’s automatic responses to stimuli. The appeal

of this approach is that it is not therapist-dependent, and can be set up in a standardized way in a variety of settings—as home therapy or in the rehabilitation clinic.

Treatments of General Arousal

Dopamine agonists have been shown to ameliorate rightward bias on letter cancellation and line bisection tests in small case series,^{77,79,80} but evidence exists on the other side that dopamine agonists may worsen exploratory bias in hemineglect,⁸⁴ possibly because the increased dopamine activates the undamaged hemisphere and increases the exploratory bias from the intact hemisphere.⁷⁸

Behavioral methods have also been used to mediate the general arousal deficit of hemineglect. In a study of “phasic alerting,” Robertson et al⁸⁵ studied 8 patients with left hemineglect who were asked to judge whether a visual stimulus on the left preceded or followed a stimulus on the right. On average, perception of left stimuli occurred 500 milliseconds after stimuli on the right. This delayed perception was lessened significantly by presenting an auditory warning in the form of a 300-millisecond tone-burst. On trials in which the warning occurred, each patient reduced the rightward bias. The effect occurred regardless of whether the auditory stimulus was presented to the left or right, thus demonstrating that increasing general alertness can ameliorate a lateralizing visuospatial bias.

Another interesting study was conducted with 7 stroke patients engaged in a car racing computer game requiring increasing alertness to avoid obstacles.⁸⁶ The investigators showed that patients improved on several alertness measures and neglect tests, but the performance reverted to pretreatment levels by 4 weeks post-treatment. Interestingly, there was an fMRI correlate which, showed increased activity in the right angular gyrus, bilateral anterior cingulate cortex, and precuneus, but which also faded along with reversion of performance at the later time.

Sensory Activation

Based on the theory that hemineglect is a state of asymmetric sensory and motor attention, many treatment strategies have focused on increasing the sensory input to the contralesional left body in an attempt to give the damaged right hemisphere additional activation.^{48,58–60} Wilson et al, for example, showed improvements on a self-care program and a picture scanning task in 2 patients tested before and after a limb activation treatment.⁴⁸ The technique works simply by asking the patient to move the left limb (upper preferred) during the performance of other tasks. Although the technique has theoretical merit if personal body representation does indeed share a neural network with peripersonal space,⁸⁷ this approach may have limitations for severely paretic patients. Further, movement of the nonparetic right hand could work against the desired activation of the right hemisphere by activating the left preferentially. More deliberate activation of left body sensory input has been tested in the form of neck muscle vibration with transcutaneous electrical stimulation,^{88,89} or the induction of left nystagmus with optokinetic stimulation^{90–92} or caloric stimulation.^{93,94} In 1 experiment, Kamath et al showed that a visual search pattern could be shifted leftward by cold water irrigation of the left ear or left sided neck muscle vibration.⁹⁵ Either stimulation seemed to have the effect of shifting egocentric reference frame leftward such that left hemineglect patients acquired the search pattern of normal controls, and control subjects had a leftward search bias induced so that they appeared to have right hemineglect. Such techniques have been shown to alter performance on neglect tasks, but seem to have limited persistence beyond the therapy session, and do not seem to generalize to improved performance on activities of daily living.

Direct Brain Stimulation

After hemispheric activity imbalance theory, techniques have been devised to alter the balance of activity in the 2 hemispheres directly. rTMS has been used over the left parietal lobe to try to deactivate that hemisphere and bring the attentional bias into balance. In 1 experiment by Oliveri et al, rTMS administered over the unaffected (left) parietal cortex reduced the line bisection abnormality during the testing session, but the results did not persist.⁸¹ Other rTMS studies have found similar results.^{82,83}

Visual Remapping

It has been known for more than a century that prism lenses can remap visual space onto the retina to result in a systematic shift of retinotopic coordinates. Fresnel lenses have been used with some success to create a shift of objects that appear further to the right than they actually are, thus shifting retinotopic space onto new coordinates of body-centered space. This technique improved performance in a group of hemineglect patients who wore the lenses for 4 weeks during therapy and also during testing before and after the treatment.⁵² Eye patching of the right eye has also been attempted to alter the visual perceptual input in an attempt to shift attentional bias. This approach is based on the rationale that each superior colliculus receives input from the contralateral hemifield of each eye, but predominantly from the contralateral eye.⁹⁶ In 1 study of 22 stroke patients comparing the eye-patching technique with the wearing of glasses that blocked the right hemifield in each eye, both groups improved on 3 tests of neglect,⁵⁷ but it is uncertain whether this approach can result in improved function of activities of daily living.

Combined Top-Down and Bottom-Up Approaches

One promising method that combines therapist driven extrinsic and sensory-motor transformation intrinsic systems is prism adaptation. This interesting approach, introduced by Rossetti et al in 1998,⁷⁶ takes advantage of the well-known short-term adaptation that takes place when subjects are exposed to prism lenses that produce a shift of the visual field. In this study, 8 patients with left hemineglect and 6 controls were asked to make 10 straight ahead pointing movements while blindfolded. Hemineglect patients pointed to the right of true midline. Patients and controls were then fitted with base-left prism glasses that shifted their visual fields 10 degrees to the right. During the adaptation period they were asked to make 50 pointing movements at objects presented 10 degrees to the left or right of the body midline. The training occurred as the hand of the subject appeared in the second half of the pointing movement, and a leftward correction was made, under the guidance and instructions of the investigator/therapist. Following this 2- to 5-minute training period, the prism goggles were removed and the subjects again pointed straight ahead while blindfolded. The resulting adaptation training produced a leftward shift on the subjective midpoint, correcting the rightward bias of the hemineglect patients to normal, and shifting the normal midpoint to the left in controls (see Fig. 2). In a separate experiment, 3 tests of hemineglect (line bisection, line cancellation, and a copying task) showed improvements on all 3 tasks following prism adaptation, a result which, interestingly, persisted 2 hours after the goggles had been removed. Subsequent studies have shown that exposures as brief as 5 to 7 minutes can produce amelioration of hemineglect that lasts up to 24 hours,⁹⁷ and twice daily adaptation sessions over a 2-week period can result in improvements in neglect of up to 5 weeks duration.⁷⁴ This prism adaptation has theoretical appeal because it cannot be due simply to a sensory or perceptual shift, but rather must alter in a lasting way central processes related to sensorimotor correspondences, perhaps stimulating the left




FIGURE 2. Prism adaptation. Patients wear prisms that shift images rightward. Pretherapy patients' straight-ahead pointing results in a shift to the right. During therapy with the prisms on, patients are trained to point straight ahead under visual guidance. When prisms are removed post-therapy, straight-ahead pointing is shifted leftward toward the true midline.

hemisphere networks demonstrated to be active in a recovery of hemineglect.³⁶ As dramatic as the results seem in prism adaptation, the treatment even so may work on only a component of hemineglect. As shown by a study by Ferber et al,⁹⁸ prism adaptation improved the exploratory eye movements of a chimeric face, but did not change the abnormal perception of the unitary object. Patients still made judgments based on the right side of the face predominantly, and also noted nothing unusual about the chimeric faces, nor of the experience of prism adaptation. The limited treatment effect speaks once again to the complex, multifaceted nature of the condition, and demonstrates how difficult hemineglect can be to treat.

One promising method that combines therapist driven extrinsic and sensory-motor transformation intrinsic systems is prism adaptation.

Efficacy of Neglect Treatments

Most of the studies described in this review have had strong theoretical foundations and were carefully designed. Results are compelling for the small case series and single subject comparisons. Efficacy, however, based on controlled clinical trials, and demonstrating lasting improvements at the level of impairment, and generalization to tasks of daily living, has been more difficult to prove. A few critical, systematic reviews of hemineglect treatments have been published in the last few years.^{99–101} These reviews set criteria for efficacy that included: (1) the impact of treatment on disability as measured by general disability scales such as the FIM, the behavioral subtests of the behavioral inattention test (BIT),¹⁰² and the Barthel Index;^{103,104} (2) the long-term effects of the treatment beyond the therapy program, and (3) the quality of evidence [eg, level 1 is a well-designed randomized controlled trial (RCT); level 2 a cohort study with a comparison group; level 3 a case-control, cross-over, or well-designed single-case study; level 4 are studies without

specific assessment of recovery or long-term effect). In the review by Luaute et al,⁹⁹ 116 articles were reviewed, 54 of which were included in the analysis. They noted 6 level 1 studies supporting the use of VST as demonstrating improvement on functional skills such as reading and writing,^{39,42,45,46,105} and demonstrating generalization of training to other activities of daily living.^{39,105,106} A study by Wiart et al combining VST and trunk rotation was a level 1 study that demonstrated both functional improvement on the FIM and a lasting effect.⁴⁴ Limb activation was supported by 2 level 1 RCTs and 6 level 3 articles. The RCTs showed that motor function was improved by LA,¹⁰⁷ but the results did not translate to ADL measures. All the single case studies showed that LA produced gains in ADL areas such as reading, walking, dressing, and feeding, and a long-term effect was reported in 4 studies.^{48,108–110} Sensory stimulation and direct brain stimulation were found to produce only a within-therapy improvement of hemineglect, and were therefore not recommended as isolated treatment modalities. The efficacy of stimulation techniques may be realized, however, when combined with VST, which may enhance VST's effect.^{64,111} Mental imagery⁴⁹ and feedback training^{112,113} had no level 1 studies to support their use, but lower-level studies seemed strong enough to warrant further investigation in RCTs. Prism adaptation was also cited as a potentially beneficial treatment, warranting further study. Several subsequent studies have shown more lasting and generalizing effects. In addition, neuroimaging studies have proposed an anatomic substrate for the effects of prism adaptation.^{70,114} Luaute et al studied 5 hemineglect patients who performed a line judgment task and a simple observation task during a PET scan before and after 2 to 5 minutes of prism adaptation.⁷⁰ The right cerebellum, left thalamus, and left temporo-occipital regions correlated positively with BIT scores, and left medial temporal and right posterior parietal regions correlated negatively with BIT improvement, suggesting involvement of these regions in the adaptation process.

A recent Cochrane meta-analysis by Bowen and Lincoln focused on controlled clinical trials, evaluating 400 patients in 15 studies. Eight of the 15 were randomized controlled trials; the others had unclear or no randomization. They found that only 6 of the studies addressed functional recovery, and only 3 of them demonstrated favorable outcomes in the treatment group. Showing benefit in their opinion were a visual cuing intervention by Kalra et al,¹⁰⁷ a VST study by Paolucci,¹⁰⁵ and the combined VST and trunk rotation study by Wiart et al.⁴⁴ This last study was the only 1 they cited that showed persistent effects over time, although results were questioned because of poor case matching (experimental group was younger and had a higher baseline FIM than the control group). The overall conclusion of this Cochrane review was that the effectiveness of cognitive rehabilitation for neglect remains unproven, but that there is enough compelling evidence to warrant further investigation with well-designed controlled clinical trials.

CONCLUSIONS

Hemineglect is a heterogeneous disorder that requires tailored treatment strategies. Most clinicians who treat hemineglect take a practical approach to treatment that involves assessing the individual deficits of the patient and devising a treatment strategy that addresses the individual's syndrome. Complete treatment approaches generally involve a combination of top-down and bottom-up approaches, with robust involvement of a therapist during the course of therapy. Some form of VST seems to be effective in most cases of neglect as long as the hemineglect includes difficulty exploring leftward into the neglected hemisphere. Prism adaptation may be

recommended for exploratory/gaze abnormalities as well, and alertness or sustained attention training may be useful for perceptual dysfunction. Further RCTs are warranted, but with restricted inclusion/exclusion criteria and outcome measures that match the appropriate syndrome subset.

ACKNOWLEDGMENT

Authors thank to Brandon Minzer for assistance with figures and manuscript preparation.

REFERENCES

- Cherney LR, Halper AS, Kwasnica CM, et al. Recovery of functional status after right hemisphere stroke: relationship with unilateral neglect. *Arch Phys Med Rehabil.* 2001;2:322–328.
- Jackson D, Thornton H, Turner-Stokes L. Can young severely disabled stroke patients regain the ability to walk independently more than three months post stroke? *Clin Rehabil.* 2000;14:538–547.
- Kinsbourne M. Mechanisms of neglect: implications for rehabilitation. *Neuropsychol Rehabil.* 1994;4:151–153.
- Bowen A, McKenna K, Tallis RC. Reasons for variability in the reported rate of occurrence of unilateral spatial neglect after stroke. *Stroke.* 1999;30:1196–1202.
- Stone SP, Wilson B, Wroot A, et al. The assessment of visuo-spatial neglect after acute stroke. *J Neurol Neurosurg Psychiatry.* 1991;54:345–350.
- Gainotti G. Emotional behavior and hemispheric side of the lesion. *Cortex.* 1972;8:41–55.
- Brozzoli C, Demattè ML, Pavani F, et al. Neglect and extinction: within and between sensory modalities. *Restor Neurol Neurosci.* 2006;24:217–232.
- Gainotti G, D'Erme P, Bartolomeo P. Early orientation of attention toward the half space ipsilateral to the lesion in patients with unilateral brain damage. *J Neurol Neurosurg Psychiatry.* 1991;54:1082–1089.
- Posner MI. Cognitive neuropsychology and the problem of selective attention. *Electroencephalogr Clin Neurophysiol.* 1987;(suppl 39):313–316.
- Binder J, Marshall R, Lazar R, et al. Distinct syndromes of hemineglect. *Arch Neurol.* 1992;49:1187–1194.
- Coslett HB, Bowers D, Fitzpatrick E, et al. Directional hypokinesia and hemispacial inattention in neglect. *Brain.* 1990;113(pt 2):475–486.
- Schwartz RL, Barrett AM, Kim M, et al. Ipsilesional intentional neglect and the effect of cueing. *Neurology.* 1999;53:2017–2022.
- Watson RT, Miller BD, Heilman KM. Nonsensory neglect. *Ann Neurol.* 1978;3:505–508.
- Meador KJ, Watson RT, Bowers D, et al. Hypometria with hemispacial and limb motor neglect. *Brain.* 1986;109(pt 2):293–305.
- Feinberg TE, Haber LD, Leeds NE. Verbal asomatognosia. *Neurology.* 1990;40:1391–1394.
- Berti A, Ladavas E, Della Corte M. Anosognosia for hemiplegia, neglect dyslexia, and drawing neglect: clinical findings and theoretical considerations. *J Int Neuropsychol Soc.* 1996;2:426–440.
- Cutting J. Study of anosognosia. *J Neurol Neurosurg Psychiatry.* 1978;41:548–555.
- Motomura N, Sawada T, Inoue N, et al. Neuropsychological and neuropsychiatric findings in right hemisphere damaged patients. *Jpn J Psychiatry Neurol.* 1988;42:747–752.
- Visser-Keizer AC, Meyboom-de Jong B, Deelman BG, et al. Subjective changes in emotion, cognition and behavior after stroke: factors affecting the perception of patients and partners. *J Clin Exp Neuropsychol.* 2002;24:1032–1045.
- Lazar RM, Festa JR, Geller AE, et al. Multi-tasking disorder from right temporo-parietal stroke. *Cog Behav Neurol.* 2007;20:157–162.
- Riddoch MJ, Humphreys GW. The effect of cueing on unilateral neglect. *Neuropsychologia.* 1983;21:589–599.
- Bailey MJ, Riddoch MJ, Crome P. Treatment of visual neglect in elderly patients with stroke: a single-subject series using either a scanning and cueing strategy or a left-limb activation strategy. *Phys Ther.* 2002;82:782–797.
- Marshall RS, Lazar RM, Krakauer JW, et al. Stimulus context in hemineglect. *Brain.* 1998;121(pt 10):2003–2010.
- Ramachandran VS, Altschuler EL, Stone L, et al. Can mirrors alleviate visual hemineglect? *Med Hypotheses.* 1999;52:303–305.
- Brain W. Visual disorientation with special reference to lesions of the right cerebral hemisphere. *Brain.* 1941;64:244–272.
- Denny-Brown D, Meyer I, Horenstein S. The significance of perceptual rivalry resulting from parietal lesion. *Brain.* 1952;75:433–471.
- Critchley M. *The Parietal Lobes.* 1950.
- Bisiach E, Bulgarelli C, Sterzi R, et al. Line bisection and cognitive plasticity of unilateral neglect of space. *Brain Cogn.* 1983;2:32–38.
- Kinsbourne M. Hemi-neglect and hemisphere rivalry. *Adv Neurol.* 1977;18:41–49.
- Heilman KM, Van Den Abell T. Right hemisphere dominance for attention: the mechanism underlying hemispheric asymmetries of inattention (neglect). *Neurology.* 1980;30:327–330.
- Chatterjee A. Cross-over, completion and confabulation in unilateral spatial neglect. *Brain.* 1995;118(pt 2):455–465.
- Anderson B. A mathematical model of line bisection behaviour in neglect. *Brain.* 1996;119(pt 3):841–850.
- Samuelsson H, Hjelmquist EK, Jensen C, et al. Nonlateralized attentional deficits: an important component behind persisting visuospatial neglect? *J Clin Exp Neuropsychol.* 1998;20:73–88.
- Corwin JV, Kanter S, Watson RT, et al. Apomorphine has a therapeutic effect on neglect produced by unilateral dorsomedial prefrontal cortex lesions in rats. *Exp Neurol.* 1968;94:683–698.
- Larisch R, Meyer W, Klimke A, et al. Left-right asymmetry of striatal dopamine D2 receptors. *Nucl Med Commun.* 1998;9:781–787.
- Corbetta M, Kincade MJ, Lewis C, et al. Neural basis and recovery of spatial attention deficits in spatial neglect. *Nat Neurosci.* 2005;8:1603–1610.
- Heilman KM, Valenstein E. Frontal lobe neglect in man. *Neurology.* 1972;22:660–664.
- Coslett HB, Bowers D, Heilman KM. Reduction in cerebral activation after right hemisphere stroke. *Neurology.* 1987;37:957–962.
- Antonucci G, Guariglia C, Judica A, et al. Effectiveness of neglect rehabilitation in a randomized group study. *J Clin Exp Neuropsychol.* 1995;17:383–389.
- Fanthome Y, Lincoln NB, Drummond A, et al. The treatment of visual neglect using feedback of eye movements: a pilot study. *Disabil Rehabil.* 1995;17:413–417.
- Robertson IH, Gray JM, Pentland B, et al. Microcomputer-based rehabilitation for unilateral left visual neglect: a randomized controlled trial. *Arch Phys Med Rehabil.* 1990;71:663–668.
- Weinberg J, Diller L, Gordon WA, et al. Visual scanning training effect on reading-related tasks in acquired right brain damage. *Arch Phys Med Rehabil.* 1977;58:479–486.
- Weinberg J, Diller L, Gordon WA, et al. Training sensory awareness and spatial organization in people with right brain damage. *Arch Phys Med Rehabil.* 1979;60:491–496.
- Wiart L, Côme AB, Debelleix X, et al. Unilateral neglect syndrome rehabilitation by trunk rotation and scanning training. *Arch Phys Med Rehabil.* 1997;78:424–429.
- Young GC, Collins D, Hren M. Effect of pairing scanning training with block design training in the remediation of perceptual problems in left hemiplegics. *J Clin Neuropsychol.* 1983;5:201–212.
- Gordon WA, Hibbard MR, Egelko S, et al. Perceptual remediation in patients with right brain damage: a comprehensive program. *Arch Phys Med Rehabil.* 1985;66:353–359.
- Robertson IH, Tegner R, Tham K, et al. Sustained attention training for unilateral neglect: theoretical and rehabilitation implications. *J Clin Exp Neuropsychol.* 1995;17:416–430.
- Wilson FC, Manly T, Coyle D, et al. The effect of contralesional limb activation training and sustained attention training for self-care programmes in unilateral spatial neglect. *Restor Neurol Neurosci.* 2000;16:1–4.
- Smania N, Bazoli F, Piva D, et al. Visuomotor imagery and rehabilitation of neglect. *Arch Phys Med Rehabil.* 1997;78:430–436.
- Niemeier JP, Cifu DX, Kishore R. The lighthouse strategy: improving the functional status of patients with unilateral neglect after stroke and brain injury using a visual imagery intervention. *Top Stroke Rehabil.* 2001;8:10–18.
- Niemeier JP. The Lighthouse Strategy: use of a visual imagery technique to treat visual inattention in stroke patients. *Brain Inj.* 1998;12:399–406.
- Rossi PW, Kheyfets S, Reding MJ. Fresnel prisms improve visual perception in stroke patients with homonymous hemianopia or unilateral visual neglect. *Neurology.* 1990;40:1597–1599.
- Vuilleumier P, Valenza N, Mayer E, et al. To see better to the left when looking more to the right: effects of gaze direction and frames of spatial coordinates in unilateral neglect. *J Int Neuropsychol Soc.* 1999;5:75–82.

54. Schindler I, Kerkhoff G. Head and trunk orientation modulate visual neglect. *Neuroreport*. 1997;8:2681–2685.
55. Barrett AM, Burkholder S. Monocular patching in subjects with right-hemisphere stroke affects perceptual-attentional bias. *J Rehabil Res Dev*. 2006;43:337–346.
56. Butter CM, Kirsch N. Combined and separate effects of eye patching and visual stimulation on unilateral neglect following stroke. *Arch Phys Med Rehabil*. 1992;73:1133–1139.
57. Beis JM, Andre JM, Baumgarten A, et al. Eye patching in unilateral spatial neglect: efficacy of two methods. *Arch Phys Med Rehabil*. 1999;80:71–76.
58. Eskes GA, Butler B, McDonald A, et al. Limb activation effects in hemispatial neglect. *Arch Phys Med Rehabil*. 2003;84:323–328.
59. Gainotti G, Perri R, Cappa A. Left hand movements and right hemisphere activation in unilateral spatial neglect: a test of the interhemispheric imbalance hypothesis. *Neuropsychologia*. 2002;40:1350–1355.
60. Robertson IH, North N. Active and passive activation of left limbs: influence on visual and sensory neglect. *Neuropsychologia*. 1993;31:293–300.
61. Bottini G, Karnath HO, Vallar G, et al. Cerebral representations for egocentric space: Functional-anatomical evidence from caloric vestibular stimulation and neck vibration. *Brain*. 2001;124:1182–1196.
62. Johannsen L, Ackermann H, Karnath HO. Lasting amelioration of spatial neglect by treatment with neck muscle vibration even without concurrent training. *J Rehabil Med*. 2003;35:249–253.
63. Karnath HO. Transcutaneous electrical stimulation and vibration of neck muscles in neglect. *Exp Brain Res*. 1995;105:321–324.
64. Schindler I, Kerkhoff G, Karnath HO, et al. Neck muscle vibration induces lasting recovery in spatial neglect. *J Neurol Neurosurg Psychiatry*. 2002;73:412–419.
65. Kerkhoff G, Keller I, Ritter V, et al. Repetitive optokinetic stimulation induces lasting recovery from visual neglect. *Restor Neurol Neurosci*. 2006;24:357–369.
66. Pizzamiglio L, Fasotti L, Jehkonen M, et al. The use of optokinetic stimulation in rehabilitation of the hemineglect disorder. *Cortex*. 2004;40:441–450.
67. Serino A, Bonifazi S, Pierfederici L, et al. Neglect treatment by prism adaptation: what recovers and for how long. *Neuropsychol Rehabil*. 2007;17:657–687.
68. Striemer C, Danckert J. Prism adaptation reduces the disengage deficit in right brain damage patients. *Neuroreport*. 2007;18:99–103.
69. Pisella L, Rode G, Farne A, et al. Prism adaptation in the rehabilitation of patients with visuo-spatial cognitive disorders. *Curr Opin Neurol*. 2006;19:534–542.
70. Luaute J, Michel C, Rode G, et al. Functional anatomy of the therapeutic effects of prism adaptation on left neglect. *Neurology*. 2006;66:1859–1867.
71. Sarri M, Kalra L, Greenwood R, et al. Prism adaptation changes perceptual awareness for chimeric visual objects but not for chimeric faces in spatial neglect after right-hemisphere stroke. *Neurocase*. 2006;12:127–135.
72. Morris AP, Kritikos A, Berberovic N, et al. Prism adaptation and spatial attention: a study of visual search in normals and patients with unilateral neglect. *Cortex*. 2004;40:703–721.
73. Girardi M, McIntosh RD, Michel C, et al. Sensorimotor effects on central space representation: prism adaptation influences haptic and visual representations in normal subjects. *Neuropsychologia*. 2004;42:1477–1487.
74. Frassinetti F, Angeli V, Meneghello F, et al. Long-lasting amelioration of visuospatial neglect by prism adaptation. *Brain*. 2002;125:608–623.
75. Tilikete C, Rode G, Rossetti Y, et al. Prism adaptation to rightward optical deviation improves postural imbalance in left-hemiparetic patients. *Curr Biol*. 2001;11:524–528.
76. Rossetti Y, Rode G, Pisella L, et al. Prism adaptation to a rightward optical deviation rehabilitates left hemispatial neglect. *Nature*. 1998;395:166–169.
77. Fleet WS, Valenstein E, Watson RT, et al. Dopamine agonist therapy for neglect in humans. *Neurology*. 1987;37:1765–1770.
78. Barrett AM, Crucian GP, Schwartz RL, et al. Adverse effect of dopamine agonist therapy in a patient with motor-intentional neglect. *Arch Phys Med Rehabil*. 1999;80:600–603.
79. Geminiani G, Bottini G, Sterzi R. Dopaminergic stimulation in unilateral neglect. *J Neurol Neurosurg Psychiatry*. 1998;65:344–347.
80. Hurford P, Stringer AY, Jann B. Neuropharmacologic treatment of hemineglect: a case report comparing bromocriptine and methylphenidate. *Arch Phys Med Rehabil*. 1998;79:346–349.
81. Oliveri M, Bisiach E, Brighina F, et al. rTMS of the unaffected hemisphere transiently reduces contralesional visuospatial hemineglect. *Neurology*. 2001;57:1338–1340.
82. Brighina F, Bisiach E, Oliveri M, et al. 1 Hz repetitive transcranial magnetic stimulation of the unaffected hemisphere ameliorates contralesional visuospatial neglect in humans. *Neurosci Lett*. 2003;336:131–133.
83. Shindo K, Sugiyama K, Huabao L, et al. Long-term effect of low-frequency repetitive transcranial magnetic stimulation over the unaffected posterior parietal cortex in patients with unilateral spatial neglect. *J Rehabil Med*. 2006;38:65–67.
84. Grujic Z, Mapstone M, Gitelman DR, et al. Dopamine agonists reorient visual exploration away from the neglected hemispace. *Neurology*. 1998;51:1395–1398.
85. Robertson IH, Mattingley JB, Rorden C, et al. Phasic alerting of neglect patients overcomes their spatial deficit in visual awareness. *Nature*. 1998;395:169–172.
86. Thimm M, Fink GR, Kust J, et al. Impact of alertness training on spatial neglect: a behavioral and fMRI study. *Neuropsychologia*. 2006;44:1230–1246.
87. Robertson IH. Cognitive rehabilitation: attention and neglect. *Trends Cogn Sci*. 1999;3:385–393.
88. Karnath HO, Christ K, Hartje W. Decrease of contralateral neglect by neck muscle vibration and spatial orientation of trunk midline. *Brain*. 1993;116(pt 2):383–396.
89. Guariglia C, Coriale G, Cosentino T, et al. TENS modulates spatial reorientation in neglect patients. *Neuroreport*. 2000;11:1945–1948.
90. Na DL, Son Y, Kim CH, et al. Effect of background motion on line bisection performance in normal subjects. *Cortex*. 2002;38:787–796.
91. Pizzamiglio L, Frasca R, Guariglia C, et al. Effect of optokinetic stimulation in patients with visual neglect. *Cortex*. 1990;26:535–540.
92. Bisiach E, Pizzamiglio L, Nico D, et al. Beyond unilateral neglect. *Brain*. 1996;119(pt 3):851–857.
93. Rubens AB. Caloric stimulation and unilateral visual neglect. *Neurology*. 1985;35:1019–1024.
94. Rode G, Perenin MT, Honore J, et al. Improvement of the motor deficit of neglect patients through vestibular stimulation: evidence for a motor neglect component. *Cortex*. 1998;34:253–261.
95. Karnath HO, Fetter M, Dichgans J. Ocular exploration of space as a function of neck proprioceptive and vestibular input—observations in normal subjects and patients with spatial neglect after parietal lesions. *Exp Brain Res*. 1996;109:333–342.
96. Hubel DH, LeVay S, Wiesel TN. Mode of termination of retinotectal fibers in macaque monkey: an autoradiographic study. *Brain Res*. 1975;96:25–40.
97. Farne A, Rossetti Y, Toniolo S, et al. Ameliorating neglect with prism adaptation: visuo-manual and visuo-verbal measures. *Neuropsychologia*. 2002;40:718–729.
98. Ferber S, Danckert J, Joannisse M, et al. Eye movements tell only half the story. *Neurology*. 2003;60:1826–1829.
99. Luaute J, Halligan P, Rode G, et al. Visuo-spatial neglect: a systematic review of current interventions and their effectiveness. *Neurosci Biobehav Rev*. 2006;30:961–982.
100. Bowen A, Lincoln NB, Dewey M. Cognitive rehabilitation for spatial neglect following stroke. *Cochrane Database Syst Rev*. 2002;CD003586.
101. Pierce SR, Buxbaum LJ. Treatments of unilateral neglect: a review. *Arch Phys Med Rehabil*. 2002;83:256–268.
102. Wilson B, Cockburn J, Halligan P. Development of a behavioral test of visuospatial neglect. *Arch Phys Med Rehabil*. 1987;68:98–102.
103. Mahoney FI, Barthel DW. Functional evaluation: the Barthel Index. *Md State Med J*. 1965;14:61–65.
104. Keith RA, Granger CV, Hamilton BB, et al. The functional independence measure: a new tool for rehabilitation. *Adv Clin Rehabil*. 1987;1:6–18.
105. Paolucci S, Antonucci G, Guariglia C, et al. Facilitatory effect of neglect rehabilitation on the recovery of left hemiplegic stroke patients: a cross-over study. *J Neurol*. 1996;243:308–314.
106. Pizzamiglio L, Antonucci G, Judica A, et al. Cognitive rehabilitation of the hemineglect disorder in chronic patients with unilateral right brain damage. *J Clin Exp Neuropsychol*. 1992;14:901–923.
107. Kalra L, Perez I, Gupta S, et al. The influence of visual neglect on stroke rehabilitation. *Stroke*. 1997;28:1386–1391.
108. Worthington A. Cueing strategies in neglect dyslexia. *Neuropsychol Rehabil*. 1996;6:1–17.
109. Robertson I, Hogg K, MacMillan TM. Rehabilitation of unilateral neglect:

- improving function by contralesional limb activation. *Neuropsychol Rehabil.* 1998;8:19–29.
110. Samuel C, Louis-Dreyfus A, Kaschel R, et al. Rehabilitation of very severe unilateral neglect by visuo-spatio-motor cueing; two single case studies. *Neuropsychol Rehabil.* 2000;10:385–399.
 111. Zoccolotti P, Guariglia C, Pizzamiglio L, et al. Good recovery in visual scanning in a patient with persistent anosognosia. *Int J Neurosci.* 1992;63:93–104.
 112. Soderback I, Bengtsson I, Ginsburg E, et al. Video feedback in occupational therapy: its effects in patients with neglect syndrome. *Arch Phys Med Rehabil.* 1992;73:1140–1146.
 113. Tham K, Ginsburg E, Fisher AG, et al. Training to improve awareness of disabilities in clients with unilateral neglect. *Am J Occup Ther.* 2001;55:46–54.
 114. Arene NU, Hillis AE. Rehabilitation of unilateral spatial neglect and neuroimaging. *Eura Medicophys.* 2007;43:255–269.