

Diagnosis and management of functional neurological disorder

Selma Aybek,¹ David L Perez²



¹Neurology Department, Psychosomatic Medicine Unit, Inselspital University Hospital, Bern, and Bern University, Bern, Switzerland

²Divisions of Cognitive Behavioral Neurology and Neuropsychiatry, Functional Neurological Disorder Unit, Departments of Neurology and Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

Correspondence to: S Aybek selma.aybek@med.unibe.ch

Cite this as: *BMJ* 2022;376:o64 <http://dx.doi.org/10.1136/bmj.o64>

Series explanation: State of the Art Reviews are commissioned on the basis of their relevance to academics and specialists in the US and internationally. For this reason they are written predominantly by US authors.

ABSTRACT

Functional neurological disorder (FND), previously regarded as a diagnosis of exclusion, is now a rule-in diagnosis with available treatments. This represents a major step toward destigmatizing the disorder, which was often doubted and deemed untreatable. FND is prevalent, generally affecting young and middle aged adults, and can cause severe disability in some individuals. An early diagnosis, with subsequent access to evidence based rehabilitative and/or psychological treatments, can promote recovery—albeit not all patients respond to currently available treatments. This review presents the latest advances in the use of validated rule-in examination signs to guide diagnosis, and the range of therapeutic approaches available to care for patients with FND. The article focuses on the two most frequently identified subtypes of FND: motor (weakness and/or movement disorders) and seizure type symptoms. Twenty two studies on motor and 27 studies on seizure type symptoms report high specificities of clinical signs (64-100%), and individual signs are reviewed. Rehabilitative interventions (physical and occupational therapy) are treatments of choice for functional motor symptoms, while psychotherapy is an emerging evidence based treatment across FND subtypes. The literature to date highlights heterogeneity in responses to treatment, underscoring that more research is needed to individualize treatments and develop novel interventions.

Introduction

Historical background

Functional neurological disorder (FND) is a prevalent, costly, and potentially disabling condition encountered by healthcare professionals in medical, clinical neuroscience, and rehabilitative specialties.^{1 2} The condition has a complex narrative in the literature that has benefited from and been hampered by the interwoven history of neurology and psychiatry.³ Labeled medicine's "silent epidemic," a "crisis" in neurology, and psychiatry's "blind spot,"⁴⁻⁶ FND has inspired renewed clinical and research interest during the past several decades. Important breakthroughs have included new diagnostic and therapeutic approaches to FND, as well as parallel advances in its pathophysiology. Figure 1 gives a brief overview of the emerging neurobiology of FND^{7 8}—depicting the condition as characterized by dysfunction within and across several brain networks.

Nosological classification

FND is classified as "conversion disorder/functional neurological symptom disorder" in the chapter "Somatic Symptom and Related disorders, code

F44.X" in the Diagnostic and Statistical Manual of Mental Disorders 5th Edition (DSM-5). In the ICD-11 (International Classification of Diseases), FND is classified as "dissociative neurological symptom disorder" in the chapter "Mental, Behavioural or Neurodevelopmental Disorders, code 6B60.X," as well as in the chapter "Diseases of the Nervous System, code 8A0X" under the term "movement disorder for parkinsonism, dystonia, and tremor" (see table 1 for details). This variability within and across classification systems is problematic, as it perpetuates a cartesian dualism and creates coding problems between mental health and neurological disorders that affect which clinical services will be reimbursed, or by which expert patients should be evaluated in medico-legal cases.

Aims of this review

In 2013, a new set of diagnostic criteria for FND appeared in the DSM-5, and emphasized the importance of making a rule-in positive diagnosis based on physical examination and semiological features. In the DSM-IV, emphasis was given to making an exclusionary diagnosis (based on all available neurological tests being normal) and




Fig 1 | Emerging pathophysiology of functional neurological disorder In the top panel, core constructs implicated in FND are highlighted, including disturbances in attention, self-agency, prediction/inference, and emotion/threat processing. In the bottom panel, the brain circuits implicated in the pathophysiology of FND (and their interactions) are displayed. As depicted, FND is a multi-network disorder involving abnormalities within and across brain circuits implicated in sense of agency, emotion/threat processing, attention, homeostatic balance, interoception, multimodal integration, and cognitive/motor control, among other functions. Circuits are described by their related dysfunction in the pathophysiology of FND. Several areas cut across multiple networks; for example, the dorsal anterior insula is most strongly interconnected with the dorsal anterior cingulate cortex, while the posterior insula receives afferent projections from the lamina I spinothalamic pathway and somatosensory cortices. Similarly, the amygdala is part of both the saliency and limbic networks. Prefrontal brain regions are interconnected with striatal-thalamic areas (not shown), and these pathways should also be factored into the neural circuitry of FND. AMY=amygdala; dIPFC=dorsolateral prefrontal cortex; FEF=frontal eye fields; HYP=hypothalamus; PAG=periaqueductal gray; pgACC=perigenual anterior cingulate cortex; sgACC=subgenual anterior cingulate cortex; SMA=supplementary motor area; TPJ=temporoparietal junction. Figures reproduced with permission from Drane et al 2020 *CNS Spectrums*

Table 1 | Nosological classification of functional neurological disorder (diagnostic entities include DSM-5, ICD-10, and ICD-11*)

DSM-5 (2013)	ICD-10 (2016)	ICD-11 (2019)
Somatic symptom and related disorders	Anxiety, dissociative, stress related, somatoform, and other non-psychotic mental disorders	06 mental, behavioral, or neurodevelopmental disorders
F44.X Conversion disorder (functional neurological symptom disorder)	Dissociative (conversion) disorders	6B60 dissociative neurological symptom disorder
F44.4: with motor symptoms (weakness or paralysis, abnormal movement, swallowing, speech)	F44.0 Dissociative amnesia	6B60.0 visual
F44.5: with attacks or seizures	F44.2 Dissociative stupor	6B60.1 auditory
F44.6: with sensory symptoms (anesthesia, sensory loss, visual/olfactory/hearing disturbance)	F44.4 Dissociative motor disorders	6B60.2 vertigo or dizziness
F44.7: with mixed symptoms	F44.5 Dissociative convulsions	6B60.3 sensory
Specify if: acute (<6 months) or persistent (>6 months)	F44.6 Dissociative anesthesia and sensory loss	6B60.4 non-epileptic seizures
Specify if: with or without psychological stressor	F44.7 Mixed dissociative (conversion) disorder	6B60.5 speech
	F44.8 Other dissociative (conversion) disorders	6B60.6 paresis or weakness
	F44.9 Dissociative (conversion) disorder, unspecified	6B60.7 gait
		6B60.8 movement
		6B60.9 cognitive
		08 Diseases of the nervous system
		8A00.3 Functional parkinsonism
		8A02.3 Functional dystonia or spasms
		8A04.4 Functional tremor

*DSM-5=Diagnostic and Statistical Manual of Mental Disorders 5th edition, published 2013. ICD =International Classification of Diseases (ICD-10 is the 10th edition, 2016, and the ICD-11 the 11th, 2019)

linking symptom onset to a psychological trigger. The presence of a psychosocial stressor is now recorded as being present or absent as an adjunctive specifier, after a rule-in diagnosis of FND has been first confirmed. This implies that neurologists (and neuropsychiatrists) are first line in the diagnostic evaluation and immediate clinical management following diagnosis. Yet this is not implemented in clinical practice worldwide, as shown by a vote at the American Academy of Neurology during a 2018 plenary session: a majority of international neurologists expressed the opinion that it is not their role to be primarily involved in the management of FND.⁹ Moreover, many neurologists still order additional tests, even if they are already convinced that the symptom is not caused by another condition¹⁰—reflecting that they do not rely on the clinical diagnosis as stated in the DSM-5.

In this review, we highlight literature that shows multidisciplinary and interdisciplinary approaches are important in the care of patients with FND.¹¹ In fact, international efforts are promoting such multidisciplinary collaborations: the American Neuropsychiatric Association Committee on Research has recently established practice recommendations^{12–13} to guide the diagnostic process, integrating both neurological and psychiatric perspectives—an approach that not only informs diagnosis but also aids the development of a biopsychosocially informed, patient centered treatment plan. A new society, the international Functional Neurological Society (www.fndsociety.org) was established in 2019, open not only to neurologists, psychiatrists, and psychologists but also to all allied healthcare professionals (eg, physiotherapists, occupational therapists, speech and language pathologists, social workers, etc). Several patient associations (fndhope.org, fndaction.org.uk) promote understanding and awareness of the disorder, develop support groups, and assist with access to appropriate medical care. Specialized FND clinics, focusing on diagnostic, therapeutic, and research aspects are being developed worldwide,^{14–16}

modeled in part on a pioneering stepped care model developed in Scotland,¹⁷ and with collaboration at its core.¹⁸ Treatment facilities are also being developed and optimized across outpatient and inpatient settings.^{14–19–22}

This clinical interest occurs in parallel with new research findings that show efficacy of specific therapeutic approaches,^{21–23–24} and elucidate underlying neurobiological mechanisms^{25–26} that open new routes into targeted treatment strategies.^{27–30}

This article presents the evidence available to guide clinicians in the diagnosis and clinical management of patients with FND. We focus on motor (F44.4 and 6B60.3/6B60.5–8) and seizure type (F44.5 and 6B60.4) clinical presentations. We do not include the less frequent subtypes such as individuals with isolated somatosensory deficits (F44.6), cognitive symptoms (6B60.9), special sensory symptoms (visual 6B60.0, auditory 6B60.1), or dizziness and vertigo (6B60.2) that have been reviewed elsewhere.^{31–32}

Incidence and prevalence of FND

FND is a frequent³³ and disabling^{34–35} condition affecting young people,³⁶ and has a poor prognosis in many patients.³⁷ The incidence rate of mixed FND is estimated at 4–12/100 000 population per year.^{38–41} Motor FND (abnormal movements and weakness) is estimated at 4–5/100 000 per year, and seizure type FND (also known as psychogenic non-epileptic or dissociative seizures) at 1.5–4.9/100 000 per year.^{42–44} Prevalence studies are scarce, but the reported rate of FND is around 50/100 000 in the population.³⁶ The prevalence of seizure type FND is estimated at 2–33/100 000.^{36–44}

Sources and selection criteria

We searched the Medline, PsycInfo, and Cochrane databases from inception to 1 November 2020 to find articles pertaining to motor FND (weakness and abnormal movements subtypes) and seizure type FND. Our search terms included “functional”,

Table 2 | Validated positive motor signs for FND (positive signs evaluated in one or more validation studies with a control group, as well as specificities and sensitivities, and with relevant reference from articles reporting on these signs)

	Positive sign	How to test	Evidence/populations tested	Reference	Note
General signs					
	Distractibility	Engage the patient in another motor or cognitive task and observe changes in the abnormal movement	In 19 functional stereotypies v 64 tardive dyskinesia; specificity 100%, sensitivity 58%	Baizabal-Carvalho 2017 ⁴⁵	Validated in stereotypies but can be seen in all FND subtypes
	Variability	Observe changes during history taking/examination/arriving or leaving the examination room: periods of unexplained improvement/disappearance of symptom	In 19 functional stereotypies v 64 tardive dyskinesia; specificity 100%, sensitivity 84%	Baizabal-Carvalho 2017 ⁴⁵	Validated in stereotypies but can be seen in all FND subtypes
	Convergence spasm	Instruct the patient to focus on your finger, 10 cm away from the face, at either extreme lateral gaze for 5 s. Move toward midline and observe the appearance of disconjugate gaze AND miosis	In 13 functional movement disorder, 11 “organic” controls, and 12 healthy: specificity 87%, sensitivity 15%. Good inter-rater reliability (κ 0.6)	Fekete 2012 ⁴⁶	
	Eye movement abnormalities during examination	Systematically test for eye movements even in patients with no symptoms: a discordance between no visual complaint and abnormal findings can be seen (excessive blinking, effortful facial expression, increased latency, gaze deviation, limited range, absent frontalis contraction during upgaze)	In 101 FND patients 43% have abnormal examination	Teodoro 2019 ⁴⁷	No “organic” control group
	Expressive behavior	Look for “expressive” behavior displaying disproportionate effort to the task during examination	In 20 FND v 20 “organic” controls, specificity 95%, sensitivity 55%. Good inter-rater reliability (κ 0.5)	Daum 2015 ⁴⁸	Can be observed throughout the examination
Gait					
	Monoplegic leg dragging	The weak leg is “dragged” like a piece of wood/inanimate object, without spastic circumduction, usually along the floor surface	Validated in 2 studies; pooled specificity 100%, sensitivity 9%. Inter-rater reliability validated in 2 studies: Moderate to good inter-rater reliability (κ 0.4 to 0.7)	Daum 2015, ⁴⁸ Stone 2010 ⁴⁹	
	“Huffing and puffing” sign	Look for 6 behaviors; huffing, grunting, grimacing, breath holding, heavy breathing, crying. Rate each on severity 0 to 4 and duration 0 to 4	In 131 FND v 37 “organic” controls, when score ≥ 2 : specificity 100%, sensitivity 44%. Moderate inter-rater reliability (κ 0.4)	Laub 2015 ⁵⁰	
	Falls toward support	The patient tends to fall in the direction of support (wall, furniture)	In 20 FND v 20 “organic” controls; specificity 93%, sensitivity 19%. Excellent inter-rater reliability (κ 0.8)	Daum 2015 ⁴⁸	
	Excessive slowness	Look for disproportionate slowness in gait (slow stepping movements contrasting with lack of limb bradykinesia)	In 20 FND v 20 “organic” controls; specificity 94%, sensitivity 32%. Moderate inter-rater reliability (κ 0.5)	Daum 2015 ⁴⁸	
	Hesitation/caution	Look for disproportionate hesitation and caution in gait (contrasting with good balance, strength, sensation)	In 20 FND v. 20 “organic” controls; specificity 100%, sensitivity 37%. Good inter-rater reliability (κ 0.7)	Daum 2015 ⁴⁸	
	Non-economic posture	Look for postures during gait that require good balance and strength such as flexed knees	In 20 FND v 20 “organic” controls; specificity 100%, sensitivity 21%. Moderate inter-rater reliability (κ 0.5)	Daum 2015 ⁴⁸	
	Sudden knee buckling	Look for sudden buckling of the knee, usually with each step. Extreme cases will show knee touching the floor at each step	In 20 FND v 20 “organic” controls; specificity 95%, sensitivity 21%. Moderate inter-rater reliability (κ 0.5)	Daum 2015 ⁴⁸	
	Chair test	In case of severe gait disorder, ask the patient to propel a chair with wheels; movements of the legs will be better than during gait	In 9 FND v 9 “organic” controls; specificity 100%, sensitivity 89%	Okun 2007 ⁵¹	
Axial					
Face					
	Hemifacial spasm	Lack of “other Babinski sign”	Observe eyebrow elevation: in “organic” cases ipsilateral to the spasm (= “other Babinski sign”), in FND absent or contralateral to the spasm	15 functional movement disorder v 37 hemifacial spasms: specificity 70%, sensitivity 100%	Baizabal-Carvalho 2017 ⁵²
	Long tonic contraction	Observe if tonic contractions are long (>3 s)	Tonic contraction >3 s: specificity 97%, sensitivity 87%	Baizabal-Carvalho 2017 ⁵²	
	Bilateral hemispasm	Observe if purely unilateral (“organic”) or also bilateral/alternating (functional)	Bilateral spasm: specificity 97%, sensitivity 40%	Baizabal-Carvalho 2017 ⁵²	
	Isolated lower face spasm	Observe if isolated lower face involvement	Isolated lower spasm: specificity 97%, sensitivity 33%	Baizabal-Carvalho 2017 ⁵²	
	Lip pulling	Observe if lower face involvement is a tonic deviation of the lower lip, often with ipsilateral platysma contraction	Lip pulling: specificity 100%, sensitivity 47%	Baizabal-Carvalho 2017 ⁵²	
	Oro-lingual dyskinesia	Lack of chewing movements	Observe if abnormal movement includes chewing	In 9 oro-lingual functional movement disorder v 50 oro-lingual tardive dyskinesia: lack of chewing movement: specificity 82%, sensitivity 78%	Baizabal-Carvalho 2017 ⁴⁵
		Lack of self biting	Observe if abnormal movement includes self-biting	Lack self-biting: specificity 64%, sensitivity 100%	Baizabal-Carvalho 2017 ⁴⁵
		Purely lingual movements	Observe if abnormal movement are both lingual and oral (“organic”) or purely lingual (functional)	Lingual without mouth movement: specificity 96%, sensitivity 44%	Baizabal-Carvalho 2017 ⁴⁵
		Oro-lingual and speech problem	Observe if speech is also affected	Accompanying abnormal speech: specificity 90%, sensitivity 44%	Baizabal-Carvalho 2017 ⁴⁵

(Continued)

Table 2 | Continued

	Positive sign	How to test	Evidence/populations tested	Reference	Note
Neck/trunk					
Weakness	Sterno-cleido-mastoid	Rotation of the head is weak (usually when turning toward the side of the limb weakness)	Validated in 2 studies; pooled specificity 93%, sensitivity 53%. Excellent inter-rater reliability (κ 0.83)	Daum 2015, ⁴⁸ Horn 2017 ⁵³	
Lack of balance	Functional Romberg	Large disbalance display during Romberg task but no falls, usually gets better with distraction (drawing number in the back, cognitive task)	In 20 FND v 20 "organic" controls: specificity 100%, sensitivity 39% Moderate inter-rater reliability (κ 0.5)	Daum 2015 ⁴⁸	
Cataplexy	Absence of hypotonic facial phenomenon	Observe face during episode: absence of ptosis, mouth opening, and tongue protrusion in functional cataplexy	In 21 FND v 30 narcolepsy patients. Good to excellent inter-rater reliability (κ 0.74 to 0.86)	Pizza 2018 ⁵⁴	
	No abrupt facial change	Observe if smile and facial expression changes	Good inter-rater reliability (κ 0.63 to 0.78)	Pizza 2018 ⁵⁴	
	No facial jerks/grimaces	Observe if brief abnormal movement occurs	Good inter-rater reliability (κ 0.67 to 0.73)	Pizza 2018 ⁵⁴	
	No postural dyscontrol	Observe if head drops and trunk falls	Excellent inter-rater reliability (κ 0.83 to 0.86)	Pizza 2018 ⁵⁴	
	Persistence of reflexes	Test tendon reflexes during the episode; in cataplexy they disappear but in functional cases they persist	Good inter-rater reliability (κ 0.77)	Pizza 2018 ⁵⁴	
Limb					
Arm/leg weakness	Discordance/inconsistency	A movement cannot be done but the same muscle can be used later in another movement	In 15 FND v 40 controls: specificity 98%, sensitivity 13%	Chabrol ⁵⁵	
Arm/leg weakness	Give way weakness	When testing strength against resistance; initially good and then sudden loss of resistance from the patient	Validated in 3 studies: pooled specificity 97%, sensitivity 67% Good inter-rater reliability (κ 0.6)	Daum 2015, ⁴⁸ Chabrol, ⁵⁵ Stone 2010 ⁴⁹	
Arm/leg weakness	Co-contraction	When testing strength, no movement at the joint (elbow, for example) occurs because co-contraction of agonist and antagonist is observed	Validated in 2 studies; pooled specificity 100%, sensitivity 28% Good inter-rater reliability (κ 0.77)	Daum 2015, ⁴⁸ Baker ⁵⁶	
Arm weakness	Drift without pronation	Arms stretched out, palms up in a full supination position, fingers adducted, eyes closed for 10 s: if a downward drift is seen, observe if a movement of pronation also occurs	Validated in 2 studies; pooled specificity 96%, sensitivity 78%. Good inter-rater reliability (κ 0.78)	Daum 2013, ⁵⁷ Daum 2015 ⁴⁸	
Arm weakness	Abductor finger sign	In severe unilateral hand weakness, ask the patient to abduct the fingers of the healthy hand against resistance; observe if involuntary abduction of the 5th finger in the weak hand occurs (functional) or not ("organic")	In 10 FND v 11 "organic" controls; specificity 100% sensitivity 100%	Tinnazzi 2008 ⁵⁸	
Arm weakness	Flex-ext sign	Arms flexed at 30°, forearms held near the wrists by examiner. Ask patient to flex the healthy arm against resistance and observe/feel if increased extension of the weak arm occurs (functional) or not ("organic"). Then ask patient to flex weak arm and observe if extension of healthy arm occurs ("organic") or not (functional)	In 10 FND v 23 "organic" controls: specificity 100% sensitivity 100%	Lombardini ⁵⁹	
Leg weakness (unilateral)	Hoover sign	Ask to flex the healthy hip against resistance and observe/feel the strength of hip extension of the weak leg (if patient lying: examiner's hand under the heel, if sitting under the thigh). Compare with voluntary hip extension of the weak leg: if involuntary strength > voluntary strength, Hoover is positive	Validated in 5 studies: pooled specificity 99.5%, sensitivity 61%	Sonoo 2004, ⁶⁰ Tinnazzi 2008, ⁵⁸ Stone 2010, ⁴⁹ Mcwirthner 2011, ⁶¹ Daum 2015 ⁴⁸	
Leg weakness (unilateral)	Abductor sign	Ask to abduct both legs against resistance: observe/feel if involuntary abduction of the weak leg occurs (functional) or not ("organic")	In 16 FND v 17 "organic" controls; specificity 100%, sensitivity 100%	Sonoo 2004 ⁶⁰	
Leg weakness	Spinal injury center test	Passively put both legs in a flexed position, sole of feet touching the bed during a lying position; observe if the weak leg stays in this position (functional) or falls back on the bed ("organic")	Validated in 2 studies: pooled specificity 96%, sensitivity 47%	Daum 2015, ⁴⁸ Yugue 2004 ⁶²	
Tremor	Distractibility	Pause during ballistic movement or during other motor/mental task or change in amplitude and frequency	In 50 FND tremor v 160 other tremors: specificity 92%, sensitivity 94%	Van der Stowe ⁶³	
	Increase in amplitude with weight	Wrap a weight (500 g) around the wrists and observe change in frequency and amplitude	In 50 FND tremor v 160 other tremors: specificity 92%, sensitivity 22%		
	Entrainment	Ask to imitate tapping motion with one hand and observe the change in tremor frequency on the other	In 50 FND tremor v 160 other tremors: specificity 91%, sensitivity 91%		
	Combination of the 3 features	Presence of at least 2 of these 3 tested in 50 FND and 160 other individuals with tremor	In 50 FND tremor v 160 other tremors; specificity 93%, sensitivity 100%		

Table 3 | Validated positive signs for seizure type FND (positive signs evaluated in one or more validation studies with a control group, as well as specificities and sensitivities, and with relevant reference from articles reporting on these signs)

Positive sign	How to test	Evidence /populations tested	Reference	Note
General signs				
Long duration	Observe if >2 minutes	In 341 FND v 441 epilepsy patients: specificity 93% sensitivity 65%	Seneviratne 2017 ⁶⁴	Other studies: Syed 2011, ⁶⁵ Bazil, ⁶⁶ Jedrzejczak 1999, ⁶⁷ De Paola 2016, ⁶⁸ Slater 1995, ⁶⁹ Azar 2008, ⁷⁰ Vogrig 2018, ⁷¹ Bazil 1997, ⁶⁶ CAVEAT: duration can be short in frontal lobe epilepsy, Saygi 1992. ⁷² Note: status epilepticus should also be considered on the differential diagnosis
Waxing and waning/ fluctuating	Observe seizure course: a decrease/ increase of motor events and/or pauses in the seizure course is suggestive of FND course	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 94%	De Paola 2016 ⁶⁸	Other studies: Syed 2011, ⁶⁵ Vogrig 2018, ⁷¹ Chen 2008 ⁷³
Clues that the patient has preserved awareness				
External influence on attack course	Observe if others can intensify or alleviate the symptoms	In 12 FND v 23 epilepsy patients: specificity 99% sensitivity 83%	Syed 2011 ⁶⁵	
Eye contact	Observe if eyes respond to environment and stimuli	In 12 FND v 23 epilepsy patients: specificity 77% sensitivity 79%	Syed 2011 ⁶⁵	
Responsiveness	Observe if patient can respond during an attack	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 36%	De Paola 2016 ⁶⁸	
Shaking without loss of consciousness	Observe if semiology looks like generalized seizure (shaking of limbs) but with NO loss of consciousness	In 20 FND v 20 epilepsy patients: specificity 100% sensitivity 20%	Devinsky 1996 ⁷⁴	
Recall of event	Ask the patient if she/he recalls the event	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 60%	De Paola 2016 ⁶⁸	
Recall of a named item	During the attack, name a color, object, or phrase and ask the patient afterwards if he/ she can recall it; if yes suggestive of FND	In 20 FND v 20 epilepsy patients: specificity 85% sensitivity 90%	Devinsky 1996 ⁷⁴	Other study: Bell 1998 ⁷⁵
Motor behavior				
Closed eyes	Observe if eyes are closed and/or resist attempted opening by examiner, and or eyelids flutter during seizure	In 52 FND v 156 epilepsy patients: specificity 98% sensitivity 96%	Chung 1996 ⁷⁶	Other studies: Syed 2011, ⁶⁵ Chen 2008, ⁷³ De Paola 2016, ⁶⁸ Gates 1985, ⁷⁷ Devinsky 1996, ⁷⁴ Detoleo 1996, ⁷⁸ Azar 2008 ⁷⁰
Asynchronous limb movements	Observe if arm or leg movements are synchronous: if not, suggestive of FND	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 84%	De Paola 2016 ⁶⁸	Other studies: Syed 2011, ⁶⁵ Chen 2008, ⁷³ Azar 2008, ⁷⁰ Gates 1985, ⁷⁷ CAVEAT: can be seen in frontal lobe seizures
Pelvis thrusting	Observe if rhythmical movements of pelvis occur	In 12 FND v 23 epilepsy patients: specificity 99% sensitivity 8%	Syed 2011 ⁶⁵	Other studies: De Paola 2016, ⁶⁸ Chen 2008, ⁷³ Azar 2008, ⁷⁰ Gates 1985, ⁷⁷ Geyer 2000, ⁷⁹ Saygi 1992 ⁷²
Side-to-side head movements	Observe if head (or body) has side-to-side lateral movements	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 66%	De Paola 2016 ⁶⁸	Other studies: Syed 2011, ⁶⁵ Chen 2008, ⁷³ Azar 2008, ⁷⁰ Gates 1985, ⁷⁷ Saygi 1992 ⁷²
Arching back ("Arc de cercle")	Observe if the patient arches back in opisthotonos-like posture	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 38%	De Paola 2011 ⁶⁸	Other study: Syed 2011 ⁶⁵
Rotation in bed	Observe if the patient rotates/changes position in bed	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 26%	De Paola 2016 ⁶⁸	Other study: Azar 2008, ⁷⁰ CAVEAT: can be seen in frontal lobe seizures, Saygi 1992 ⁷²
Accompanying behavior				
Ictal crying	Observe if the patient cries during seizure	In 12 FND v 23 epilepsy patients: specificity 98% sensitivity 8%	Syed 2011 ⁶⁵	Other studies: De Paola 2016, ⁶⁸ Devinsky 1996, ⁷⁴ Chen 2008, ⁷³ Slater 1995, ⁶⁹ Walczak 1996, ⁸⁰ Asadi-Pooya, 2016 ⁸¹
Ictal whispering	If the patient talks, observe if whispering	In 12 FND v 23 epilepsy patients: specificity 91% sensitivity 49%	Syed 2011 ⁶⁵	
Ictal stuttering	If the patient talks, observe if stuttering	In 117 FND v 113 epilepsy patients: specificity 100% sensitivity 9%	Vossler 2004 ⁸²	
Ictal hyperventilation	Observe if patients hyperventilates throughout the seizure	In 50 FND v 20 epilepsy patients: specificity 100% sensitivity 14%	De Paola 2016 ⁶⁸	Other studies: Devinsky 1996 ⁷⁴
Post-ictal behavior				
Short, irregular, shallow breathing pattern	Observe breathing pattern: in FND often irregular, rapid, and shallow (hyperventilation pattern), no snoring, not loud and shorter duration	In 24 FND v 23 generalized epilepsy and 20 frontal lobe epilepsy patients: Mean duration in FND 94 ±55 seconds, in generalized epilepsy 347 ±118 seconds and in frontal lobe epilepsy 64 ±26 (P<0.001)	Azar 2008 ⁷⁰	Other studies: Chen 2008 ⁷³
Rapid recovery	Observe if patient has immediate recovery	In 12 FND v 23 epilepsy patients: specificity 85% sensitivity 73%	Syed 2011 ⁶⁵	Other studies: Izadyar 2018 ⁸³ CAVEAT: rapid recovery also occurs in frontal lobe epilepsy, Saygi 1992, ⁷² Azar2008 ⁷⁰
No confusion	Observe if patient has confusion or disorientation; if not suggestive of FND	In 24 FND v 23 epilepsy patients (generalized); specificity 100% sensitivity 88%	Azar 2008 ⁷⁰	Slater 1995 ⁶⁹ CAVEAT: no confusion also occurs in frontal lobe epilepsy, Saygi 1992 ⁷²
Abrupt signs of recovery	Observe 3 specific behaviors: 1. Blink or brief head shaking after the event. 2. Looking around, 3. Question "what happened?"	In 64 FND v 42 epilepsy patients: if at least one of the 3 behaviors: specificity 100% sensitivity 45%	Izadyar 2018 ⁸³	

(Continued)

Table 3 | Continued

Positive sign	How to test	Evidence /populations tested	Reference	Note
Additional clues				
Teddy Bear sign	Observe if a stuffed animal is brought by the patient during vEEG monitoring: when yes, suggestive of FND	In 104 FND v 147 epilepsy patients: specificity 88% sensitivity 13%	Cervenka ⁸⁴	Not a semiology sign per se
Signs suggestive of epilepsy				
Occurrence from physiological sleep	During video-EEG monitoring, assess if events occur during physiological sleep	In 280 non-epileptic events v 622 epileptic events: specificity 100% sensitivity 31%	Bazil 1997 ⁶⁶	Other study: De Paola 2016, ⁶⁸ If present, is specific for epilepsy. Requires video EEG, not a bedside test. Apparent clinical sleep but without EEG sleep pattern (pseudosleep) occurs in FND (100% specificity, 56% sensitivity) Benbadis 1996 ⁸⁵
Vocalization during attack	Observe if brief vocalization occurs during the seizure (epileptic cry)	In 25 FND v 25 epilepsy patients: specificity 100% sensitivity 60%	Gates 1985 ⁷⁷	If present is specific for epilepsy. In FND, vocalization such as groaning, moaning can occur before the seizure Other study: De Paola 2016 ⁶⁸
Oral (lateral tongue) laceration	Observe if lateral tongue laceration occurred during the event	In 18 FND vs. 66 epilepsy patients: specificity 100% sensitivity 26%	Oliva 2008 ⁸⁶	If present is specific for epilepsy. Tested in inpatient setting with objective evaluation by staff. Other study: Dufresne 2019. ⁸⁷ Of note, REPORTED tongue laceration (not objectively assessed) has been reported equally frequently in epilepsy and FND (around 25%) ^{88,89}
Self-injury	Observe if event caused injury	In 32 FND v 42 epilepsy patients: specificity 97% sensitivity 19%	Slater 1995 ⁶⁹	If present is specific for epilepsy. Of note, reported injuries are less frequent in FND but can occur (found in up to 40% of cases in Peguero 1995 ⁸⁸)
Post-ictal Babinski sign	Test for plantar reflex within 5 minutes of the event	In 13 FND v 40 patients with epilepsy: specificity 100%, sensitivity 43%	Walczak 1994 ⁹⁰	If Babinski sign, specific for epilepsy

“psychogenic”, “conversion”, “dissociative”, and “hysterical.” More comprehensive details on search terms are available in the supplementary file.

We reviewed titles and abstracts of studies that provided sufficient information on clinical diagnosis and treatment. We excluded case reports, studies not available in English, French, or German, and articles focusing on mechanistic related research questions. In addition, we reviewed the reference lists of selected articles and included other relevant articles. We prioritized original articles but also included noteworthy systematic reviews, narrative reviews, and expert opinions. For the review of positive signs, we summarized in table 2 and table 3 those which had some form of validation (controlled designs to test for specificity and sensitivity or data on inter-rater reliability). To this end, we selected 22 studies^{45-63 91-93} reporting on 37 bedside clinical tests or groups of tests for motor FND (functional weakness and functional movement disorder) as well as 27 studies^{64-87 90 94 95} reporting on 23 bedside clinical tests or group of tests for seizure type FND. In addition, we also discuss important case series or reviews that highlight relevant bedside clinical tests. We excluded studies that had a primary focus on laboratory tests, as our aim was to provide evidence informing the use of clinical neurological examination findings at the bedside.

Diagnosis of FND

Criterion A of the DSM-5 requires “One or more symptoms of altered voluntary motor or sensory function.”⁹⁶ This means that patients present with neurological symptoms, and when these concern motor function (F44.4) they can be divided into two broad categories: negative symptoms (lack of movement, weakness) or positive symptoms

(abnormal movement such as tremor, jerks, dystonia, etc). Symptoms can also occur in brief episodes and resemble an epileptic seizure, encompassing the functional seizure (F44.5) diagnosis. To frame subtypes under the overarching diagnostic category of FND, this article refers to functional seizures as seizure type FND.

Criterion B of the DSM-5 requires that “Clinical findings provide evidence of incompatibility between the symptom and recognized neurological or medical conditions.” Demonstrating this incompatibility (eg, noting that subacute arm or leg weakness is distinct from lesional cortico-spinal tract disruptions as might be present in multiple sclerosis or ischemic stroke) is achieved through evaluating for positive signs during the physical examination. The presence of these signs, such as a give-way pattern of weakness when performing confrontation strength testing, or appreciation of tremor entrainment (ie, the rhythmicity of the tremor can be modulated by performance of paced volitional movements performed in a different body part) confirms criterion B and enables a physician to make a positive rule-in diagnosis. This approach is fundamentally different from an exclusionary process and helps not only to make the diagnosis but also to initiate treatment. Indeed, showing or explaining to the patient how diagnosis was reached—by appreciating positive physical examination signs—has been suggested by expert opinion to help individuals understand their disorder and not feel as though “everything is normal in my tests” so the doctor is jumping to the conclusion that “it is functional.”

Most of these clinical bedside positive signs were described a century ago by Jean-Martin Charcot,⁹⁷ Charles Hoover,⁹⁸ and Joseph Babinski.⁹⁹ In the era of evidence based medicine, recent studies have looked

at the sensitivity and specificity of these signs, and a few tested their inter-rater reliability. Below, we detail the current evidence on the formal validation of these rule-in signs.

Positive diagnosis of motor FND (F44.4)

Our search found 37 bedside clinical tests or groups of tests for motor FND (functional weakness and functional movement disorder) that had some formal validation (table 2). Sample size varied between 8 and 107 patients with FND. Most investigations of positive signs were conducted in a single study or a small number of studies (maximum five for the Hoover sign), allowing for the calculation of pooled specificity and sensitivity by merging data from different studies. Five studies reported on the inter-rater reliability of positive signs.

Overall, the specificities of validated signs are high, ranging from 64% to 100%; however, the sensitivities are lower, ranging from 9% to 100%. Inter-rater reliability of these signs is overall good to excellent (defined as κ values: <0.2 poor/0.21-0.4 fair/0.41-0.6 moderate/0.61-0.8 good/>0.8 excellent).

What to look for at the bedside

General signs common to all FND presentations are: variability of the symptom, which can be observed during history taking and examination, and effortful or grimacing expression while following the examiner's instructions during examination. If suspecting a functional movement disorder, test oculomotor function to show abnormal eye movements and in particular convergence spasm,¹⁰⁰ even if the patient did not endorse this as a concern during history taking.

When assessing gait, look for typical positive signs such as monoplegic leg dragging, excessive visible effort ("huffing and puffing" sign),⁵⁰ falling toward support (chair nearby, table, wall), excessive slowness, hesitation or caution, non-economic posture (for example knee flexed), and knee buckling (sudden loss of tone at each step).¹⁰¹ Asking a patient with severe gait disorder to propel a chair while sitting on it will show improvement in FND.⁵¹

When assessing hemifacial spasm, look for typical signs such as long contraction of more than three seconds, lip pulling (tonic deviation of the lip, often the lower one) sometimes with platysma contraction, and lack of "other Babinski sign" for hemifacial spasm (other Babinski sign=eyebrow elevation on the side of the spasm).¹⁰² Positive signs for functional orofacial movements in comparison with tardive dyskinesia are: lack of chewing movements, lack of self-biting, lingual movements without mouth movements, and abnormal speech.⁴⁵ A large case series (61 patients) that focused on facial functional movement¹⁰³ reported involvement of the lip as the most frequent (60.7%, with the lip pulling feature).

When assessing movements of the trunk, look for the typical positive sign of asymmetry in strength of the sterno-cleido-mastoid muscle.⁵³ A "functional

Romberg" sign is described as large movements of imbalance with sudden steps and no falls and improvement with cognitive distraction or numbers drawn on the back.⁴⁸

When assessing episodes of cataplexy (brief, symmetrical loss of muscle tone with retained consciousness precipitated by strong emotions) look for positive signs⁵⁴ such as lack of sudden facial expression change, facial jerks or grimaces, postural dyscontrol (head drop, trunk fall), in addition to preserved tendon reflexes (which typically disappear during cataplexy associated with narcolepsy).

When assessing upper arm weakness, look for discordance or inconsistency in strength (at different instances during the examination), as well as a give-way/collapsing pattern, drift without pronation, and/or co-contractions of agonist and antagonist muscles preventing movement of the tested joint. As a cautionary note, give-way/collapsing pattern of weakness is common in patients with pain limited weakness (and pain limited weakness should not be mistaken for functional limb weakness).¹⁰⁴ In cases of complete hand plegia, involuntary abduction of the fifth finger can be seen when the patient is asked to do a forced abduction against the examiner's resistance on the healthy hand.⁵⁸ The flex-ext sign, which is the equivalent of the Hoover sign,¹⁰⁵ can be elicited as follows: the involuntary flexion of the arm at the elbow that occurs when the patient focuses on extending the healthy elbow against the examiner's resistance is better than the voluntary flexion.⁵⁹

When assessing lower limb weakness, also look for discordance/inconsistency, give-way/collapsing weakness, co-contractions, and the Hoover sign.⁶¹ The classical way to describe a positive Hoover sign is when the involuntary hip extension (when the patient focuses on flexing the healthy leg against the examiner's resistance) is stronger than the voluntary hip extension. A similar pattern can be found during leg abduction⁶⁰: when the patient is asked to do a forced abduction with both legs against the examiner's resistance, the weak leg will have a stronger involuntary abduction than when the voluntary abduction is tested. In patients with severe unilateral leg weakness, positioning passively the leg in flexion with the soles on the bed (spinal injury test)⁶² shows a discordance in strength as the weak leg will not fall on the side, as expected in complete weakness.

When assessing tremor, typical signs are distractibility, entrainment, and increase in amplitude with weight load on the wrists. In addition, look for variability in amplitude, frequency, and direction of tremor.⁶³ A "whack a mole" sign can be seen¹⁰⁶: when the limb affected by tremor is immobilized by the examiner, the tremor appears in another body segment (head, trunk, other arm, or legs).

No validated clinical signs are available for assessing dystonia, but a pattern of adult sudden onset fixed dystonia (typically clenched fist sparing thumb and index finger¹⁰⁷) or equinovarus foot is

suggestive of functional dystonia.¹⁰⁸ Associated prominent pain and other FND signs can help support the diagnosis.¹⁰⁹

When assessing tics, no validated signs are available but clinical clues can help identify functional tics^{110 111}: lack of premonitory urge and inability to suppress the movement, female preponderance, additional FND symptoms, lack of response to anti-tic medication, and absence of family history. In functional tics, the cranial region is less affected, the type of tic is often “blocking” (ie, interferes with voluntary action) and palin, echo, and copro phenomenon are less common.

Overall, the evidence for rule-in motor signs shows very high specificity, which advocates for their routine use in clinical practice. A range of educational pictorial and video libraries illustrate many of these signs.^{2 101 102 112 113} Too much emphasis on a single sign, however, can lead to false positives. In a cohort of 190 patients diagnosed with a neurological disorder, 37 (20%) had at least one positive functional neurological sign.⁹¹ Interestingly, regression analysis showed that this 20% of the cohort had typical risk factors known in patients with FND, suggesting that the presence of positive signs in this subgroup could either be false positives or indicate the presence of an FND comorbidity. Keep in mind the possibility that the patient has both FND and another neurological disorder: recent reports describe functional neurological signs in a subset of patients with Parkinson’s disease^{114 115} or multiple sclerosis.¹¹⁶ Overall, data from a systematic review and a prospective study underscore that rates of misdiagnosis in FND since 1970 (once confirmed) are low, and between 1% and 4%.^{117 118}

Recently, efforts have been made to integrate additional clinical features in the process of diagnosis, such as, for example, abrupt onset, fluctuations of the motor symptom, comorbid pain, and fatigue.¹¹⁹ The presence of these features should raise the index of suspicion and prompt a more systematic search of signs positive for FND.

Positive diagnosis of seizure type FND (F44.5)

Our search highlighted 23 bedside clinical tests or groups of tests for seizure type FND that had some formal validation (table 3). Sample size varied between 11 and 341 patients with seizure type FND. Most positive signs were investigated in multiple studies with different control groups, so we decided not to calculate pooled specificities and sensitivities. We report in table 3 data from the main study (usually the largest cohort) of the corresponding sign but added citations of all others that tested the same sign. Four studies looked at inter-rater reliability of the signs.^{65 79 94 95}

Overall, the specificities of validated signs are high, ranging from 77% to 100%. Sensitivities are lower, ranging from 9% to 96%. The inter-rater agreement of two independent raters was excellent ($\kappa > 0.87$) for 40 signs.⁶⁵

What to look for at the bedside

Duration is an important feature helping to differentiate seizure type FND from epileptic seizures: a duration of more than two minutes is highly specific for FND (although caution should be taken not to miss status epilepticus in emergent situations). A pattern of waxing and waning, irregular course, or pauses in the event are also typical.

Preserved consciousness can be shown when patients recall the event or are able to encode a memory (it is useful to ask the patient to memorize items during the event and ask afterwards if that information can be recalled). Indirect evidence of preserved awareness includes patients responding to eye contact, exploring the room visually, or having their event influenced by other people (alleviating or aggravating it).

Specific behaviors can be observed, such as closed eyes or even resistance to passive opening attempts by the examiner, asynchronous limb movements (can also occur in frontal lobe seizures), pelvis thrusting, side-to-side head movements, and arching back.

Vocalization or abnormal speech can take the form of moaning or groaning (usually before the event), crying or weeping, whispering, or stuttering. Vocalization in the form of a sudden cry during the seizure is suggestive of epileptic seizures (during the tonic phase).

After the event, rapid recovery is typical in seizure type FND with no post-ictal confusion. Motor signs indicating an abrupt stop of the event can be observed, such as blinking or brief head shaking indicating the end, looking around, asking “what happened?” Breathing is different from epileptic stertorous breathing (low pitch sound during inspiration) and takes the form of regular hyperventilation.

Features that suggest epileptic seizures include: occurrence from physiological sleep and post-ictal Babinski sign. Note that urinary incontinence¹²⁰ and self-injury⁸⁸ can occur in individuals with epileptic seizures and in those with the seizure type of FND. Also, an oral tongue laceration (usually lateral part) documented by medical staff is specific for epilepsy; reports of tongue laceration by the patient are less specific, as it has been reported in up to 21% of non-epileptic events compared with 27% of epileptic seizures.⁸⁹

Overall, the evidence for positive signs of seizure type FND also shows very high specificity, which advocates for their routine use clinically. Along with these clinical signs, electroencephalographic (EEG) recording of the event can help differentiate FND from epilepsy as the event will not be accompanied by the typical electrographic features on EEG. The gold standard for diagnosis is thus to capture a typical event with video EEG (if needed with specific provocation techniques¹²¹). Video EEG is not always available, therefore the International League Against Epilepsy has developed staged criteria to make a diagnosis of seizure type FND, including possible, probable, clinically established, and documented criteria.¹²² A diagnosis is “possible” when the history

and description from a witness is suggestive of FND and an interictal EEG is normal. The diagnosis reaches the level of “probable” when a clinician with experience diagnosing seizure disorders observes the positive signs either by witnessing an event or a video recording and an interictal EEG is normal. Thus, it is helpful to ask patients to obtain home videos of typical events when possible.¹²³

The diagnosis is “clinically established” when these semiological features are observed (in person or on video) and an ambulatory ictal EEG of a typical event is normal (separated in time). It is “documented” when a typical event is concurrently captured on video and EEG. Keep in mind the possibility that a patient has both FND and epilepsy, as around 20% of patients with seizure type FND also have epileptic seizures.^{124 125}

The experience of the clinician assessing the presence and interpretation of these positive signs is also important. A study¹²⁶ showed that emergency department staff (physicians or nurses) without specific neurology training only recognized 44-58% of functional neurological events, while neurology-trained staff recognized 70-73%, and epileptologists 88% of seizure type FND events. These observations underscore the need for increased educational efforts pertaining to the recognition of highly specific signs for seizure type FND.

Recent efforts have been made to integrate self-reported subjective features in the diagnosis process, such as, for example, amnesia of the event, mind-body-world disconnection, or sensory experiences.¹²⁷ The presence of these features can help raise the index of suspicion for a given diagnosis, prompting diagnostic confirmation based on semiological features and EEG data. Emerging research also looks at how to optimally combine self-report and objective signs to support the clinical diagnosis, such as, for example, history of migraine and female sex associated with long duration and eye closure being strongly suggestive of FND.¹²⁸ Before good clinical scales with precise cut-off scores are available, subjective reports should be regarded cautiously, and objective assessment through video EEG remains the gold standard.

Role of exclusion (alternative diagnoses) in making the diagnosis

Criterion C of the DSM-5 requires that “the symptom or deficit is not better explained by another medical or mental (health) disorder.” This should not be mistakenly interpreted to represent that a patient cannot have FND AND a comorbid neurological condition. In instances of diagnostic uncertainty, the diagnosis for a subset of patients with functional movement symptoms can benefit from the use of an electromyogram, an accelerometer, or an EEG (to seek Bereitschaft potential in cases of jerky movements, for example) to help make a laboratory assisted diagnosis of FND (for a review of adjunctive diagnostic tests in FND, see¹²⁹). In particular, validated criteria for tremor (which can capture

features of entrainment or pauses that may not be readily appreciable through visual inspection) are now available to help reach a “laboratory supported” level of certainty about diagnosis.¹³⁰⁻¹³² Similarly, for patients with seizure of diagnostic uncertainty, using adjunctive diagnostic tests,¹³³ such as nuclear medicine brain imaging approaches, may help provide additional diagnostic clarification.¹³

Impact on daily activities and quality of life

Criterion D of the DSM-5 requires that “the symptom or deficit causes clinically significant distress or impairment in social, occupational, or other important areas of functioning or warrants medical evaluation.” In clinical practice, the fact that patients seek medical attention helps support this criterion, as their symptoms often have a significant impact on their daily activities. Across both motor and seizure type FND, health related quality of life (HRQoL) is comparable with, if not worse than, that observed in other major neurological disorders. The most robust literature is found in those with seizure type FND, where this population has consistently reported lower HRQoL than populations with epilepsy.^{134 135} In a 20 year retrospective study, patients with seizure type FND also showed a standardized mortality ratio 2.5 times above that of the general population—rates comparable with people observed in treatment for refractory epilepsy.¹³⁶ Notably, frequency of seizures has not been found to have a dose dependent relationship with HRQoL in patients with seizure type FND; achieving complete resolution of seizures, however, does positively correlate with improvements in HRQoL.^{137 138} As detailed in a systematic review, a range of psychiatric and psychosocial factors influence HRQoL in patient with seizure type FND.¹³⁹ These factors include depression, dissociation, other somatic (bodily) symptoms, escape-avoidance coping, negative illness perceptions, medication side effects, family functioning, and perceived stigma regarding a diagnosis of seizure type FND.^{134 135 140-147}

Patients with motor FND reported similar HRQoL to patients with Parkinson’s disease,^{148 149} and worse HRQoL compared with individuals with primary dystonia.¹⁴⁹ Non-motor symptoms were closely related to HRQoL in patients with motor FND—most notably fatigue, cognitive complaints, and anxiety.^{150 151} A study of 107 patients with motor FND reported them to have similarly impaired HRQoL to patients with other neurological causes for limb weakness.⁴⁹ Two separate long term follow-up studies showed that HRQoL remained impaired in patients with motor FND at levels comparable with patients who had other neurological conditions (eg, multiple sclerosis).^{117 152} In one 14 year follow-up study, patients with motor FND showed a 1.48 standardized mortality ratio.¹¹⁷ Lastly, in a study of mixed FND, positive associations were observed between social network size and physical and mental HRQoL, underscoring the importance of the biopsychosocial formulation in determinants of health status in patients with FND.¹⁵³

How to explain the diagnosis

Central to delivering a diagnosis of FND is the language used—a topic that has been debated for decades. Pejorative terms such as “hysteria” or “pseudoseizures” are no longer acceptable.^{154 155} However, debate continues regarding use of “functional” versus “psychogenic.” Support for the term “functional” includes: (1) it is a neutral framing that embraces the causes and mechanistic heterogeneity of this condition within the biopsychosocial model; (2) it parallels the transient reversibility of symptoms reflected in diagnostic criteria; (3) it aids acceptance given the lack of overt connotations to a mental health condition; and (4) it avoids mind-body dualism.¹⁵⁶⁻¹⁵⁹ Arguments in favor of “psychogenic” include: (1) it is a term that closely couples the condition with psychiatric or psychological care; (2) the term “functional” has a complicated history itself—and patients experience their symptoms as “dysfunctional”; and (3) “functional” can contribute to a sometimes unhelpful function-versus-structure dualism (replacing one misguided dualistic framing with another).¹⁶⁰⁻¹⁶²

Practitioners in movement disorders have shifted from “psychogenic” to “functional movement disorders,” while epileptologists continue to debate “functional seizures” versus “dissociative seizures” versus “psychogenic nonepileptic seizures” versus other terms (eg, non-epileptic attack disorder).¹⁶³⁻¹⁶⁵ Another complicating factor is lack of agreement regarding use of “seizure” for paroxysmal FND versus “events” or “attacks.”^{166 167}

We support “functional neurological disorder” as the overarching diagnostic category, and thus our opinion is that FND subtypes should carry the qualifier “functional.” This should not be interpreted, however, as invalidating that FND remains at the intersection of neurology and psychiatry, and research suggests that the language used around the diagnosis is also important in framing the condition.¹⁶⁸ Lastly, the “organic” versus “non-organic” label is falsely dualistic of brain and mind, and calls for its removal are well founded.¹⁶⁹

Efforts have been made to operationalize the communication of a FND diagnosis—framing this as the first step in treatment.¹⁷⁰ Early strategies to conceptualize the diagnosis as “good news” have fallen out of favor as this can be perceived as invalidating.¹⁷⁰ In the literature, communication approaches recommended by Hall-Patch and colleagues include: (1) validating symptoms as genuine and common; (2) naming the condition; (3) providing a brief mechanistic explanation (eg, “brain becomes overloaded and shuts down”); (4) addressing effective and ineffective treatments; and (5) fostering a hopeful sentiment of improvement (eg, pointing out that treatments are available).¹⁷¹

Describing FND as akin to a “software rather than a hardware problem” and symptoms occurring when the “computer crashes” is another helpful mechanistic explanation.^{172 173} Showing patients their “rule-in” signs and referring back to those

features when discussing the diagnosis is an expert recommendation that has been widely adopted¹⁷⁴; similarly, focusing on the “what” of diagnosis rather than the “why” can help avoid overly simplistic attempts to link FND symptoms to stress.¹⁷⁵ High yield “how to” articles written by FND experts have further contributed to the dissemination of good clinical practices regarding communication approaches^{172 176 177} (an illustration on how to deliver a diagnosis is provided in video 1, supplementary file 1). Additionally, providing patients with written materials (eg, www.neurosymptoms.org) can further enhance their understanding.¹⁵⁶ Improved education for physicians is needed around delivery of the diagnosis,¹⁷⁸ given that neurologists find FND a challenging condition to discuss with patients and to document within the medical records.¹⁷⁹⁻¹⁸² A clinical practice survey in 2018 from the International Parkinson and Movement Disorder Society, however, showed that members were more likely to communicate a FND diagnosis without ordering unnecessary tests compared with 10 years prior.¹⁸³

Treatments

Therapeutic options range from explanation alone to complex multidisciplinary rehabilitation. Triaging patients in the appropriate pathway is important and should, when possible, be individualized to specific clinical characteristics (an illustrated guide on triaging managing decisions is presented in video 2, supplementary file 2).

Communicating the diagnosis as the first step

In patients with seizure type FND, delivery of the diagnosis has been shown in a small minority of patients to result in cessation of seizures or decreased seizure frequency.¹⁸⁴⁻¹⁸⁹ For example, in 54 patients with seizure type FND, 27% achieved seizure remission in the week following communication of the diagnosis.¹⁸⁵ Studies have also shown that post diagnosis, emergency department visits and inpatient hospitalizations decrease,⁴³ with a shift toward increased use of outpatient psychiatric services.¹⁹⁰ However, it remains unclear if fewer patients overall are utilizing healthcare.¹⁹¹ The long term benefit of communicating a diagnosis of seizure type FND, if used in isolation, also remains unclear.¹⁹² More research is needed on the immediate impact of the delivery of a diagnosis of FND in other subtypes.

Psychoeducation

Several studies have investigated the efficacy of augmented educational interventions.¹⁹³⁻¹⁹⁷ One RCT evaluated the benefit of three monthly psychoeducation sessions (n=34) versus routine follow-up (n=30) in patients with seizure type FND.¹⁹⁴ No differences in frequency of seizures were recorded at a group level; however, the intervention group reported significantly improved psychosocial functioning at three and six months. In a mixed FND cohort (n=193) and their relatives (n=152)

attending a single 105 minute multidisciplinary education session, significant increases in diagnostic understanding, acceptance, belief of treatability, and hopefulness were observed across all in attendance.¹⁹⁶ A large RCT in 186 patients with functional motor symptoms investigated the efficacy of online education and self-help interventions (n=93) compared with usual care alone (n=93), and showed no incremental benefits in terms of health status.¹⁹⁷ Thus, while patient satisfaction is generally high for educational initiatives, their use in isolation does not appear to positively affect recovery from FND.

Physiotherapy and other rehabilitation

Physiotherapy is a first line treatment for patients with motor FND, although more research is needed to optimize and personalize its use. A major advance was the publication of the 2015 consensus recommendations for physiotherapy for functional motor symptoms.¹⁹⁸ General components of physiotherapy for motor FND as outlined in this article include: education on FND, creating a positive expectation of improvement, open and consistent communication between the multidisciplinary team and patient, limited hands-on interventions, encouraging early weight bearing, focusing on task completion and automatic movements, avoiding use of adaptive equipment where possible, and physiotherapy should be psychologically informed, including recognizing and exploring unhelpful thoughts and behaviors. In terms of rehabilitation strategies, an important principle of physiotherapy for motor FND is emphasis on motor retraining in the context of diverted attention (leveraging observations that functional neurological symptoms improve with distraction and worsen when attention is drawn toward the body).

Two RCTs have shown the efficacy of physiotherapy for motor FND to date.^{199 200} In the first, 60 patients with a functional gait disorder were randomized to a three week inpatient multidisciplinary rehabilitation (n=31) versus a wait list control (n=29).¹⁹⁹ The intervention, performed by a multidisciplinary rehabilitation team, consisted of “adapted physical activity with an educational and cognitive behavioral frame of reference.” Core treatments included symptom explanation and actively reinforcing normal movements. The intervention was associated with a significant improvement in walking ability and quality of life, with gait improvements sustained at one month and one year post treatment. In the second study, 60 patients with motor FND were randomized to five consecutive days in a day hospital for specialized neurophysiotherapy (eight 45-90 minute sessions; n=30) versus referral to non-specialist local neurophysiotherapy (n=30).²⁰⁰ At six months, 72% in the intervention group rated their symptoms as improved (versus 18% in controls). Significant improvements in physical and social functioning were also reported in patients assigned to specialized neurophysiotherapy.

Additionally, a 2013 systematic review reported on 29 cohort studies that also underscored the efficacy of physiotherapy in most patients treated.²⁰¹ Nonetheless, unanswered questions remain, including optimal and cost effective treatment settings (eg, inpatient versus day hospital versus outpatient versus tele-physiotherapy) as well as the optimal frequency and intensity.^{19 202 203} More research is needed regarding the management of commonly present non-motor symptoms, such as pain and fatigue.²⁰⁴ A large scale RCT for physiotherapy for motor FND currently under way will likely help clarify some of these questions.²⁰⁵

Another rehabilitative treatment that requires more research is occupational therapy. Recently published consensus recommendations for occupational therapy²⁰⁶ will help standardize interventions in this area. While beyond the scope of this article, speech and language interventions for functional speech/voice disorders may also be helpful in patients with FND—with consensus recommendations published in 2021.²⁰⁷ Additionally, given the phenotypic heterogeneity found in FND, published consensus recommendations to standardize outcomes measures for clinical trials across core FND symptoms, other physical and mental health symptoms, life impact, healthcare economics, and adverse event reporting, will positively influence research on FND in future.^{208 209}

Psychotherapy

RCTs have examined the efficacy of cognitive behavioral therapy (CBT) in treating patients with seizure type FND.²¹⁰ A 2010 pilot RCT performed in seizure type FND (n=66) compared 12 weekly conventional CBT sessions plus standard medical care (SMC) (n=33) versus SMC alone (n=33).²¹¹ This study showed that at the end of treatment the CBT group exhibited a reduced frequency of seizures compared with controls (CBT group: monthly seizure frequency at end of treatment 2.0 (interquartile range (IQR) 6.0) versus 6.75 (IRQ 38.63) in the SMC group. However, at six months after the end of treatment this effect did not remain significant. A four arm pilot RCT compared 12 session, weekly CBT informed psychotherapy (n=9) versus CBT plus sertraline (n=9) versus sertraline (n=9) versus standard care (n=7), identifying significant seizure frequency reductions (>50%) in both CBT alone (post-treatment/pre-treatment seizure ratio, mean=0.49; standard error (SE)=0.1; 95% confidence interval (CI) 0.28 to 0.84) and CBT plus sertraline (post-treatment/pre-treatment seizure ratio, mean=0.41 (SE=0.1); 95% CI 0.21 to 0.79).²¹² Patient and psychotherapist manuals for the 2014 study have been published.²¹³

The largest published RCT to date, the cognitive behavioral therapy versus standardized medical care for adults with dissociative non-epileptic seizures trial, compared 186 patients receiving 12 session, weekly conventional CBT plus standardized medical care versus 182 patients receiving standardized medical care alone. At 12 months post randomization,

no significant differences in four week seizure frequency (CBT plus standardized medical care: median four seizures (IQR 0-20) versus standardized medical care: median seven seizures (IQR 1-35)) were observed across treatment arms—albeit both groups improved from baseline. Additionally, nine of 16 secondary outcomes (eg, somatic symptoms, psychosocial functioning) improved differentially in the CBT plus standardized medical care treatment group. In a separate pilot RCT of 60 patients with seizure type FND, the addition of motivational interviewing to CBT (versus CBT alone) improved treatment adherence and decreased frequency of seizures.²¹⁴

CBT has also been evaluated in RCTs across several other FND populations. In a mixed cohort, 127 patients were randomized to self-guided CBT plus usual care (n=64) versus usual care alone (n=63), with a significant (but modest) increased proportion of individuals in the CBT arm reporting being “better” or “much better” at three months²¹⁵ (adjusted common odds ratio for an improved outcome with CBT plus usual care of 2.36 (95% CI 1.17 to 4.74). Health anxiety and somatic symptom burden also improved in the CBT arm; the self-guided CBT manual used in this study has been published.²¹⁶ A pilot 12 week RCT in motor FND compared CBT (n=14) with CBT plus adjunctive physical activity (n=15) with SMC (n=8), and showed significant improvements in FND severity, as well as secondary measures (ie, depression, anxiety, somatic symptom scales) in the CBT alone and CBT plus activity groups compared with SMC.²¹⁷

A systematic review of prospective psychotherapy clinical trials published in FND to date was published in 2020.²¹⁸

Psychopharmacology

Several studies have investigated the efficacy of serotonergic based medications in the treatment of patients with FND.²¹⁹⁻²²¹ In general, selective serotonin reuptake inhibitors and serotonin norepinephrine reuptake inhibitors (or other indicated psychotropic medications) are used to manage concurrent mental health symptoms, but are not themselves indicated for the direct treatment of FND.²²²

Emerging treatments

Studies have set out to evaluate the efficacy of mindfulness based therapy,²²³ prolonged exposure,²²⁴ psychodynamic psychotherapy,²²⁵ and group psychotherapy (including dialectical behavioral therapy)²²⁶⁻²²⁸ in patients with FND. Additionally, hypnosis has been studied in RCTs across inpatient²²⁹ and outpatient settings,²³⁰ with data supporting that outpatients with functional motor symptoms receiving hypnosis improved relative to baseline and a wait list control.²³⁰ Telepsychotherapy also appears to be a viable option in patients with FND.²³¹ Given the mixed psychotherapy results observed in the FND field to date, it remains to

be determined if guiding treatment selection based on clinical (biopsychosocial) formulations may yield more efficacious and consistent results.²³²

In addition to psychotherapy, evidence from RCTs supports using botulinum neurotoxin in the management of functional motor symptoms, although the mechanism of action may be related to placebo effects.^{233 234} Emerging evidence, yet limited, supports transcranial magnetic stimulation, although it remains unclear if potential efficacy is owing to circuit-specific neuromodulation or placebo response.^{29 235-238} More research is needed to identify potential therapeutic roles for therapeutic sedation, botulinum neurotoxin treatment, placebo, hypnosis, and virtual reality interventions among other promising management strategies.^{234 239-243}

In addition to targeting functional motor and seizure symptoms, many patients with FND experience mental health concerns (depression, anxiety, post-traumatic stress disorder, etc) and/or non-motor symptoms (pain, fatigue, dizziness, cognitive symptoms, etc) that require active consideration in the development of patient-centered treatment plans. Additionally, while only a subset of patients with FND reports adverse life experiences (eg, childhood maltreatment),²⁴⁴ these factors can be linked to increased severity of symptoms and higher rates of psychiatric comorbidities.²⁴⁵ More work is needed regarding how to optimally consider concurrently present mental health factors and non-motor physical symptoms in the treatment of patients with FND.

Guidelines

The American Neuropsychiatric Association's Committee on Research has established expert opinion statements regarding the diagnostic approach to motor FND¹² and seizure type FND,¹³ integrating both neurological and psychiatric aspects. However, these articles do not offer guidelines regarding how to optimally manage prominent mental health problems and/or pain, fatigue, and cognitive symptoms when present. A task force from the International League Against Epilepsy proposed in 2013 minimum requirements for the diagnosis of seizure type FND,¹²² yet a need persists to better integrate neurological and psychiatric diagnostic criteria for FND. It is anticipated that the FND Society will be working to further operationalize the diagnostic and therapeutic approach to FND.

Conclusion

The diagnosis of the two most frequent presentations of functional neurological disorder (motor and seizure type) is now a rule-in approach where the careful use of positive clinical signs and neurological expertise is mandatory. A thorough physical examination is necessary and the diagnosis should not rely solely on medical record review and/or a history of associated risk factors (eg, a traumatic life event) that is non-specific for the diagnosis. The use of adjunctive electrophysiological tests such as movement

recordings for tremor or EEG for seizure type FND help ascertain the diagnosis in diagnostically challenging cases; capturing a typical seizure event on video EEG and appreciating the absence of an electrographic correlate helps make a diagnosis of documented seizure type FND. Other neurological tests are not required for the diagnosis but may help if an underlying coexisting neurological disorder is suspected. A dual diagnosis of FND and another neurological disorder should always be considered. Once the diagnosis is made, a clear explanation should be provided to the patient, avoiding the term “medically unexplained.” Educational materials in the form of written or web based information can help patients understand the disorder, build social supports, and achieve diagnostic understanding through collaboration with patient support groups if desired. Early treatment should be based on the patient’s physical symptoms and informed by the biopsychosocial model, taking into consideration relevant mental health and sociocultural factors. First line treatments for both motor FND and seizure type FND include education and psychotherapy; physiotherapy is also first line treatment for motor FND. Many patients benefit from a multidisciplinary approach that incorporates neuropsychiatric and rehabilitation perspectives. Neurological follow-up should be routinely scheduled, as new symptoms can arise that require clinical triage. Lastly, research efforts are under way to aid the development of novel, biologically informed treatments.

QUESTIONS FOR FUTURE RESEARCH

- Given that individual positive signs vary in their sensitivities and specificities, what would be the added value to combine signs in a score and what cut off could bring the best positive predictive value to improve the diagnosis of FND?
- Other than additional comorbidities (eg, fatigue, pain, anxiety, and depression) can we identify prognostic markers that can help clinicians and patients choose the optimal treatment path for a given patient?
- In addition to physiotherapy, psychotherapy, and multidisciplinary treatments, can we refine neuromodulation protocols (informed by neural circuit models of FND) to obtain a clinically significant therapeutic effect?

PATIENT INVOLVEMENT

The CEO and founder of a charity dedicated to patients with FND (FND Hope) reviewed a draft of this manuscript and made suggestions and edits on the content and presentation; the main suggestions concerned avoiding a lengthy historical background section which we agreed with and general feedback on the terminology used for FND.

Contributors: SA drafted the outline of the draft, designed collection of data, collected and analyzed data, critically appraised data, and wrote and edited the final manuscript. She is guarantor.

DLP drafted the outline of the draft, designed collection of data, collected and analyzed data, critically appraised data, and wrote and edited the final manuscript. He is guarantor.

Both SA and DLP selected the relevant articles, reviewed critically the literature, and wrote the manuscript.

The CEO and Founder of a charity contributed to editing the final manuscript.

Funding: SA was supported by the Swiss National Science Foundation grant PP00P3_176985.

DLP was supported by the Sidney R Baer Jr Foundation.

Competing interests: SA has received honorariums for continuing medical education lectures in functional neurological disorder, is a member of the research committee of the American Neuropsychiatric Association, and is on the editorial board of *Cognitive Neuropsychiatry*.

DLP has received honorariums for continuing medical education lectures in functional neurological disorder, is a member of the research committee of the American Neuropsychiatric Association, and is on the editorial board of *Epilepsy and Behavior*.

Provenance and peer review: Commissioned; externally peer reviewed.

- 1 Stephen CD, Fung V, Lungu CI, Espay AJ. Assessment of emergency department and inpatient use and costs in adult and pediatric functional neurological disorders. *JAMA Neurol* 2021;78:88-101. doi:10.1001/jamaneurol.2020.3753
- 2 Stone J, Burton C, Carson A. Recognising and explaining functional neurological disorder. *BMJ* 2020;371:m3745. doi:10.1136/bmj.m3745
- 3 Trimble M, Reynolds EH. A brief history of hysteria: from the ancient to the modern. In: Hallett M, Stone J, Carson A, eds. *Handbook of Clinical Neurology*. Elsevier, 2016: 3-10.
- 4 Burke MJ. “It’s all in your head” —Medicine’s silent epidemic. *JAMA Neurol* 2019;76:1417-8. doi:10.1001/jamaneurol.2019.3043
- 5 Hallett M. Psychogenic movement disorders: a crisis for neurology. *Curr Neurol Neurosci Rep* 2006;6:269-71. doi:10.1007/s11910-006-0015-x
- 6 Keynejad RC, Carson AJ, David AS, Nicholson TR. Functional neurological disorder: psychiatry’s blind spot. *Lancet Psychiatry* 2017;4:e2-3. doi:10.1016/S2215-0366(17)30036-6
- 7 Aybek S. Corticolimbic fast-tracking in functional neurological disorders: towards understanding of the ‘dynamic lesion’ of Jean-Martin Charcot. *J Neurol Neurosurg Psychiatry* 2019;90:845. doi:10.1136/jnnp-2019-320597
- 8 Drane DL, Fani N, Hallett M, Khalsa SS, Perez DL, Roberts NA. A framework for understanding the pathophysiology of functional neurological disorder. *CNS Spectr* 2020;Sep 4:1-7.
- 9 Perez DL, Haller AL, Espay AJ. Should neurologists diagnose and manage functional neurologic disorders? It is complicated. *Neurol Clin Pract* 2019;9:165-7. doi:10.1212/CPJ.0000000000000573
- 10 Espay AJ, Goldenhar LM, Voon V, Schrag A, Burton N, Lang AE. Opinions and clinical practices related to diagnosing and managing patients with psychogenic movement disorders: An international survey of movement disorder society members. *Mov Disord* 2009;24:1366-74. doi:10.1002/mds.22618
- 11 Aybek S, Lidstone SC, Nielsen G, et al. What is the role of a specialist assessment clinic for FND? Lessons from three national referral centers. *J Neuropsychiatry Clin Neurosci* 2020;32:79-84. doi:10.1176/appi.neuropsych.19040083
- 12 Perez DL, Aybek S, Popkirov S, et al. On behalf of the American Neuropsychiatric Association Committee for Research. A review and expert opinion on the neuropsychiatric assessment of motor functional neurological disorders. *J Neuropsychiatry Clin Neurosci* 2021;33:14-26. doi:10.1176/appi.neuropsych.19120357
- 13 Baslet G, Bajestan SN, Aybek S, et al. Evidence-based practice for the clinical assessment of psychogenic nonepileptic seizures: a report from the American Neuropsychiatric Association Committee on Research. *J Neuropsychiatry Clin Neurosci* 2021;33:27-42. doi:10.1176/appi.neuropsych.19120354
- 14 Jacob AE, Smith CA, Jablonski ME, et al. Multidisciplinary clinic for functional movement disorders (FMD): 1-year experience from a single centre. *J Neurol Neurosurg Psychiatry* 2018;89:1011-2. doi:10.1136/jnnp-2017-316523
- 15 Perez DL, Young SS, King JN, et al. Preliminary predictors of initial attendance, symptom burden, and motor subtype in a US functional neurological disorders clinic population. *Cogn Behav Neurol* 2016;29:197-205. doi:10.1097/WNN.000000000000106
- 16 Aybek S, Lidstone SC, Nielsen G, et al. What is the role of a specialist assessment clinic for FND? Lessons from three national referral centers. *J Neuropsychiatry Clin Neurosci* 2020;32:79-84. doi:10.1176/appi.neuropsych.19040083

- 17 Stone JC. A. [I couldn't view this link – could you check?] [Authors to check]https://www.google.com/url?sa=t&rtct=j&q=&esrc=s&source=web&cd=&ved=2ahUKewj0-jg1aHxhAhVwEC-wKHYVqAQsQFjAeegQIAxAD&url=http%3A%2F%2Fhealthcareimprovement.scotland.org%2Ffidoc.ashx%3Fdocid%3D6d302668-ff12-4f53-b547-72c19531e37a%26version%3D1&usq=AOVaw0zz6Pgrlc-z05U79zHIRSY. 2012.
- 18 Edwards MJ. Functional neurological disorder: an ethical turning point for neuroscience. *Brain* 2019;142:1855-7. doi:10.1093/brain/awz194
- 19 Jacob AE, Kaelin DL, Roach AR, Ziegler CH, LaFaver K. Motor retraining (MoRe) for functional movement disorders: outcomes from a 1-week multidisciplinary rehabilitation program. *PM R* 2018;10:1164-72. doi:10.1016/j.pmrj.2018.05.011
- 20 McCormack R, Moriarty J, Mellers JD, et al. Specialist inpatient treatment for severe motor conversion disorder: a retrospective comparative study. *J Neurol Neurosurg Psychiatry* 2014;85:895-900. doi:10.1136/jnnp-2013-305716
- 21 Nielsen G, Ricciardi L, Demartini B, Hunter R, Joyce E, Edwards MJ. Outcomes of a 5-day physiotherapy programme for functional (psychogenic) motor disorders. *J Neurol* 2015;262:674-81. doi:10.1007/s00415-014-7631-1
- 22 McKee K, Glass S, Adams C, et al. The inpatient assessment and management of motor functional neurological disorders: an interdisciplinary perspective. *Psychosomatics* 2018;59:358-68. doi:10.1016/j.psym.2017.12.006
- 23 Goldstein LH, Robinson EJ, Mellers JDC, et al, CODES study group. Cognitive behavioural therapy for adults with dissociative seizures (CODES): a pragmatic, multicentre, randomised controlled trial. *Lancet Psychiatry* 2020;7:491-505. doi:10.1016/S2215-0366(20)30128-0
- 24 Stone J. Functional neurological disorders: the neurological assessment as treatment. *Neurophysiol Clin* 2014;44:363-73. doi:10.1016/j.neucli.2014.01.002
- 25 Edwards MJ, Adams RA, Brown H, Pareés I, Friston KJ. A Bayesian account of 'hysteria'. *Brain* 2012;135:3495-512. doi:10.1093/brain/awt129
- 26 Perez DL, Dworetzky BA, Dickerson BC, et al. An integrative neurocircuit perspective on psychogenic nonepileptic seizures and functional movement disorders: neural functional unawareness. *Clin EEG Neurosci* 2015;46:4-15. doi:10.1177/1550059414555905
- 27 Garcin B, Roze E, Mesrati F, et al. Transcranial magnetic stimulation as an efficient treatment for psychogenic movement disorders. *J Neurol Neurosurg Psychiatry* 2013;84:1043-6. doi:10.1136/jnnp-2012-304062
- 28 McWhirter L, Ludwig L, Carson A, McIntosh RD, Stone J. Transcranial magnetic stimulation as a treatment for functional (psychogenic) upper limb weakness. *J Psychosom Res* 2016;89:102-6. doi:10.1016/j.jpsychores.2016.08.010
- 29 Peterson KT, Kosior R, Meek BP, Ng M, Perez DL, Modirrousta M. Right temporoparietal junction transcranial magnetic stimulation in the treatment of psychogenic nonepileptic seizures: a case series. *Psychosomatics* 2018;59:601-6. doi:10.1016/j.psym.2018.03.001
- 30 Baslet G, Ehlerl A, Oser M, Dworetzky BA. Mindfulness-based therapy for psychogenic nonepileptic seizures. *Epilepsy Behav* 2020;103(Pt A):106534. doi:10.1016/j.yebeh.2019.106534
- 31 Ball HA, McWhirter L, Ballard C, et al. Functional cognitive disorder: dementia's blind spot. *Brain* 2020;143:2895-903. doi:10.1093/brain/awaa224
- 32 Popkirov S, Staab JP, Stone J. Persistent postural-perceptual dizziness (PPPD): a common, characteristic and treatable cause of chronic dizziness. *Pract Neurol* 2018;18:5-13. doi:10.1136/practneurol-2017-001809
- 33 Stone J, Carson A, Duncan R, et al. Who is referred to neurology clinics?—The diagnoses made in 3781 new patients. *Clin Neurol Neurosurg* 2010;112:747-51. doi:10.1016/j.clineuro.2010.05.011
- 34 Carson AJ, Ringbauer B, Stone J, McKenzie L, Warlow C, Sharpe M. Do medically unexplained symptoms matter? A prospective cohort study of 300 new referrals to neurology outpatient clinics. *J Neurol Neurosurg Psychiatry* 2000;68:207-10. doi:10.1136/jnnp.68.2.207
- 35 Carson A, Stone J, Hibberd C, et al. Disability, distress and unemployment in neurology outpatients with symptoms 'unexplained by organic disease'. *J Neurol Neurosurg Psychiatry* 2011;82:810-3. doi:10.1136/jnnp.2010.220640
- 36 Carson A, Lehn A. Epidemiology. *Handb Clin Neurol* 2016;139:47-60. doi:10.1016/B978-0-12-801772-2.00005-9
- 37 Gelauff J, Stone J, Edwards M, Carson A. The prognosis of functional (psychogenic) motor symptoms: a systematic review. *J Neurol Neurosurg Psychiatry* 2014;85:220-6. doi:10.1136/jnnp-2013-305321
- 38 Stefánsson JG, Messina JA, Meyerowitz S. Hysterical neurosis, conversion type: clinical and epidemiological considerations. *Acta Psychiatr Scand* 1976;53:119-38. doi:10.1111/j.1600-0447.1976.tb00066.x
- 39 Stevens DL. Neurology in Gloucestershire: the clinical workload of an English neurologist. *J Neurol Neurosurg Psychiatry* 1989;52:439-46. doi:10.1136/jnnp.52.4.439
- 40 Binzer M, Andersen PM, Kullgren G. Clinical characteristics of patients with motor disability due to conversion disorder: a prospective control group study. *J Neurol Neurosurg Psychiatry* 1997;63:83-8. doi:10.1136/jnnp.63.1.83
- 41 Stone J, Warlow C, Sharpe M. The symptom of functional weakness: a controlled study of 107 patients. *Brain* 2010;133:1537-51. doi:10.1093/brain/awq068
- 42 Sigurdardóttir KR, Olafsson E. Incidence of psychogenic seizures in adults: a population-based study in Iceland. *Epilepsia* 1998;39:749-52. doi:10.1111/j.1528-1157.1998.tb01161.x
- 43 Razvi S, Mulhern S, Duncan R. Newly diagnosed psychogenic nonepileptic seizures: health care demand prior to and following diagnosis at a first seizure clinic. *Epilepsy Behav* 2012;23:7-9. doi:10.1016/j.yebeh.2011.10.009
- 44 Villagrán A, Eldøen G, Duncan R, Aaberg KM, Hofoss D, Lossius MI. Incidence and prevalence of psychogenic nonepileptic seizures in a Norwegian county: A 10-year population-based study. *Epilepsia* 2021;62:1528-35. doi:10.1111/epi.16949
- 45 Baizabal-Carvallo JF, Jankovic J. Functional (psychogenic) stereotypes. *J Neurol* 2017;264:1482-7. doi:10.1007/s00415-017-8551-7
- 46 Fekete R, Baizabal-Carvallo JF, Ha AD, Davidson A, Jankovic J. Convergence spasm in conversion disorders: prevalence in psychogenic and other movement disorders compared with controls. *J Neurol Neurosurg Psychiatry* 2012;83:202-4. doi:10.1136/jnnp-2011-300733
- 47 Teodoro T, Cunha JM, Abreu LF, Yogarajah M, Edwards MJ. Abnormal eye and cranial movements triggered by examination in people with functional neurological disorder. *Neuroophthalmology* 2019;43:240-3. doi:10.1080/01658107.2018.1536998
- 48 Daum C, Gheorghita F, Spatola M, et al. Interobserver agreement and validity of bedside 'positive signs' for functional weakness, sensory and gait disorders in conversion disorder: a pilot study. *J Neurol Neurosurg Psychiatry* 2015;86:425-30. doi:10.1136/jnnp-2013-307381
- 49 Stone J, Warlow C, Sharpe M. The symptom of functional weakness: a controlled study of 107 patients. *Brain* 2010;133:1537-51. doi:10.1093/brain/awq068
- 50 Laub HN, Dwivedi AK, Revilla FJ, Duker AP, Pecina-Jacob C, Espay AJ. Diagnostic performance of the "Huffing and Puffing" sign in psychogenic (functional) movement disorders. *Mov Disord Clin Pract* 2015;2:29-32. doi:10.1002/mdc3.12102
- 51 Okun MS, Rodriguez RL, Foote KD, Fernandez HH. The "chair test" to aid in the diagnosis of psychogenic gait disorders. *Neurologist* 2007;13:87-91. doi:10.1097/01.nrl.0000256358.52613.cc
- 52 Baizabal-Carvallo JF, Jankovic J. Distinguishing features of psychogenic (functional) versus organic hemifacial spasm. *J Neurol* 2017;264:359-63. doi:10.1007/s00415-016-8356-0
- 53 Horn D, Galli S, Berney A, Vingerhoets F, Aybek S. Testing head rotation and flexion is useful in functional limb weakness. *Mov Disord Clin Pract* 2017;4:597-602. doi:10.1002/mdc3.12492
- 54 Pizzi F, Antelmi E, Vandi S, et al. The distinguishing motor features of cataplexy: a study from video-recorded attacks. *Sleep* 2018;41:41. doi:10.1093/sleep/zy026
- 55 Chabrol H, Peresson G, Claret M. Lack of specificity of the traditional criteria for conversion disorders. *Eur Psychiatry* 1995;10:317-9. doi:10.1016/0924-9338(96)80314-2
- 56 Baker JH, Silver JR. Hysterical paraplegia. *J Neurol Neurosurg Psychiatry* 1987;50:375-82. doi:10.1136/jnnp.50.4.375
- 57 Daum C, Aybek S. Validity of the "Drift without pronation" sign in conversion disorder. *BMC Neurol* 2013;13:31. doi:10.1186/1471-2377-13-31
- 58 Tinazzi M, Simonetto S, Franco L, et al. Abduction finger sign: a new sign to detect unilateral functional paralysis of the upper limb. *Mov Disord* 2008;23:2415-9. doi:10.1002/mds.22268
- 59 Lombardi TL, Barton E, Wang J, et al. The elbow flex-ex: a new sign to detect unilateral upper extremity non-organic paresis. *J Neurol Neurosurg Psychiatry* 2014;85:165-7. doi:10.1136/jnnp-2012-304314
- 60 Sonoo M. Abductor sign: a reliable new sign to detect unilateral non-organic paresis of the lower limb. *J Neurol Neurosurg Psychiatry* 2004;75:121-5.
- 61 McWhirter L, Stone J, Sandercock P, Whiteley W. Hoover's sign for the diagnosis of functional weakness: a prospective unblinded cohort study in patients with suspected stroke. *J Psychosom Res* 2011;71:384-6. doi:10.1016/j.jpsychores.2011.09.003
- 62 Yugué I, Shiba K, Ueta T, Iwamoto Y. A new clinical evaluation for hysterical paralysis. *Spine (Phila Pa 1976)* 2004;29:1910-3. discussion 1913. doi:10.1097/01.brs.0000137055.55350.37
- 63 van der Stouwe AMM, Elting JW, van der Hoeven JH, et al. How typical are 'typical' tremor characteristics? Sensitivity and specificity of five tremor phenomena. *Parkinsonism Relat Disord* 2016;30:23-8. doi:10.1016/j.parkreldis.2016.06.008

- 64 Seneviratne U, Minato E, Paul E. How reliable is ictal duration to differentiate psychogenic nonepileptic seizures from epileptic seizures? *Epilepsy Behav* 2017;66:127-31. doi:10.1016/j.yebeh.2016.10.024
- 65 Syed TU, LaFrance WC Jr, Kahrman ES, et al. Can semiology predict psychogenic nonepileptic seizures? A prospective study. *Ann Neurol* 2011;69:997-1004. doi:10.1002/ana.22345
- 66 Bazil CW, Walczak TS. Effects of sleep and sleep stage on epileptic and nonepileptic seizures. *Epilepsia* 1997;38:56-62. doi:10.1111/j.1528-1157.1997.tb01077.x
- 67 Jedrzejczak J, Owczarek K, Majkowski J. Psychogenic pseudoepileptic seizures: clinical and electroencephalogram (EEG) video-tape recordings. *Eur J Neurol* 1999;6:473-9. doi:10.1046/j.1468-1331.1999.640473.x
- 68 De Paola L, Terra VC, Silvano CE, et al. Improving first responders' psychogenic nonepileptic seizures diagnosis accuracy: Development and validation of a 6-item bedside diagnostic tool. *Epilepsy Behav* 2016;54:40-6. doi:10.1016/j.yebeh.2015.10.025
- 69 Slater JD, Brown MC, Jacobs W, Ramsay RE. Induction of pseudoseizures with intravenous saline placebo. *Epilepsia* 1995;36:580-5. doi:10.1111/j.1528-1157.1995.tb02571.x
- 70 Azar NJ, Tayah TF, Wang L, Song Y, Abou-Khalil BW. Postictal breathing pattern distinguishes epileptic from nonepileptic convulsive seizures. *Epilepsia* 2008;49:132-7. doi:10.1111/j.1528-1167.2007.01215.x
- 71 Vogrig A, Hsiang JC, Ng J, Rolnick J, Cheng J, Parvizi J. A systematic study of stereotypy in epileptic seizures versus psychogenic seizure-like events. *Epilepsy Behav* 2019;90:172-7. doi:10.1016/j.yebeh.2018.11.030
- 72 Saygi S, Katz A, Marks DA, Spencer SS. Frontal lobe partial seizures and psychogenic seizures: comparison of clinical and ictal characteristics. *Neurology* 1992;42:1274-7. doi:10.1212/WNL.42.7.1274
- 73 Chen DK, Graber KD, Anderson CT, Fisher RS. Sensitivity and specificity of video alone versus electroencephalography alone for the diagnosis of partial seizures. *Epilepsy Behav* 2008;13:115-8. doi:10.1016/j.yebeh.2008.02.018
- 74 Devinsky O, Sanchez-Villaseñor F, Vazquez B, Kothari M, Alper K, Luciano D. Clinical profile of patients with epileptic and nonepileptic seizures. *Neurology* 1996;46:1530-3. doi:10.1212/WNL.46.6.1530
- 75 Bell WL, Park YD, Thompson EA, Radtke RA. Ictal cognitive assessment of partial seizures and pseudoseizures. *Arch Neurol* 1998;55:1456-9. doi:10.1001/archneur.55.11.1456
- 76 Chung SS, Gerber P, Kirlin KA. Ictal eye closure is a reliable indicator for psychogenic nonepileptic seizures. *Neurology* 2006;66:1730-1. doi:10.1212/01.wnl.0000218160.31537.87
- 77 Gates JR, Ramani V, Whalen S, Loewenson R. Ictal characteristics of pseudoseizures. *Arch Neurol* 1985;42:1183-7. doi:10.1001/archneur.1985.04060110065017
- 78 DeToledo JC, Ramsay RE. Patterns of involvement of facial muscles during epileptic and nonepileptic events: review of 654 events. *Neurology* 1996;47:621-5. doi:10.1212/WNL.47.3.621
- 79 Geyer JD, Payne TA, Drury I. The value of pelvic thrusting in the diagnosis of seizures and pseudoseizures. *Neurology* 2000;54:227-9. doi:10.1212/WNL.54.1.227
- 80 Walczak TS, Bogoljubov A. Weeping during psychogenic nonepileptic seizures. *Epilepsia* 1996;37:208-10. doi:10.1111/j.1528-1157.1996.tb00013.x
- 81 Asadi-Pooya AA, Wyeth D, Sperling MR. Ictal crying. *Epilepsy Behav* 2016;59:1-3. doi:10.1016/j.yebeh.2016.03.012
- 82 Vossler DG, Haltiner AM, Schepp SK, et al. Ictal stuttering: a sign suggestive of psychogenic nonepileptic seizures. *Neurology* 2004;63:516-9. doi:10.1212/01.WNL.0000133208.57562.CB
- 83 Izadyar S, Shah V, James B. Comparison of postictal semiology and behavior in psychogenic nonepileptic and epileptic seizures. *Epilepsy Behav* 2018;88:123-9. doi:10.1016/j.yebeh.2018.08.020
- 84 Cervenka MC, Lesser R, Tran TT, Fortuné T, Muthugovindan D, Miglioretti DL. Does the teddy bear sign predict psychogenic nonepileptic seizures? *Epilepsy Behav* 2013;28:217-20. doi:10.1016/j.yebeh.2013.05.016
- 85 Benbadis SR, Lancman ME, King LM, Swanson SJ. Preictal pseudosleep: a new finding in psychogenic seizures. *Neurology* 1996;47:63-7. doi:10.1212/WNL.47.1.63
- 86 Oliva M, Pattison C, Carino J, Roten A, Matkovic Z, O'Brien TJ. The diagnostic value of oral lacerations and incontinence during convulsive "seizures". *Epilepsia* 2008;49:962-7. doi:10.1111/j.1528-1167.2008.01554.x
- 87 Dufresne D, Dubovec K, So NK, Kotagal P. Ictal biting injuries in the epilepsy monitoring unit, a cohort study of incidence and semiological significance. *Seizure* 2019;66:39-41. doi:10.1016/j.seizure.2019.02.005
- 88 Peguero E, Abou-Khalil B, Fakhoury T, Mathews G. Self-injury and incontinence in psychogenic seizures. *Epilepsia* 1995;36:586-91. doi:10.1111/j.1528-1157.1995.tb02572.x
- 89 Stone J, Duncan R. Tongue biting in pseudoseizures and epilepsy. *Pract Neurol* 2006;6:64-5. doi:10.1136/jnnp.2005.012345
- 90 Walczak TS, Rubinsky M. Plantar responses after epileptic seizures. *Neurology* 1994;44:2191-3. doi:10.1212/WNL.44.11.2191
- 91 Aboud O, Al-Salaimeh A, Kumar Raina S, Sahaya K, Hinduja A. Positive clinical signs in neurological diseases - An observational study. *J Clin Neurosci* 2019;59:141-5. doi:10.1016/j.jocn.2018.10.113
- 92 Baizabal-Carvallo JF, Alonso-Juarez M, Jankovic J. Functional gait disorders, clinical phenomenology, and classification. *Neurol Sci* 2020;41:911-5. doi:10.1007/s10072-019-04185-8
- 93 Criswell S, Sterling C, Swisher L, Evanoff B, Racette BA. Sensitivity and specificity of the finger tapping task for the detection of psychogenic movement disorders. *Parkinsonism Relat Disord* 2010;16:197-201. doi:10.1016/j.parkrel.2009.11.007
- 94 Kuyk J, Spinhoven P, van Dyck R. Hypnotic recall: a positive criterion in the differential diagnosis between epileptic and pseudoepileptic seizures. *Epilepsia* 1999;40:485-91. doi:10.1111/j.1528-1157.1999.tb00745.x
- 95 Sen A, Scott C, Sisodiya SM. Stertorous breathing is a reliably identified sign that helps in the differentiation of epileptic from psychogenic non-epileptic convulsions: an audit. *Epilepsy Res* 2007;77:62-4. doi:10.1016/j.eplepsyres.2007.07.009
- 96 Association AP. Diagnostic and Statistical Manual of Mental Disorders (DSM-5), 5th ed. Washington, DC:2013.
- 97 Charcot J. *Clinical lectures on diseases of the nervous system*. New Sydenham Society, 1889.
- 98 Koehler PJ, Okun MS. Important observations prior to the description of the Hoover sign. *Neurology* 2004;63:1693-7. doi:10.1212/01.WNL.0000142977.21104.94
- 99 Babinski J. Diagnostic différentiel de l'hémiplégie organique et de l'hémiplégie hystérique. *Gazette des Hôpitaux de Paris* 1900;73:533-7.
- 100 Vanek J, Prasko J, Ociskova M, et al. Sleep disturbances in patients with nonepileptic seizures. *Nat Sci Sleep* 2021;13:209-18. doi:10.2147/NSS.S289190
- 101 Nonnekens J, Růžička E, Serranová T, Reich SG, Bloem BR, Hallett M. Functional gait disorders: A sign-based approach. *Neurology* 2020;94:1093-9. doi:10.1212/WNL.00000000000009649
- 102 Kaski D, Bronstein AM, Edwards MJ, Stone J. Cranial functional (psychogenic) movement disorders. *Lancet Neurol* 2015;14:1196-205. doi:10.1016/S1474-4422(15)00226-4
- 103 Fasano A, Valadas A, Bhatia KP, et al. Psychogenic facial movement disorders: clinical features and associated conditions. *Mov Disord* 2012;27:1544-51. doi:10.1002/mds.25190
- 104 Perez DL, Hunt A, Sharma N, Flaherty A, Caplan D, Schmammann JD. Cautionary notes on diagnosing functional neurologic disorder as a neurologist-in-training. *Neurol Clin Pract* 2020;10:484-7. doi:10.1212/CPJ.0000000000000779
- 105 Anderson JR, Nakhate V, Stephen CD, Perez DL. Functional (psychogenic) neurological disorders: assessment and acute management in the emergency department. *Semin Neurol* 2019;39:102-14.
- 106 Park JE, Maurer CW, Hallett M. The 'whack-a-mole' sign in functional movement disorders. *Mov Disord Clin Pract* 2015;2:286-8. doi:10.1002/mdc3.12177
- 107 Schrag A, Trimble M, Quinn N, Bhatia K. The syndrome of fixed dystonia: an evaluation of 103 patients. *Brain* 2004;127:2360-72. doi:10.1093/brain/awh262
- 108 Frucht L, Perez DL, Callahan J, et al. Functional dystonia: differentiation from primary dystonia and multidisciplinary treatments. *Front Neurol* 2021;11:605262. doi:10.3389/fneur.2020.605262
- 109 Lang AE. Psychogenic dystonia: a review of 18 cases. *Can J Neurol Sci* 1995;22:136-43. doi:10.1017/S031716710004021X
- 110 Ganos C, Martino D, Espay AJ, Lang AE, Bhatia KP, Edwards MJ. Tics and functional tic-like movements: Can we tell them apart? *Neurology* 2019;93:750-8. doi:10.1212/WNL.00000000000008372
- 111 Demartini B, Ricciardi L, Paresi I, Ganos C, Bhatia KP, Edwards MJ. A positive diagnosis of functional (psychogenic) tics. *Eur J Neurol* 2015;22:527-e36. doi:10.1111/ene.12609
- 112 Roper LS, Saifee TA, Paresi I, Rickards H, Edwards MJ. How to use the entrainment test in the diagnosis of functional tremor. *Pract Neurol* 2013;13:396-8. doi:10.1136/practneurol-2013-000549
- 113 Thenganatt MA, Jankovic J. Psychogenic tremor: a video guide to its distinguishing features. *Tremor Other Hyperkinet Mov (N Y)* 2014;4:253. doi:10.5334/tohm.228
- 114 Wissel BD, Dwivedi AK, Merola A, et al. Functional neurological disorders in Parkinson disease. *J Neurol Neurosurg Psychiatry* 2018;89:566-71. doi:10.1136/jnnp-2017-317378
- 115 Frasca Polara G, Fleury V, Stone J, et al. Prevalence of functional (psychogenic) parkinsonism in two Swiss movement disorders clinics and review of the literature. *J Neurol Sci* 2018;387:37-45. doi:10.1016/j.jns.2018.01.022

- 116 Walzl D, Solomon AJ, Stone J. Functional neurological disorder and multiple sclerosis: a systematic review of misdiagnosis and clinical overlap. *J Neurol* 2021. doi:10.1007/s00415-021-10436-6.
- 117 Gelauff JM, Carson A, Ludwig L, Tijssen MAJ, Stone J. The prognosis of functional limb weakness: a 14-year case-control study. *Brain* 2019;142:2137-48. doi:10.1093/brain/awz138
- 118 Stone J, Smyth R, Carson A, et al. Systematic review of misdiagnosis of conversion symptoms and "hysteria". *BMJ* 2005;331:989. doi:10.1136/bmj.38628.466898.55
- 119 Lagrand T, Tuitert I, Klamer M, et al. Functional or not functional; that's the question: Can we predict the diagnosis functional movement disorder based on associated features? *Eur J Neurol* 2021;28:33-9. doi:10.1111/ene.14488
- 120 Brigo F, Nardone R, Ausserer H, et al. The diagnostic value of urinary incontinence in the differential diagnosis of seizures. *Seizure* 2013;22:85-90. doi:10.1016/j.seizure.2012.10.011
- 121 Popkirov S, Grönheit W, Wellmer J. Hyperventilation and photic stimulation are useful additions to a placebo-based suggestive seizure induction protocol in patients with psychogenic nonepileptic seizures. *Epilepsy Behav* 2015;46:88-90. doi:10.1016/j.yebeh.2015.04.020
- 122 LaFrance WCJr, Baker GA, Duncan R, Goldstein LH, Reuber M. Minimum requirements for the diagnosis of psychogenic nonepileptic seizures: a staged approach: a report from the International League Against Epilepsy Nonepileptic Seizures Task Force. *Epilepsia* 2013;54:2005-18. doi:10.1111/epi.12356
- 123 Tatum WO, Hirsch LJ, Gelfand MA, et al, OSmartVIE Investigators. Assessment of the predictive value of outpatient smartphone videos for diagnosis of epileptic seizures. *JAMA Neurol* 2020;77:593-600. doi:10.1001/jamaneurol.2019.4785
- 124 Goleva SB, Lake AM, Torstenson ES, Haas KF, Davis LK. Epidemiology of functional seizures among adults treated at a university hospital. *JAMA Netw Open* 2020;3:e2027920. doi:10.1001/jamanetworkopen.2020.27920
- 125 Kutlubaev MA, Xu Y, Hackett ML, Stone J. Dual diagnosis of epilepsy and psychogenic nonepileptic seizures: Systematic review and meta-analysis of frequency, correlates, and outcomes. *Epilepsy Behav* 2018;89:70-8. doi:10.1016/j.yebeh.2018.10.010
- 126 Wasserman D, Herskovitz M. Epileptic vs psychogenic nonepileptic seizures: a video-based survey. *Epilepsy Behav* 2017;73:42-5. doi:10.1016/j.yebeh.2017.04.020
- 127 Reuber M, Chen M, Jamnadas-Khoda J, et al. Value of patient-reported symptoms in the diagnosis of transient loss of consciousness. *Neurology* 2016;87:625-33. doi:10.1212/WNL.0000000000002948
- 128 Lenio S, Kerr WT, Watson M, et al. Validation of a predictive calculator to distinguish between patients presenting with dissociative versus epileptic seizures. *Epilepsy Behav* 2021;116:107767. doi:10.1016/j.yebeh.2021.107767
- 129 Thomsen BLC, Teodoro T, Edwards MJ. Biomarkers in functional movement disorders: a systematic review. *J Neurol Neurosurg Psychiatry* 2020;91:1261-9. doi:10.1136/jnnp-2020-323141
- 130 Schwingenschuh P, Saifee TA, Katschnig-Winter P, et al. Validation of "laboratory-supported" criteria for functional (psychogenic) tremor. *Mov Disord* 2016;31:555-62. doi:10.1002/mds.26525
- 131 Schwingenschuh P, Katschnig P, Seiler S, et al. Moving toward "laboratory-supported" criteria for psychogenic tremor. *Mov Disord* 2011;26:2509-15. doi:10.1002/mds.23922
- 132 Zeuner KE, Shoge RO, Goldstein SR, Dambrosia JM, Hallett M. Accelerometry to distinguish psychogenic from essential or parkinsonian tremor. *Neurology* 2003;61:548-50. doi:10.1212/01.WNL.0000076183.34915.CD
- 133 Sundararajan T, Tesar GE, Jimenez XF. Biomarkers in the diagnosis and study of psychogenic nonepileptic seizures: A systematic review. *Seizure* 2016;35:11-22. doi:10.1016/j.seizure.2015.12.011
- 134 Szaflarski JP, Hughes C, Szaflarski M, et al. Quality of life in psychogenic nonepileptic seizures. *Epilepsia* 2003;44:236-42. doi:10.1046/j.1528-1157.2003.35302.x
- 135 Testa SM, Scheff BK, Szaflarski JP, Yeh HS, Privitera MD. Mood, personality, and health-related quality of life in epileptic and psychogenic seizure disorders. *Epilepsia* 2007;48:973-82. doi:10.1111/j.1528-1167.2006.00965.x
- 136 Nightscales R, McCartney L, Auvrez C, et al. Mortality in patients with psychogenic nonepileptic seizures. *Neurology* 2020;95:e643-52. doi:10.1212/WNL.0000000000009855
- 137 Quigg M, Armstrong RF, Farace E, Fountain NB. Quality of life outcome is associated with cessation rather than reduction of psychogenic nonepileptic seizures. *Epilepsy Behav* 2002;3:455-9. doi:10.1016/S1525-5050(02)00524-3
- 138 Walther K, Volbers B, Erdmann L, et al. Psychological long-term outcome in patients with psychogenic nonepileptic seizures. *Epilepsia* 2019;60:669-78. doi:10.1111/epi.14682
- 139 Jones B, Reuber M, Norman P. Correlates of health-related quality of life in adults with psychogenic nonepileptic seizures: A systematic review. *Epilepsia* 2016;57:171-81. doi:10.1111/epi.13268
- 140 LaFrance WCJr, Syc S. Depression and symptoms affect quality of life in psychogenic nonepileptic seizures. *Neurology* 2009;73:366-71. doi:10.1212/WNL.0b013e3181b04c83
- 141 LaFrance WCJr, Alcoso ML, Davis JD, et al. Impact of family functioning on quality of life in patients with psychogenic nonepileptic seizures versus epilepsy. *Epilepsia* 2011;52:292-300.
- 142 Cronje G, Pretorius C. The coping styles and health-related quality of life of South African patients with psychogenic nonepileptic seizures. *Epilepsy Behav* 2013;29:581-4. doi:10.1016/j.yebeh.2013.09.045
- 143 Wolf LD, Hentz JG, Ziemba KS, et al. Quality of life in psychogenic nonepileptic seizures and epilepsy: the role of somatization and alexithymia. *Epilepsy Behav* 2015;43:81-8. doi:10.1016/j.yebeh.2014.12.010
- 144 Rawlings GH, Brown I, Reuber M. Predictors of health-related quality of life in patients with epilepsy and psychogenic nonepileptic seizures. *Epilepsy Behav* 2017;68:153-8. doi:10.1016/j.yebeh.2016.10.035
- 145 Robson C, Myers L, Pretorius C, Lian OS, Reuber M. Health related quality of life of people with non-epileptic seizures: The role of socio-demographic characteristics and stigma. *Seizure* 2018;55:93-9. doi:10.1016/j.seizure.2018.01.001
- 146 Karakis I, Janocko NJ, Morton ML, et al. Stigma in psychogenic nonepileptic seizures. *Epilepsy Behav* 2020;111:107269. doi:10.1016/j.yebeh.2020.107269
- 147 Mitchell JW, Ali F, Cavanna AE. Dissociative experiences and quality of life in patients with non-epileptic attack disorder. *Epilepsy Behav* 2012;25:307-12. doi:10.1016/j.yebeh.2012.08.022
- 148 Anderson KE, Gruber-Baldini AL, Vaughan CG, et al. Impact of psychogenic movement disorders versus Parkinson's on disability, quality of life, and psychopathology. *Mov Disord* 2007;22:2204-9. doi:10.1002/mds.21687
- 149 Gendre T, Carle G, Mesrati F, et al. Quality of life in functional movement disorders is as altered as in organic movement disorders. *J Psychosom Res* 2019;116:10-6. doi:10.1016/j.jpsychores.2018.11.006
- 150 Gelauff JM, Kingma EM, Kalkman JS, et al. Fatigue, not self-rated motor symptom severity, affects quality of life in functional motor disorders. *J Neurol* 2018;265:1803-9. doi:10.1007/s00415-018-8915-7
- 151 Věchetová G, Slovák M, Kemlink D, et al. The impact of non-motor symptoms on the health-related quality of life in patients with functional movement disorders. *J Psychosom Res* 2018;115:32-7. doi:10.1016/j.jpsychores.2018.10.001
- 152 Stone J, Sharpe M, Rothwell PM, Warlow CP. The 12 year prognosis of unilateral functional weakness and sensory disturbance. *J Neurol Neurosurg Psychiatry* 2003;74:591-6. doi:10.1136/jnnp.74.5.591
- 153 Ospina JP, Larson AG, Jalilianhasanpour R, et al. Individual differences in social network size linked to nucleus accumbens and hippocampal volumes in functional neurological disorder: A pilot study. *J Affect Disord* 2019;258:50-4. doi:10.1016/j.jad.2019.07.061
- 154 Mac CJ, Trimble MR. 'Hysteria', 'functional' or 'psychogenic'? A survey of British neurologists' preferences. *J R Soc Med* 1991;84:471-5. doi:10.1177/014107689108400808
- 155 Stone J, Wojcik W, Durrance D, et al. What should we say to patients with symptoms unexplained by disease? The "number needed to offend". *BMJ* 2002;325:1449-50. doi:10.1136/bmj.325.7378.1449
- 156 Stone J, Carson A, Hallett M. Explanation as treatment for functional neurologic disorders. *Handb Clin Neurol* 2016;139:543-53. doi:10.1016/B978-0-12-801772-2.00044-8
- 157 Edwards MJ, Stone J, Lang AE. From psychogenic movement disorder to functional movement disorder: it's time to change the name. *Mov Disord* 2014;29:849-52. doi:10.1002/mds.25562
- 158 LaFaver K, Hallett M. Functional or psychogenic: what's the better name? *Mov Disord* 2014;29:1698-9. doi:10.1002/mds.26035
- 159 Ding JM, Kanaan RA. Conversion disorder: A systematic review of current terminology. *Gen Hosp Psychiatry* 2017;45:51-5. doi:10.1016/j.genhosppsych.2016.12.009
- 160 Fahn S, Olanow CW. "Psychogenic movement disorders": they are what they are. *Mov Disord* 2014;29:853-6. doi:10.1002/mds.25899
- 161 Jankovic J. "Psychogenic" versus "functional" movement disorders? That is the question. *Mov Disord* 2014;29:1697-8. doi:10.1002/mds.26040
- 162 Bègue I, Adams C, Stone J, Perez DL. Structural alterations in functional neurological disorder and related conditions: a software and hardware problem? *Neuroimage Clin* 2019;22:101798. doi:10.1016/j.nicl.2019.101798
- 163 Tolchin B, Perez DL, Szaflarski JP, et al. What's in a name? *Epilepsy Behav* 2020;112:107364. doi:10.1016/j.yebeh.2020.107364
- 164 Asadi-Pooya AA, Brigo F, Mildon B, Nicholson TR. Terminology for psychogenic nonepileptic seizures: Making the case for "functional seizures". *Epilepsy Behav* 2020;104(Pt A):106895. doi:10.1016/j.yebeh.2019.106895

- 165 Kerr WT, Stern JM. We need a functioning name for PNES: Consider dissociative seizures. *Epilepsy Behav* 2020;105:107002. doi:10.1016/j.yebeh.2020.107002
- 166 LaFrance WC Jr. Psychogenic nonepileptic "seizures" or "attacks"? It's not just semantics: seizures. *Neurology* 2010;75:87-8. doi:10.1212/WNL.Ob013e3181e62181
- 167 Benbadis SR. Psychogenic nonepileptic "seizures" or "attacks"? It's not just semantics: attacks. *Neurology* 2010;75:84-6. doi:10.1212/WNL.Ob013e3181e6216f
- 168 Ding JM, Kanaan RA. What should we say to patients with unexplained neurological symptoms? How explanation affects offence. *J Psychosom Res* 2016;91:55-60. doi:10.1016/j.jpsychores.2016.10.012
- 169 Stone J, Carson A. 'Organic' and 'non-organic': a tale of two turnips. *Pract Neurol* 2017;17:417-8. doi:10.1136/practneurol-2017-001660
- 170 Shen W, Bowman ES, Markand ON. Presenting the diagnosis of pseudoseizure. *Neurology* 1990;40:756-9. doi:10.1212/WNL.40.5.756
- 171 Hall-Patch L, Brown R, House A, et al, NEST collaborators. Acceptability and effectiveness of a strategy for the communication of the diagnosis of psychogenic nonepileptic seizures. *Epilepsia* 2010;51:70-8. doi:10.1111/j.1528-1167.2009.02099.x
- 172 Carson A, Lehn A, Ludwig L, Stone J. Explaining functional disorders in the neurology clinic: a photo story. *Pract Neurol* 2016;16:56-61. doi:10.1136/practneurol-2015-001242
- 173 Stone J, Carson A. Functional neurologic disorders. *Continuum (Minneapolis)* 2015;21(3 Behavioral Neurology and Neuropsychiatry):818-37. doi:10.1212/01.CON.0000466669.02477.45
- 174 Stone J, Edwards M. Trick or treat? Showing patients with functional (psychogenic) motor symptoms their physical signs. *Neurology* 2012;79:282-4. doi:10.1212/WNL.Ob013e31825dff63
- 175 Perez DL, Hunt A, Sharma N, Flaherty A, Caplan D, Schmahmann JD. Cautionary notes on diagnosing functional neurologic disorder as a neurologist-in-training. *Neurol Clin Pract* 2020;10:484-7. doi:10.1212/CJP.0000000000000779
- 176 Reuber M. Dissociative (non-epileptic) seizures: tackling common challenges after the diagnosis. *Pract Neurol* 2019;19:332-41. doi:10.1136/practneurol-2018-002177
- 177 Rockliffe-Fidler C, Willis M. Explaining dissociative seizures: a neuropsychological perspective. *Pract Neurol* 2019;19:259-63. doi:10.1136/practneurol-2018-002100
- 178 Medina M, Giambareri L, Lazarow SS, et al. Using patient-centered clinical neuroscience to deliver the diagnosis of functional neurological disorder (FND): results from an innovative educational workshop. *Acad Psychiatry* 2021;45:185-9. doi:10.1007/s40596-020-01324-8
- 179 Kanaan RA, Armstrong D, Wessely SC. Neurologists' understanding and management of conversion disorder. *J Neurol Neurosurg Psychiatry* 2011;82:961-6. doi:10.1136/jnnp.2010.233114
- 180 Kanaan RA, Armstrong D, Barnes P, Wessely S. In the psychiatrist's chair: how neurologists understand conversion disorder. *Brain* 2009;132:2889-96. doi:10.1093/brain/awp060
- 181 Kanaan RA, Armstrong D, Wessely S. Limits to truth-telling: neurologists' communication in conversion disorder. *Patient Educ Couns* 2009;77:296-301. doi:10.1016/j.pec.2009.05.021
- 182 Altalib HH, Elzamzamy K, Pugh MJ, et al. Communicating diagnostic certainty of psychogenic nonepileptic seizures - a national study of provider documentation. *Epilepsy Behav* 2016;64(Pt A):4-8. doi:10.1016/j.yebeh.2016.08.032
- 183 LaFaver K, Lang AE, Stone J, et al. Opinions and clinical practices related to diagnosing and managing functional (psychogenic) movement disorders: changes in the last decade. *Eur J Neurol* 2020;27:975-84. doi:10.1111/ene.14200
- 184 Farias ST, Thieman C, Alsaadi TM. Psychogenic nonepileptic seizures: acute change in event frequency after presentation of the diagnosis. *Epilepsy Behav* 2003;4:424-9. doi:10.1016/S1525-5050(03)00143-4
- 185 Duncan R, Horwood J, Razvi S, Mulhern S. Psychogenic nonepileptic seizures that remit when the diagnosis is given: just good luck? *Epilepsy Behav* 2020;102:106667. doi:10.1016/j.yebeh.2019.106667
- 186 Gambini O, Demartini B, Chiesa V, Turner K, Barbieri V, Canevini MP. Long-term outcome of psychogenic nonepileptic seizures: the role of induction by suggestion. *Epilepsy Behav* 2014;41:140-3. doi:10.1016/j.yebeh.2014.09.076
- 187 Mayor R, Brown RJ, Cock H, et al. Short-term outcome of psychogenic non-epileptic seizures after communication of the diagnosis. *Epilepsy Behav* 2012;25:676-81. doi:10.1016/j.yebeh.2012.09.033
- 188 Duncan R, Razvi S, Mulhern S. Newly presenting psychogenic nonepileptic seizures: incidence, population characteristics, and early outcome from a prospective audit of a first seizure clinic. *Epilepsy Behav* 2011;20:308-11. doi:10.1016/j.yebeh.2010.10.022
- 189 Arain AM, Hamadani AM, Islam S, Abou-Khalil BW. Predictors of early seizure remission after diagnosis of psychogenic nonepileptic seizures. *Epilepsy Behav* 2007;11:409-12. doi:10.1016/j.yebeh.2007.07.017
- 190 Nunez-Wallace KR, Murphey DK, Proto D, et al. Health resource utilization among US veterans with psychogenic nonepileptic seizures: A comparison before and after video-EEG monitoring. *Epilepsy Res* 2015;114:114-21. doi:10.1016/j.eplepsyres.2015.05.002
- 191 Salinsky M, Storzbach D, Goy E, Kellogg M, Boudreau E. Health care utilization following diagnosis of psychogenic nonepileptic seizures. *Epilepsy Behav* 2016;60:107-11. doi:10.1016/j.yebeh.2016.04.007
- 192 Brough JL, Moghaddam NG, Gresswell DM, Dawson DL. The impact of receiving a diagnosis of Non-Epileptic Attack Disorder (NEAD): A systematic review. *J Psychosom Res* 2015;79:420-7. doi:10.1016/j.jpsychores.2015.09.009
- 193 Mayor R, Brown RJ, Cock H, et al. A feasibility study of a brief psycho-educational intervention for psychogenic nonepileptic seizures. *Seizure* 2013;22:760-5. doi:10.1016/j.seizure.2013.06.008
- 194 Chen DK, Maheshwari A, Franks R, Trolley GC, Robinson JS, Hrachovy RA. Brief group psychoeducation for psychogenic nonepileptic seizures: a neurologist-initiated program in an epilepsy center. *Epilepsia* 2014;55:156-66. doi:10.1111/epi.12481
- 195 Drane DL, LaRoche SM, Ganesh GA, Teagarden D, Loring DW. A standardized diagnostic approach and ongoing feedback improves outcome in psychogenic nonepileptic seizures. *Epilepsy Behav* 2016;54:34-9. doi:10.1016/j.yebeh.2015.10.026
- 196 Cope SR, Smith JG, Edwards MJ, Holt K, Agrawal N. Enhancing the communication of functional neurological disorder diagnosis: a multidisciplinary education session. *Eur J Neurol* 2021;28:40-7. doi:10.1111/ene.14525
- 197 Gelauff JM, Rosmalen JGM, Carson A, et al. Internet-based self-help randomized trial for motor functional neurologic disorder (SHIFT). *Neurology* 2020;95:e1883-96. doi:10.1212/WNL.0000000000010381
- 198 Nielsen G, Stone J, Matthews A, et al. Physiotherapy for functional motor disorders: a consensus recommendation. *J Neurol Neurosurg Psychiatry* 2015;86:1113-9. doi:10.1136/jnnp-2014-309255
- 199 Jordbru AA, Smedstad LM, Klungsøy O, Martinsen EW. Psychogenic gait disorder: a randomized controlled trial of physical rehabilitation with one-year follow-up. *J Rehabil Med* 2014;46:181-7. doi:10.2340/16501977-1246
- 200 Nielsen G, Buszewicz M, Stevenson F, et al. Randomised feasibility study of physiotherapy for patients with functional motor symptoms. *J Neurol Neurosurg Psychiatry* 2017;88:484-90. doi:10.1136/jnnp-2016-314408
- 201 Nielsen G, Stone J, Edwards MJ. Physiotherapy for functional (psychogenic) motor symptoms: a systematic review. *J Psychosom Res* 2013;75:93-102. doi:10.1016/j.jpsychores.2013.05.006
- 202 Maggio JB, Ospina JP, Callahan J, Hunt AL, Stephen CD, Perez DL. Outpatient physical therapy for functional neurological disorder: a preliminary feasibility and naturalistic outcome study in a US Cohort. *J Neuropsychiatry Clin Neurosci* 2020;32:85-9. doi:10.1176/appi.neuropsych.19030068
- 203 Demartini B, Bombieri F, Goeta D, Gambini O, Ricciardi L, Tinazzi M. A physical therapy programme for functional motor symptoms: A telemedicine pilot study. *Parkinsonism Relat Disord* 2020;76:108-11. doi:10.1016/j.parkrel.2019.05.004
- 204 Maggio J, Alluri PR, Paredes-Echeverri S, et al. Briquet syndrome revisited: implications for functional neurological disorder. *Brain Commun* 2020;2:a156. doi:10.1093/braincomms/fcaa156
- 205 Nielsen G, Stone J, Buszewicz M, et al. Physio4FMD Collaborative Group. Physio4FMD: protocol for a multicentre randomised controlled trial of specialist physiotherapy for functional motor disorder. *BMC Neurol* 2019;19:242. doi:10.1186/s12883-019-1461-9
- 206 Nicholson C, Edwards MJ, Carson AJ, et al. Occupational therapy consensus recommendations for functional neurological disorder. *J Neurol Neurosurg Psychiatry* 2020;91:1037-45. doi:10.1136/jnnp-2019-322281
- 207 Baker J, Barnett C, Cavalli L, et al. Management of functional communication, swallowing, cough and related disorders: consensus recommendations for speech and language therapy. *J Neurol Neurosurg Psychiatry* 2021;92:1112-25. doi:10.1136/jnnp-2021-326767
- 208 Pick S, Anderson DG, Asadi-Pooya AA, et al. Outcome measurement in functional neurological disorder: a systematic review and recommendations. *J Neurol Neurosurg Psychiatry* 2020;91:638-49. doi:10.1136/jnnp-2019-322180
- 209 Nicholson TR, Carson A, Edwards MJ, et al, and the FND-COM (Functional Neurological Disorders Core Outcome Measures) Group. FND-COM group collaborators are as follows. Outcome measures for functional neurological disorder: a review of the theoretical complexities. *J Neuropsychiatry Clin Neurosci* 2020;32:33-42. doi:10.1176/appi.neuropsych.19060128

- 210 Carlson P, Nicholson Perry K. Psychological interventions for psychogenic non-epileptic seizures: A meta-analysis. *Seizure* 2017;45:142-50. doi:10.1016/j.seizure.2016.12.007
- 211 Goldstein LH, Chalder T, Chigwedere C, et al. Cognitive-behavioral therapy for psychogenic nonepileptic seizures: a pilot RCT. *Neurology* 2010;74:1986-94. doi:10.1212/WNL.Ob013e3181e39658
- 212 LaFrance WCJr, Baird GL, Barry JJ, et al, NES Treatment Trial (NEST-T) Consortium. Multicenter pilot treatment trial for psychogenic nonepileptic seizures: a randomized clinical trial. *JAMA Psychiatry* 2014;71:997-1005. doi:10.1001/jamapsychiatry.2014.817
- 213 Reiter J, Andrews D, Reiter C, LaFrance WCJr. *Taking control of your seizures: workbook*. Oxford University Press, 2015. doi:10.1093/med:psych/9780199335015.001.0001
- 214 Tolchin B, Baslet G, Suzuki J, et al. Randomized controlled trial of motivational interviewing for psychogenic nonepileptic seizures. *Epilepsia* 2019;60:986-95. doi:10.1111/epi.14728
- 215 Sharpe M, Walker J, Williams C, et al. Guided self-help for functional (psychogenic) symptoms: a randomized controlled efficacy trial. *Neurology* 2011;77:564-72. doi:10.1212/WNL.Ob013e318228c0c7
- 216 Williams C, Carson A, Smith S, Sharpe M, Cavanagh J, Kent C. *Overcoming functional neurological symptoms: a five area approach*. CRC. Press, 2017. doi:10.1201/b13567
- 217 Dallochio C, Tinazzi M, Bombieri F, Arnó N, Erro R. Cognitive behavioural therapy and adjunctive physical activity for functional movement disorders (conversion disorder): a pilot, single-blinded, randomized study. *Psychother Psychosom* 2016;85:381-3. doi:10.1159/000446660
- 218 Gutkin M, McLean L, Brown R, Kanaan RA. Systematic review of psychotherapy for adults with functional neurological disorder. *J Neurol Neurosurg Psychiatry* 2020;Nov 5:321926.
- 219 Voon V, Lang AE. Antidepressant treatment outcomes of psychogenic movement disorder. *J Clin Psychiatry* 2005;66:1529-34. doi:10.4088/JCP.v66n1206
- 220 LaFrance WCJr, Keitner GI, Papandonatos GD, et al. Pilot pharmacologic randomized controlled trial for psychogenic nonepileptic seizures. *Neurology* 2010;75:1166-73. doi:10.1212/WNL.Ob013e3181f4d5a9
- 221 Pintor L, Baillés E, Matrai S, et al. Efficiency of venlafaxine in patients with psychogenic nonepileptic seizures and anxiety and/or depressive disorders. *J Neuropsychiatry Clin Neurosci* 2010;22:401-8. doi:10.1176/jnp.2010.22.4.401
- 222 Bravo TP, Hoffman-Snyder CR, Wellik KE, et al. The effect of selective serotonin reuptake inhibitors on the frequency of psychogenic nonepileptic seizures: a critically appraised topic. *Neurologist* 2013;19:30-3. doi:10.1097/NRL.Ob013e31827c6bfd
- 223 Baslet G, Ehler A, Oser M, Dworetzky BA. Mindfulness-based therapy for psychogenic nonepileptic seizures. *Epilepsy Behav* 2020;103(Pt A):106534. doi:10.1016/j.yebeh.2019.106534
- 224 Myers L, Vaidya-Mathur U, Lancman M. Prolonged exposure therapy for the treatment of patients diagnosed with psychogenic nonepileptic seizures (PNES) and post-traumatic stress disorder (PTSD). *Epilepsy Behav* 2017;66:86-92. doi:10.1016/j.yebeh.2016.10.019
- 225 Kompoliti K, Wilson B, Stebbins G, Bernard B, Hinson V. Immediate vs. delayed treatment of psychogenic movement disorders with short term psychodynamic psychotherapy: randomized clinical trial. *Parkinsonism Relat Disord* 2014;20:60-3. doi:10.1016/j.parkreldis.2013.09.018
- 226 Bullock KD, Mirza N, Forte C, Trockel M. Group dialectical-behavior therapy skills training for conversion disorder with seizures. *J Neuropsychiatry Clin Neurosci* 2015;27:240-3. doi:10.1176/appi.neuropsych.13120359
- 227 Barry JJ, Wittenberg D, Bullock KD, Michaels JB, Classen CC, Fisher RS. Group therapy for patients with psychogenic nonepileptic seizures: a pilot study. *Epilepsy Behav* 2008;13:624-9. doi:10.1016/j.yebeh.2008.06.013
- 228 Conwill M, Oakley L, Evans K, Cavanna AE. CBT-based group therapy intervention for nonepileptic attacks and other functional neurological symptoms: a pilot study. *Epilepsy Behav* 2014;34:68-72. doi:10.1016/j.yebeh.2014.03.012
- 229 Moene FC, Spinhoven P, Hoogduin KA, van Dyck R. A randomised controlled trial on the additional effect of hypnosis in a comprehensive treatment programme for in-patients with conversion disorder of the motor type. *Psychother Psychosom* 2002;71:66-76. doi:10.1159/000049348
- 230 Moene FC, Spinhoven P, Hoogduin KA, van Dyck R. A randomized controlled clinical trial of a hypnosis-based treatment for patients with conversion disorder, motor type. *Int J Clin Exp Hypn* 2003;51:29-50. doi:10.1076/1051.1.29.14067
- 231 LaFrance WCJr, Ho WLN, Bhatla A, Baird GL, Altalib HH, Godleski L. Treatment of psychogenic nonepileptic seizures (PNES) using video telehealth. *Epilepsia* 2020;61:2572-82. doi:10.1111/epi.16689
- 232 Perez DL. The CODES trial for dissociative seizures: a landmark study and inflection point. *Lancet Psychiatry* 2020;7:464-5. doi:10.1016/S2215-0366(20)30143-7
- 233 Vizcarra JA, Lopez-Castellanos JR, Dwivedi AK, Schmerler DA, Ries S, Espay AJ. OnabotulinumtoxinA and cognitive behavioral therapy in functional dystonia: A pilot randomized clinical trial. *Parkinsonism Relat Disord* 2019;63:174-8. doi:10.1016/j.parkreldis.2019.02.009
- 234 Dreissen YEM, Dijk JM, Gelauff JM, et al. Botulinum neurotoxin treatment in jerky and tremulous functional movement disorders: a double-blind, randomised placebo-controlled trial with an open-label extension. *J Neurol Neurosurg Psychiatry* 2019;90:1244-50. doi:10.1136/jnnp-2018-320071
- 235 Nicholson TRJ, Voon V. Transcranial magnetic stimulation and sedation as treatment for functional neurologic disorders. *Handb Clin Neurol* 2016;139:619-29. doi:10.1016/B978-0-12-801772-2.00050-3
- 236 Taib S, Ory-Magne F, Brefel-Courbon C, et al. Repetitive transcranial magnetic stimulation for functional tremor: A randomized, double-blind, controlled study. *Mov Disord* 2019;34:1210-9. doi:10.1002/mds.27727
- 237 Garcin B, Mesrati F, Hubsch C, et al. Impact of transcranial magnetic stimulation on functional movement disorders: cortical modulation or a behavioral effect? *Front Neurol* 2017;8:338. doi:10.3389/fneur.2017.00338
- 238 Pick S, Hodsoll J, Stanton B, et al. Trial Of Neurostimulation In Conversion Symptoms (TONICS): a feasibility randomised controlled trial of transcranial magnetic stimulation for functional limb weakness. *BMJ Open* 2020;10:e037198.
- 239 Stone J, Hoeritzauer I, Brown K, Carson A. Therapeutic sedation for functional (psychogenic) neurological symptoms. *J Psychosom Res* 2014;76:165-8. doi:10.1016/j.jpsychores.2013.10.003
- 240 Deeley Q. Hypnosis as therapy for functional neurological disorders. *Handb Clin Neurol* 2016;139:585-95. doi:10.1016/B978-0-12-801772-2.00047-3
- 241 Bullock K, Won AS, Bailenson J, Friedman R. Virtual reality-delivered mirror visual feedback and exposure therapy for FND: a midpoint report of a randomized controlled feasibility study. *J Neuropsychiatry Clin Neurosci* 2020;32:90-4. doi:10.1176/appi.neuropsych.19030071
- 242 Edwards MJ, Bhatia KP, Cordivari C. Immediate response to botulinum toxin injections in patients with fixed dystonia. *Mov Disord* 2011;26:917-8. doi:10.1002/mds.23562
- 243 Burke MJ, Faria V, Cappon D, Pascual-Leone A, Kaptchuk TJ, Santarnecchi E. Leveraging the shared neurobiology of placebo effects and functional neurological disorder: a call for research. *J Neuropsychiatry Clin Neurosci* 2020;32:101-4. doi:10.1176/appi.neuropsych.19030077
- 244 Ludwig LPJ, Nicholson T, Aybek S, et al. Life events, stress and psychological trauma in conversion (functional neurological) disorder: systematic review and meta-analysis of case-control studies. *Lancet Psychiatry* 2018;5:307-20. doi:10.1016/S2215-0366(18)30051-8
- 245 Selkirk M, Duncan R, Oto M, Pelosi A. Clinical differences between patients with nonepileptic seizures who report antecedent sexual abuse and those who do not. *Epilepsia* 2008;49:1446-50. doi:10.1111/j.1528-1167.2008.01611.x

Supplementary file 1: Search terms

Supplementary file 2: Video 1 How to explain the diagnosis of FND. Key elements that can be discussed with patients when delivering a diagnosis of FND. Video made using Doodly software version 2.6.13

Supplementary file 3: Video 2 Triaging initial treatment recommendations in patients with FND. Key elements that warrant consideration when providing initial treatment recommendations to patients with FND. Video made using Doodly software version 2.6.13